

CLINICAL STUDIES OF THE BLOOD VOLUME. V. HYPERTHYROIDISM AND MYXEDEMA

By JOHN G. GIBSON, 2d, AND ALFRED W. HARRIS

(From the Department of Medicine, Harvard Medical School and the Medical Clinic of the Peter Bent Brigham Hospital, Boston)

(Received for publication September 7, 1938)

In a study of the blood volume in normal human beings (1) a relationship between the total circulating blood volume and the basal metabolic rate was observed. In a group of 99 individuals the distribution of cases above and below the average total blood volume was similar to the distribution of basal metabolic rates above and below average normal values, and the decline in total blood volume with advancing age was parallel to the decline in basal metabolic rate. An increase in total blood volume, the degree of which was related to the severity of the condition, was found in patients with congestive heart failure (2) in which it may be assumed that the oxygen carrying mechanism was inefficient in meeting the tissue oxygen requirement. This observation suggested that the level of oxygen consumption might exert a considerable effect on the circulating blood volume, and prompted us to undertake a study of the blood volume in hyperthyroidism and myxedema in which the metabolism is severely disturbed.

Previous studies have indicated that hyperthyroidism is characterized by an abnormally high (3, 4, 5) and myxedema by an abnormally low circulating blood volume (6, 7, 8). Chang (4), using the CO method, found no direct relationship between the degree of increase in basal metabolic rate and increase in blood volume but observed decreases in blood volume with clinical recovery in all his cases. Goldbloom and Libin (5), using a dye method, found the increases above their normal values in hyperthyroidism great enough to lead them to believe that determination of blood volume was of value in the differential diagnosis of hyperthyroidism. Thompson (6) found the plasma volume about 30 per cent below normal in untreated cases of myxedema. On treatment, with the return of the basal metabolic rate to normal, the plasma volume rose to within normal limits.

MATERIAL AND METHODS

Twenty females and 5 males with clinically proven hyperthyroidism were studied. The basal metabolic rate was more than 15 per cent above normal in all cases. In 5 males and 10 females the changes in plasma and total blood volume were followed during the course of therapy, and determinations were made in all of these cases at intervals varying from 5 to 22 days after subtotal thyroidectomy was performed. In 2 males and 4 females blood volume was determined pre-operatively after the administration of Lugol's solution.

Single blood volume determinations were made in 5 females and 2 males with clinically proven myxedema. The basal metabolic rate was more than 15 per cent below normal in all cases. One man with pernicious anemia developed myxedema during a relapse, and another man developed myxedema some time after subtotal thyroidectomy.

Basal metabolic rates were determined by the standard technique of Benedict and Roth (9); plasma and total blood volume were determined by the direct method of Gibson and Evans (10); venous pressure was determined by the direct method of Evans (11) and circulation time by the decholin method of Winternitz, Deutsch, and Brüll (12).

RESULTS

Surface area rather than height was taken as the basis for prediction of normal plasma and total blood volume, inasmuch as the basal metabolic rates were based upon surface area. In almost every case the clinical history revealed the loss of some weight prior to admission to the hospital, and it is possible that the predicted volumes based upon surface area may represent values that were too high in some instances. The actual total blood volumes determined in these cases may well represent, therefore, greater percentage deviations from the normal volumes than indicated.

The course of the basal metabolism, plasma and total blood volume, hematocrit, venous pressure, and circulation time in 25 cases of hyperthyroidism and in 7 cases of myxedema is shown in Table I.

TABLE I

Absolute plasma, cell, and total blood volume; normal total blood volume predicted from surface area, venous pressure, and circulation time in 25 cases of hyperthyroidism and 7 cases of myxedema

Case number	Date	Sex	Age	Height	Weight	Surface area	Predicted total blood volume	Venous pressure	Circulation time	Basal metabolic rate	Blood volume			Hematocrit	Percentage deviation from predicted total blood volume		Remarks
											Plasma	Cell	Total blood		+	-	
			years	cm.	kgm.	square meters	cc.	mm. H ₂ O	seconds	per cent	cc.	cc.	cc.	per cent of cells			
25 PATIENTS WITH HYPERTHYROIDISM																	
9	December 10, 1934	F	40	160.0	55.0	1.55	4025	90	8	+27	2380	1970	4350	45.3	8.1		
12	December 13, 1934	F	23	165.6	66.0	1.71	4175	80	11	+55	3105	1665	4770	36.1	14.3		
18	April 8, 1935	F	42	156.8	51.7	1.49	3850	80	10	+51	2350	1650	4000	41.6	3.9		
23A	May 4, 1935	F	41	152.5	56.0	1.50	3900	75	10	+61	2335	1575	3910	40.7	0.3		
23B	May 24, 1935			152.5	56.4	1.515	3925	90	10	+3	2270	1200	3470	36.1	11.6	8 days postoperatively	
28A	May 17, 1935	F	34	167.5	54.5	1.60	4100	70	9	+47	2740	1610	4350	37.4	6.1		
28B	June 15, 1935			167.5	52.4	1.58	4075	65	10	+6	2680	1345	4025	34.0	1.2	10 days postoperatively	
31A	May 22, 1935	M	22	164.0	43.2	1.40	3450	55	11	+70	2490	1900	4380	42.4	27.0		
31B	June 12, 1935			164.0	44.6	1.44	3725	105	11	+7	2250	1590	3940	41.8	3.1	8 days postoperatively	
60A	July 8, 1935	F	38	163.8	70.0	1.75	4175	60	10	+46	2665	1765	4430	40.5	6.1		
60B	July 25, 1935			163.8	64.6	1.69	4175	55	11	-16	2240	1540	3780	40.8	8.8	9 days postoperatively	
40	June 17, 1935	F	28	158.8	48.4	1.47	3800	100	8	+66	1900	1760	3660	38.2	3.7		
64A	July 13, 1935	F	48	157.4	52.3	1.50	3900	65	12	+58	2470	2160	4630	37.5	18.7		
84A	September 20, 1935	F	42	156.4	62.4	1.61	4100	60	12	+85	3090	2000	5090	39.3	24.1		
84B	September 30, 1935			156.4	60.0	1.58	4075	65	12	+31(?)	2535	1675	4210	39.9	3.3	Iodine administration	
85A	October 1, 1935	F	28	171.5	50.6	1.58	4075	65	11	+30	2270	1600	3870	41.5	5.0		
85B	October 16, 1935			171.5	50.0	1.57	4050	65	12	+11	1990	1330	3320	41.2	17.9	7 days postoperatively	
179	May 12, 1936	F	37	162.6	52.0	1.53	3950	30	9	+20	2265	1365	3630	39.8	8.1		
233A	September 11, 1936	M	36	174.5	65.3	1.78	5250	75	14	+24	2600	2220	4820	46.0	8.2		
233B	September 23, 1936			174.5	65.5	1.79	5275	45	14	+14	2980	2380	5360	44.4	1.6	Iodine administration	
233C	October 16, 1936			174.5	66.8	1.79	5300	20	18	+3	2900	2225	5125	43.4	3.3	22 days postoperatively	
232A	September 11, 1936	F	28	152.5	47.6	1.41	3525	90	12	+38	2075	1435	3510	40.7	0.4		
232B	September 23, 1936			152.5	55.8	1.51	3925	80	12	+14	2150	1320	3470	37.9	9.2	Iodine administration	
232C	October 16, 1936			152.5	54.1	1.49	3875	100	17	-19	2180	1135	3315	34.3	11.5	21 days postoperatively	
241A	October 22, 1936	M	39	176.5	63.7	1.78	5250	40	22	+17	2955	2495	5450	45.8	3.8		
241B	November 2, 1936			176.5	66.3	1.81	5350	40	22	-4	2530	2000	4530	44.3	15.3	Iodine administration	
338A	March 1, 1938	F	44	172.7	76.4	1.89	4175	75	19	+23	2600	2160	4760	45.5	14.0		
340	March 21, 1938	F	63	166.3	74.4	1.81	4175	18	18	+46	2900	2040	4940	41.1	18.3		
344	March 30, 1938	F	39	171.5	86.9	1.97	4175	14	14	+30	2810	2060	4870	42.3	16.6		
345A	April 1, 1938	F	39	162.5	60.8	1.63	4125	10	14	+48	2350	1650	4000	41.2	8.0		
345B	April 20, 1938			162.5	59.3	1.62	4100	10	13(?)	+28	2000	1530	3530	43.4	13.0	Iodine administration	
346A	April 12, 1938	F	52	158.8	57.0	1.57	4025	8	13	+47	2180	1580	3760	41.9	6.5		
346B	April 29, 1938			158.8	55.4	1.55	4000	10	13	+1	2220	1440	3660	39.3	9.1	7 days postoperatively	
348A	April 26, 1938	F	38	149.5	49.8	1.42	3650	10	14	+47	2180	1900	4080	46.5	11.8		
348B	May 10, 1938			149.5	46.9	1.37	3425	10	14	+28	1900	1280	3180	40.4	7.1	11 days postoperatively	
350A	May 16, 1938	M	62	176.0	84.2	1.99	6000	15	15	+49	4330	2800	7130	39.3	18.9		
350C	June 11, 1938			176.0	82.2	1.98	5975	24	24	-2	4100	2380	6480	36.7	8.5	5 days postoperatively	
351A	May 19, 1938	F	55	148.8	59.6	1.53	3950	16	16	+60	2440	1610	4050	39.6	2.5		
351B	June 10, 1938			148.8	56.0	1.48	3825	17	17	+26	1720	1470	3190	46.1	16.6	9 days postoperatively	
352A	May 26, 1938	M	50	175.0	68.6	1.82	5400	12	12	+53	3250	2690	5940	45.3	10.0		
352B	June 22, 1938			175.0	69.4	1.83	5450	16	16	+6	3000	2170	5170	42.2	5.1	9 days postoperatively	
356	June 6, 1938	F	35	160.0	58.6	1.59	4075	12	12	+46	2150	1580	3730	42.3	8.5		
7 PATIENTS WITH MYXEDEMA																	
10	December 11, 1934	F	53	165.5	72.1	1.78	4150	90	23	-15	3070	1170	4170	26.4	0.5		
29	May 18, 1935	F	62	161.5	63.6	1.66	4150	65	45	-17	2030	1530	3560	43.0	14.2		
234A	September 22, 1936	F	45	172.5	53.3	1.61	4100	95	26	-43	2490	1210	3700	32.7	9.8		
234B	October 13, 1936			172.5	54.3	1.63	4125	100	20	-20	2485	1235	3720	33.2	8.8	On thyroid medication	
239	October 20, 1936	F	42	158.8	66.6	1.68	4150	60	23	-18	2120	1400	3520	39.8	15.2		
241C	December 11, 1936	M	39	176.5	76.0	1.91	5725	45	35	-21	2680	2020	4700	43.0	17.9	Post-total thyroidectomy	
271D	April 1, 1937	M	72	177.5	75.4	1.92	5775	90	22	-23	2730	2240	4970	45.0	13.9	Pericious anemia	
353A	May 26, 1938	F	59	163.8	69.2	1.66	4150	50	22	-33	1960	1290	3150	40.8	24.1		
353B	July 7, 1938			163.8	65.7	1.62	4125	10	22	-12	2450	1410	3860	36.5	8.9	On thyroid medication	

In Figure 1 is shown the distribution of cases of hyperthyroidism and myxedema above and below normal total blood volume as predicted on the basis of surface area. It is evident that in comparison with the curve of distribution of total blood volumes in normals, the cases with hyperthyroidism form a group with the volumes definitely increased above normal, and that the cases

with myxedema form a group with definitely subnormal volumes. Thus in the 25 cases of hyperthyroidism 17 cases had a total blood volume which was within plus or minus 10 per cent of predicted normal value, 8 were above 10 per cent above normal, while none were less than 10 per cent below normal. In the group of patients with myxedema only one case had a volume in




FIG. 1. THE DISTRIBUTION OF THE PERCENTAGE OF CASES ABOVE AND BELOW PREDICTED NORMAL TOTAL BLOOD VOLUME IN 25 CASES OF HYPERTHYROIDISM AND 7 CASES OF MYXEDEMA COMPARED TO THE DISTRIBUTION OF THE PERCENTAGE OF CASES ABOVE AND BELOW AVERAGE NORMAL TOTAL BLOOD VOLUME IN 99 NORMAL HUMANS

excess of predicted normal value, only 2 cases fell within limits of plus or minus 10 per cent of normal, and 5 cases had volumes more than 10 per cent below normal.

The percentage deviation from predicted normal total blood volume based on surface area is shown in relation to basal metabolic rate in Figure 2. A direct relationship between deviation from predicted normal total blood volume and the levels of basal metabolic rate was observed. Prior to treatment, the circulation time was below normal in all except one case of hyperthyroidism and above normal in all the cases of myxoedema. Venous pressures were within the limits of normality of the method employed in all cases in which the determination was made. The direct relationship between the speed of blood flow and

basal metabolic rate reported by Blumgart, Gargill, and Gilligan (13) in hyperthyroidism was confirmed.

No significant relationship between the percentage deviation from predicted normal volume based on surface area and circulation time or venous pressure was observed.

The change in the absolute total blood volume in 5 males and 10 females in relation to change in basal metabolic rate occurring during the course of therapy is shown in Figure 3. In all cases the basal metabolic rate was lower after operation, and in all except 1 case the total blood volume was less postoperatively than it was at the initial determination. Reduction in total blood volume postoperatively ranged from 540 to 770 cc., or an average of 650 cc. in males, and from 100 to 900




FIG. 2. PERCENTAGE DEVIATION FROM NORMAL TOTAL BLOOD VOLUME AS PREDICTED ON THE BASIS OF SURFACE AREA IN 25 CASES OF HYPERTHYROIDISM AND 7 CASES OF MYXEDEMA IN RELATION TO BASAL METABOLIC RATE

cc. or an average of 500 cc. in females. In 1 male and 3 females the total blood volume decreased during the course of iodine administration. The total blood volume determined postoperatively was higher than that following iodine administration in 1 male and lower in 1 male and 1 female.

In 2 cases of myxedema, following the increase in basal metabolic rate resulting from thyroid therapy the total blood volume was unchanged in one and definitely increased in the other.

DISCUSSION

Our findings are somewhat at variance with those reported in the literature. Chang (4), using the CO method, found an average increase over his accepted normal value for total blood

volume of about 17 per cent in 21 cases of hyperthyroidism. Goldbloom and Libin (5), employing a modification of the dye method of Seydhelm and Lampe, found the total blood volume increased by 50 per cent over normal in 9 cases with hyperthyroidism. The average increase over normal in our series of 25 cases was only 6 per cent. Thompson (6) found the total blood volume to be about 25 per cent below normal in 9 cases of myxedema. The average in our group of 7 myxedematous patients was 15 per cent below normal.

In our opinion certain errors inherent in the earlier dye techniques, principally those arising from differences in the time required for the dye to become completely mixed in the blood stream




FIG. 3. DECREASE IN TOTAL BLOOD VOLUME IN HYPERTHYROIDISM WITH DECREASE IN BASAL METABOLIC RATE FOLLOWING THERAPY

Dotted lines indicate the change taking place under treatment with Lugol's solution only; solid lines indicate the change following subtotal thyroidectomy.

in individuals with extremely rapid or slow circulation times (10), may be responsible for the discrepancies in results obtained by the above authors and those obtained in this study. The higher values found by Chang (4) may be attributed to the fact that the CO method probably measures myohemoglobin in addition to circulating oxyhemoglobin.

In both groups of patients in this series there were considerable variations in individual cases from the trend exhibited by the group as a whole, and therefore the determination of the blood volume has little, if any, practical value in the differential diagnosis of either hyperthyroidism or myxedema. Several factors account for these variations. In the determination of basal metabolic rates, oxygen consumption can be determined only within rather wide limits. It is common knowledge that errors are more prone to be

in the direction of too high rather than too low rates, and that the technical difficulties of accurate determinations increase with the severity of the clinical condition of the patient. In addition, as stated above, standards of normality are at best arbitrary values when applied to individuals and while the conclusions drawn from a sufficiently large group of cases may be valid, the findings in an individual case may not be in keeping with average trends.

Further evidence that in hyperthyroidism total blood volume determined at the height of the rise in metabolism is increased over the level of the individual in a normal metabolic state may be adduced from the fact that in all cases successful treatment of the disorder, as evidenced by a reduction in the metabolic rate, was accompanied by a prompt and considerable fall in the total blood volume. This reduction was greater than could

be accounted for by rest, or by the minor hemorrhage incident to operation. The reduction occurring coincident with the lowering of the metabolic rate was shared equally by plasma and red cell volume. The percentage of reduction in red cell volume occurring after therapy bears a linear relationship to the fall in metabolic rate as is shown in Figure 4.

In our opinion the total blood volume is definitely related to the oxygen requirement as expressed by the basal metabolic determination. Other than an increase above normal requirements in the total number of circulating red cells in hyperthyroidism and a decrease in myxedema, no explanation of the mechanism of the volume

change characteristic of these diseases has come to light.

None of these cases of hyperthyroidism had any evidence of valvular diseases or hypertensive heart disease nor did they exhibit any of the physical signs of congestive heart failure. The greatest increase in total blood volume above normal in this series was 28 per cent and the average about 6 per cent. In frank congestive failure the average increase in total blood volume above normal is 22 per cent (2). It would appear that even in the presence of an increased cardiac burden, the mechanical disadvantage imposed by the degree of hypervolemia experienced by these thyrotoxic patients was not enough to




FIG. 4. PERCENTAGE REDUCTION FROM INITIAL CIRCULATING RED CELL VOLUME OCCURRING AFTER THERAPY IN HYPERTHYROIDISM IN RELATION TO THE ABSOLUTE DECREASE IN BASAL METABOLIC RATE

precipitate congestive heart failure. However, in the absence of known specific pathological changes in the thyroid heart (14, 15), the hypervolemia brought about by the increased oxygen requirement of this diseased state may offer some explanation as to the mechanism of congestive heart failure in cases of hyperthyroidism.

CONCLUSIONS

(1) In 25 cases of hyperthyroidism the total blood volume was increased above normal on an average of 5.45 per cent; and 15.5 per cent below normal in 7 cases of myxedema.

(2) The deviation from normal in the untreated hyperthyroid state bears a linear relationship to the oxygen requirement as measured by the determination of the basal metabolic rate.

(3) In hyperthyroidism successful treatment is accompanied by a decrease in red cell and total blood volume commensurate with the lowering in basal metabolic rate.

We wish to express our thanks to Drs. H. L. Blumgart and M. D. Altschule of the Beth Israel Hospital, Boston, for permission to utilize clinical material. Miss Evelyn Berstein gave valuable technical assistance.

BIBLIOGRAPHY

- Gibson, J. G., 2d, and Evans, Wm. A., Jr., Clinical studies of the blood volume. II. The relation of plasma and total blood volume to venous pressure, blood velocity rate, physical measurements, age and sex in ninety normal humans. *J. Clin. Invest.*, 1937, **16**, 317.
- Gibson, J. G., 2d, and Evans, Wm. A., Jr., Clinical studies of the blood volume. III. Changes in blood volume, venous pressure and blood velocity rate in chronic congestive heart failure. *J. Clin. Invest.*, 1937, **16**, 851.
- Rowntree, L. G., and Brown, G. E., *The Volume of the Blood and Plasma in Health and Disease*, W. B. Saunders Co., Philadelphia, 1929.
- Chang, H., The blood volume in hyperthyroidism. *J. Clin. Invest.*, 1931, **10**, 475.
- Goldbloom, A. A., and Libin, I., Clinical studies in circulatory adjustments. I. Clinical evaluation of studies of circulating blood volume. *Arch. Int. Med.*, 1935, **55**, 484.
- Thompson, W. O., Studies in blood volume. I. The blood volume in myxedema with a comparison of plasma volume changes in myxedema and cardiac edema. *J. Clin. Invest.*, 1926, **2**, 477.
- Blumgart, H. L., Gargill, S. L., and Gilligan, D. R., Studies on velocity of blood flow. XIV. The circulation in myxedema with a comparison of the velocity of blood flow in myxedema and thyrotoxicosis. *J. Clin. Invest.*, 1931, **9**, 91.
- Holböll, S. A., Über die Grösse der blutmenge bei patienten mit myxödem. Studien der blutmenge. *Acta med. Scandinav.*, 1930, **73**, 538.
- Roth, P., Modifications of apparatus and improved technique adaptable to the Benedict type of respiration apparatus. *Boston M. and S. J.*, 1922, **186**, 457.
- Gibson, J. G., 2d, and Evans, Wm. A., Jr., Clinical studies of the blood volume. I. Clinical application of a method employing the azo dye "Evans Blue" and the spectrophotometer. *J. Clin. Invest.*, 1937, **16**, 301.
- Evans, Wm. A., Jr., Venous pressure. *New England J. Med.*, 1932, **207**, 1934.
- Winternitz, M., Deutsch, J., and Brüll, Z., Eine Klinisch brauchbare Bestimmungsmethode der Blutumlaufzeit mittels Decholininjektion. *Med. Klin.*, 1931, **27**, 986.
- Blumgart, H. L., Gargill, S. L., and Gilligan, D. R., Studies on velocity of blood flow. XIII. The circulatory response to thyrotoxicosis. *J. Clin. Invest.*, 1930, **9**, 69.
- Rake, G., and McEachern, D., A study of the heart in hyperthyroidism. *Am. Heart J.*, 1932, **8**, 19.
- Weller, C. V., Wanstrom, R. C., Gordon, H., and Bugher, J. C., Cardiac histopathology in thyroid disease. *Am. Heart J.*, 1932, **8**, 8.