

PRESERVATION OF NORMAL HUMAN PLASMA IN THE LIQUID STATE. IV. STUDIES ON ISOHEMAGGLUTININ TITERS

Eugene L. Lozner, Lloyd R. Newhouser

J Clin Invest. 1944;23(3):361-363. <https://doi.org/10.1172/JCI101503>.

Research Article

Find the latest version:

<https://jci.me/101503/pdf>



PRESERVATION OF NORMAL HUMAN PLASMA IN THE LIQUID STATE. IV. STUDIES ON ISOHEMAGGLUTININ TITERS¹

BY EUGENE L. LOZNER² AND LLOYD R. NEWHOUSER³

(From the Naval Medical Research Institute and the Naval Medical School,
National Naval Medical Center, Bethesda, Maryland)

(Received for publication November 29, 1943)

The presence of isohemagglutinins in plasma has prompted at least two observers to attribute untoward reactions following the administration of *pooled* plasma to this factor (1, 2). While it is conceivable that in plasma from a *single* donor, the same hazard may theoretically exist as in blood from a "dangerous universal donor," it is nevertheless a fact, as has been pointed out by Thalhimer (3), that for purposes of bacteriologic control, the preparation of plasma on a large scale invariably involves pooling. It is the purpose of this communication therefore to report isohemagglutinin titrations on 1000 pools of plasma as compared with 100 monovalent controls and to present a preliminary investigation of the clinical significance of the isoagglutinin titers of pooled plasma.

METHODS

One thousand specimens of pools of plasma, prepared by the Blood Plasma Department of the U. S. Naval Medical School, were analyzed for their titer of isohemagglutinin against "A" and "B" red cell suspensions. The method of plasma preparation has been described previously (4). The method of titration was that in which 0.5 ml. of plasma was diluted geometrically with 0.9 per cent sodium chloride in a rack of 10 × 75 mm. test tubes. To each dilution is added 0.5 ml. of a 1 per cent washed red cell suspension from a panel of "A" and "B" individuals. The tubes are allowed to stand without agitation for 2 hours at 37° C. Following this, each tube is gently agitated and the end-point is read as the last tube in which macroscopic agglutinates are present. The first tube, *i.e.*, that in which 0.5 ml. undiluted plasma has been added to 0.5 ml. red cell suspension, is called "1 : 2" inasmuch as a 50 per cent dilution of the plasma is present. All titrations were done in duplicate and each time a series of titrations was run,

¹ This article has been released for publication by the Division of Publications of the Bureau of Medicine and Surgery of the U. S. Navy. The opinions are those of the writers and are not to be considered as reflecting the policies of the Navy Department.

² Lieutenant, Medical Corps, United States Naval Reserve.

³ Captain, Medical Corps, United States Navy.

controls consisting of plasma from a known "A" and a known "B" individual were also run. In all, 100 "A" and 100 "B" monovalent control titrations were done. The pooled plasma varied with respect to age of preservation in the liquid state and to the number of donors of the pool.

In connection with the statistical study of blood plasma administration presented in the first paper of this series (5), it was possible to compare the distribution of agglutinin titers in those lots of plasma which had been reported as being followed by untoward reactions with a sample of those lots of plasma which had been reported as not followed by reactions. It was also possible to determine the reaction percentage in the administration of those lots of plasma which turned out to have agglutinin titers over "1 : 16."

RESULTS

The distribution of isohemagglutinin titers in 1000 pools of plasma and 100 monovalent controls and the correlations with the age of plasma and with the number of donors contributing to the pools are presented in the accompanying tables (Tables I and II).

It will be observed that pooling alone reduces the percentage of the group with titers of "1 : 16" and over from 72 per cent in the controls to 7.7 per cent in the pools, with respect to anti-A agglutinin, and from 78 per cent in the controls to 15.2 per cent in the pools, with respect to anti-B agglutinin.

The difference between pools with only 5 to 8 donors contributing to them and those with 9 to 12 donors is striking. Whereas the former group had percentages of titers of "1 : 16" and over, of 8.1 per cent and 15.7 per cent of anti-A and anti-B agglutinin, respectively, the latter had percentages of 1.5 per cent and 8.7 per cent.

It is also evident that aging the plasma in the liquid state over 4 months also decreases considerably the percentage of the group with titers of "1 : 16" and over. In 145 pools over 4 months old, none had a titer of "1 : 16" or over of anti-A agglutinin and 7 had titers of "1 : 16" to "1 : 64," but none over "1 : 64," of anti-B

TABLE II
Analysis of anti-B agglutinin titrations on 1000 pools of plasma

Groups		0	1 : 2	1 : 4	1 : 8	1 : 16	1 : 32	1 : 64	1 : 128	1 : 256	1 : 512	Total
Monovalent controls	Number	0	0	4	18	32	25	10	5	4	2	100
	Per cent	0	0	4	18	78 per cent						100 per cent
Total pools	Number	329	187	198	134	92	42	13	2	2	1	1000
	Per cent	32.9	18.7	19.8	13.4	15.2 per cent						100 per cent
Pools with 5 to 8 donors	Number	299	171	185	130	90	40	11	2	2	1	931
	Per cent	32.1	18.4	19.9	14	15.7 per cent						100 per cent
Pools with 9 to 12 donors	Number	30	16	13	4	2	2	2	0	0	0	69
	Per cent	43.5	23.2	18.8	5.8	8.7 per cent						100 per cent
Pools under 1 month old	Number	148	113	130	93	59	27	10	2	2	1	585
	Per cent	25.3	19.3	22.2	15.9	17.3 per cent						100 per cent
Pools 1 to 4 months old (Liquid)	Number	92	52	50	32	28	14	2	0	0	0	270
	Per cent	34.1	19.3	18.5	11.8	16.3 per cent						100 per cent
Pools over 4 months old (Liquid)	Number	89	22	18	9	5	1	1	0	0	0	145
	Per cent	61.4	15.2	12.4	6.2	4.8 per cent						100 per cent

demonstrated to be harmful and it may be concluded therefore that preliminary cross-matching of *pooled* plasma does not appear to be justified experimentally.

SUMMARY AND CONCLUSIONS

1. An analysis of isohemagglutinin titrations on 1000 pools of plasma and 100 monovalent controls is presented.

2. Pooling the plasma reduced the percentage with titers of "1 : 16" and over, of anti-A isohemagglutinin from 72 per cent in the controls to 7.7 per cent in the pools, and of anti-B isohemagglutinin from 78 per cent in the controls to 15.2 per cent.

3. Pools with 9 to 12 donors have a lower percentage of titers of isohemagglutinins of "1 : 16" and over than pools with 5 to 8 donors.

4. Pools over 4 months old have a much lower percentage of titers over "1 : 16" than pools under 4 months old.

5. To date, 15 administrations of plasma with titers over "1 : 16" have been performed without untoward reaction. There was also no

significant difference in titers of isohemagglutinin in lots of plasma reported as being followed by untoward reactions, compared with lots of plasma which were reported as not followed by reactions.

6. It may be concluded that thus far there is no evidence for any harmful effect from the titer of isohemagglutinins present in *pooled* plasma, and, consequently, no justification for preliminary cross-matching of such plasma prior to administration.

BIBLIOGRAPHY

- Polayes, S. H., and Squillace, J. A., Near fatal reaction to transfusion with dried human plasma solution. *J.A.M.A.*, 1942, 118, 1050.
- Downs, T. McK., A reaction to dried pooled human plasma. *U. S. Nav. M. Bull.*, 1943, 41, 189.
- Thalhimer, W., Intravenous injection of pooled normal plasma or serum: Is it dangerous? *J.A.M.A.*, 1942, 120, 1263.
- Newhouser, L. R., and Kendrick, D. B., *Human Plasma Manual*. U. S. Naval Medical School, Washington, D. C., 1941.
- Lozner, E. L., and Newhouser, L. R., Preservation of normal human plasma in the liquid state. I. A statistical study of 1751 administrations. *J. Clin. Invest.*, 1944, 23, 343.