

**DEPRESSION OF PHAGOCYTOSIS BY SOLUTES IN
CONCENTRATIONS FOUND IN THE KIDNEY AND URINE**

Irwin Chernew, A. I. Braude

J Clin Invest. 1962;41(10):1945-1953. <https://doi.org/10.1172/JCI104652>.

Research Article

Find the latest version:

<https://jci.me/104652/pdf>



DEPRESSION OF PHAGOCYTOSIS BY SOLUTES IN CONCENTRATIONS FOUND IN THE KIDNEY AND URINE *

BY IRWIN CHERNEW AND A. I. BRAUDE

(From the Department of Medicine, University of Pittsburgh School of Medicine,
Pittsburgh, Pa.)

(Submitted for publication May 18, 1962; accepted July 5, 1962)

Immunologic processes in the kidney encounter wide variations in solute concentration and osmotic pressure that exist in no other extracellular fluids (1). It can be anticipated, for example, that leukocytes in the kidney might be exposed to 425 mM per L sodium and 850 mM per L urea, tissue concentrations demonstrated in the renal papillae of mammals. Between the extremes of maximum diuresis and antidiuresis, phagocytic cells might be exposed to tubular fluid that ranges from under 50 to over 1,300 mOsm per L, or approximately one-sixth to four times the osmotic pressure of plasma (1). In order to determine how leukocytes function in this environment, the present study was made of phagocytosis in urine and in improvised fluids containing solutes in the concentrations found in the kidney and urinary tract. The results show that such concentrations of urinary solutes depress phagocytosis and may thereby contribute to the unique susceptibility of the kidney to infection by bacteria that seldom produce disease elsewhere.

METHODS

A. Technique for studying phagocytosis

Four-tenths (0.4) ml fresh heparinized human venous blood or 0.2 ml plasma-leukocyte suspensions were added to 2 ml of test solution and mixed with 0.2 ml of bacterial suspensions in sterile pyrogen-free glass tubes 10 to 11 mm in diameter and 10 cm long. The tubes were immediately stoppered and rotated mechanically at 5 revolutions per minute at 37° C. Tubes containing staphylococci were rotated for 30 minutes and those containing *Escherichia coli* for 2 hours, periods found in preliminary studies to be required for heavy phagocytosis of either species of bacteria. After rotation the tubes were inverted 10 to 15 times, and smears prepared on glass slides for staining with combined Wright-Giemsa's stain.

The number of bacteria per 100 neutrophils and the per cent of cells exhibiting phagocytosis were compared

* Supported in part by grant H-3220, U. S. Public Health Service.

with control values. The results are indicated in the tables by the following symbols: %P = per cent neutrophils exhibiting phagocytosis; B/N_x = number bacteria ingested per 100 neutrophils in test solution; B/N_c = number bacteria ingested per 100 neutrophils in control solution; %C, phagocytic index, = $B/N_x \div B/N_c \times 100$; a subscript indicates the number of neutrophils containing too many bacteria to count, e.g., $B/N_x = 2500_s$ indicates that 2,250 bacteria were counted in 95 neutrophils and 5 neutrophils contained too many bacteria to count. For purposes of calculating %C, it was arbitrarily assumed that such phagocytic cells contained 50 bacteria each.

B. Preparation of plasma-leukocyte suspensions

The red cells in heparinized venous human blood were allowed to become sediment at 37° C and the plasma-leukocyte supernatant fluid was removed for study. The leukocyte count was 1.5 to 2.0 times that of the original whole blood. Plasma-leukocyte suspensions were used as indicated below; otherwise the experiments were done with fresh heparinized whole blood to avoid the injury to leukocytes that might occur during separation from red cells. The presence of plasma constituents and erythrocytes in the test suspensions was considered desirable from the standpoint of reproducing conditions of phagocytosis that exist in the kidney and urine during phagocytosis.

C. Preparation of bacterial suspensions

1. *Staphylococci*. A strain of *Staphylococcus epidermidis* isolated from infected human urine was grown in trypticase soy broth for 18 to 20 hours at 37° C and then killed by heating for 30 minutes at 72° C. The bacterial count of this suspension was 2.0×10^6 per mm^3 as determined in a hemocytometer chamber. The dead non-pathogenic staphylococci were used because they permitted a study of phagocytosis of particles offering no discernible resistance to ingestion.

2. *Escherichia coli*. A strain of *E. coli* (Serotype 0:113) isolated from infected human urine was cultured in trypticase soy broth for approximately 18 hours and the bacterial growth resuspended in Ringer's solution. A fresh culture of *E. coli* was prepared for each experiment and the concentration of bacilli determined with the hemocytometer. Most suspensions contained about 10^6 bacteria per mm^3 with a range from 3.6×10^6 per mm^3 to 1.8×10^6 per mm^3 . After mixture with blood or plasma-leukocytes, the concentration of bacteria was about 7 to 8 per cent of the original. The ratios of bacteria to

leukocytes ranged from 70:1 to 400:1 and were kept constant in each experiment. These ratios of *E. coli* to leukocytes were selected in order to reproduce the rigorous challenge to phagocytosis encountered in heavily infected pyuric urines. The high consistency of the results recorded below were not improved when extracellular bacteria were removed by washing the leukocytes in lactated Ringer's solution just before preparing the smears.

D. Preparation of test solutions

Lactated Ringer's solution was used as the basic medium for phagocytosis and the solutes were altered according to the aims of the experiments; cation concentrations were adjusted by changing the concentration of their chloride salts. Ringer-Locke solution was used for titrating the effect of variable sodium concentrations on phagocytosis. The formula of the lactated Ringer's solution (Abbott) was 130 mM per L Na, 4 mM per L K, 1.5 mM per L Ca, 109 mM per L Cl, and 28 mM per L lactate. Ringer-Locke solution was prepared by adding the desired quantities of solute to distilled water; it contained 130 to 150 mM per L NaCl, 3 to 5 mM per L KCl, 1.5 mM per L CaCl₂, 100 to 200 mg per cent glucose, and 4.0 mM per L NaHCO₃. The exact concentrations of solutes are recorded for individual experiments.¹

E. Preparation of urine

Most studies were conducted with whole fresh urines, or with the cell-free supernatant fluids of freshly obtained pyuric urines. A few urines were stored at -20° C before the experiment.

Osmolarities of the urines were measured in the Fiske osmometer and pH of the final mixtures in the Beckman pH meter. Sodium and potassium were determined by photometry, urea by urease nesslerization, and chloride by titration with mercuric ion. The relative degree of phagocytosis in urine was based on control values determined simultaneously in lactated Ringer's solution.

F. Adjustment of pH

The effect of pH on phagocytosis was studied after diluting 0.8 ml of a plasma-leukocyte suspension with 4.0 ml lactated Ringer's solution; then 0.06 N HCl was

added until the desired pH was obtained as determined in the Beckman pH meter. Two ml of each mixture was removed to determine pH; the remaining 2.8 ml was inoculated with 0.2 ml of the standard *E. coli* suspension and phagocytosis determined in the usual fashion. The white blood cell (WBC) count of the plasma-leukocyte suspension before dilution with lactated Ringer's solution was approximately twice that of blood, and its erythrocyte count, approximately 10⁶ per mm³. The addition of each 0.1 ml of 0.06 N HCl raised chloride concentration approximately 3 to 4 mM per L.

RESULTS

I. Phagocytosis in improvised fluids

A. Sodium. As shown in Table I, sodium in concentrations of 300 mM per L and greater markedly reduced phagocytosis of both staphylococci and *E. coli*. At concentrations of sodium below 300 mM per L, hypertonic solutions produced little depression of phagocytosis. Phagocytosis also appeared to be reduced when the concentration of sodium fell below 100 mM per L.

Disturbed phagocytosis in high and low sodium concentration was associated with morphologic abnormalities. The cells in solutions of high sodium concentrations were shrunken, and their nuclei were often fused or in the process of being extruded. Clumping of cells, observed in the control solutions, also failed to occur. In the dilute solutions, the neutrophils were sometimes swollen and clumped.

B. Urea. As shown in Tables II and III, phagocytosis was depressed at least 50 per cent at urea concentrations above 500 mM per L, and impairment became progressively reduced until marked inhibition was observed at 1,000 mM per L. The cells were usually reduced in size in the highly concentrated solution. The nuclear lobes of some neutrophils had fused.

C. Sodium and urea in combination. In order to determine if the deleterious effects of high concentrations of sodium and urea were additive, both were studied in the same solution. The results are given in Table IV.

Addition of urea to solutions already hypertonic with 250 mM per L sodium further depressed phagocytosis of *E. coli*, and to a much greater degree than when those concentrations of urea were added to solutions isotonic with 150 mM per L sodium. In hypotonic solutions of sodium, the addition of urea also impaired phago-

¹The final concentration of solute depends not only on the concentration in the original test solution, the plasma, and the bacterial suspension, but also on the amount of solute diffusing into the cells. Only the concentrations in the original solution are listed. The final concentration of solutes, such as urea, that are present in lower concentration in the blood than in the test solution would be about 25 per cent less than the recorded value. In the solutions studied for the effects of high sodium concentrations, the actual concentrations of this solute would be 1 to 10 per cent less after blood and bacteria were added. In test solutions with low sodium values, the final difference was not greater than 10 per cent except for the lowest concentration, which doubled in the final solution.

TABLE I
Phagocytosis in varying concentrations of sodium chloride*

		mM/L Sodium								
		392	324	244	167	157†	92	11.3		
Phagocytosis of <i>Staphylococci</i>										
Subject 1, whole blood	%P	37	64	100	100	100	100	59		
	B/N _x	166	296	1567 ₋₄	2810 ₋₈	2620 ₋₆	2414 ₋₃	981 ₋₃		
	%C	6.3	11.3	59.8	100	100	92.1	43.2		
Subject 2, whole blood	%P	29	32	64	90	81	78	26		
	B/N _x	104	159	931 ₋₅	867 ₋₃	1020 ₋₆	926	225		
	%C	10.2	15.6	91.3	94.8	100	90.8	22.1		
		400	350	300	250	200	159†	100	50	25
Phagocytosis of <i>E. coli</i>										
Subject 3, whole blood	%P	32		47		91	87	89	71	72
	B/N _x	36		237		699	871	694	423	308
	%C	4.1		27.2		80.2	100	79.7	48.6	35.4
Plasma- leukocyte suspension	%P	1	12	15	20	37	82	94		
	B/N _x	2	48	68	78	183	289	387		
	%C	7	16.6	23.5	27	63.3	100	133.4		

* %P = per cent neutrophils exhibiting phagocytosis; B/N_x = number bacteria ingested per 100 neutrophils in test solution; B/N_c = number bacteria ingested per 100 neutrophils in control solution; %C, phagocytic index, = B/N_x ÷ B/N_c × 100; a subscript indicates the number of neutrophils containing too many bacteria to count. See Methods, section A, for details.

† This is the control solution. The basic solution is a Ringer-Locke solution varied only in sodium concentration.

cytosis, but less than in hypertonic solution of sodium.

The observation that phagocytosis in 250 mM per L sodium was only slightly reduced but became markedly depressed with the addition of 120 mM per L urea might indicate that an osmotic pressure critical to inhibition of phagocytosis had been reached. A concentration of 324 mM per L sodium chloride sharply inhibited phagocytosis, while 244 mM per L sodium chloride was only moderately inhibitory (see Table I). The addition of 120 mM per L urea is equivalent to about 60 mM per L sodium chloride in osmolarity.

Thus, osmolarity contributed by 324 mM per L sodium chloride would be approximately equivalent to that contributed by 250 mM per L sodium chloride and 120 mM per L urea.

In high concentrations of urea and sodium, the cells were usually small; in low sodium and 0 mM per L urea, cells were swollen, vacuolated, and clumped.

D. Potassium, calcium, creatinine, ammonium, and citrate. The effects on phagocytosis of each solute was determined when its concentration was manipulated in Ringer-Locke solution to represent values near or beyond the extreme that might

TABLE II
Phagocytosis of *Staphylococci* in varying concentrations of urea*

		mM/L Urea†					
		0‡	102	360	600	900	1,200
Subject 3, whole blood	%P	100	99	96	61	41	44
	B/N _x	3,083 ₋₃	2,187 ₋₄	1,704	477	440	292
	%C	100	70.9	55.3	15.3	14.3	9.5

* Symbols as in Table I.

† Because of dilution by blood, the final concentration is approximately 25 per cent less than that recorded here in the test solution (see Methods, section D). The final concentration after addition of test solution with 900 mM per L urea is approximately 675 mM per L, a figure well below that in the dog renal medulla (1).

‡ Control solution.

TABLE III
Phagocytosis of *E. coli* in varying concentrations of urea*

		mM/L Urea										
		0†	100	200	300	400	500	600	700	800	900	1,000
Subject 4, whole blood ‡	%P	97	80				61	62				
	B/N _x	100	511				324	348				
	%C	100	71.8				45.5	48.9				
Subject 5, whole blood ‡	%P	90	87				69	51				
	B/N _x	429	359				250	107				
	%C	100	83.7				58.3	25.0				
Subject 6, whole blood ‡	%P	98	98	97	90	95	85	72	51	25	33	11
	B/N _x	749	518	636	431	519	329	213	187	150	88	44
	%C	100	59	85	58	69	44	28	25	20	12	6
Subject 7, plasma- leukocyte suspension‡	%P	99			87	69	81	33				4
	B/N _x	452			355	257	356	143				9
	%C	100			78.5	56.9	78.8	31.6				2.0

* Symbols as in Table I.

† Control solution.

‡ Suspended in lactated Ringer's solution as described in Methods, section A.

TABLE IV
Combined effect of sodium and urea on phagocytosis of *E. coli* in Ringer-Locke solution*

mM/L Sodium	mM/L Urea	250	250	250	150	150	150	150	50	50	50	50
		0	120	400	0	120	400	600	0	120	400	600
Subject 7	%P	63	27	19	90	87	69	37	60	71	41	44
	B/N _x	218	59	34	429	359	250	107	175	219	106	91
	%C	50.8	13.8	7.9	100	83.7	58.3	25.0	40.8	51.0	24.7	21.2
Subject 8	%P	85	69	63	97	99	92	78	56	93	83	49
	B/N _x	562	284	253	872	783	516	365	263	496	363	236
	%C	64.0	32.0	29.0	100	90	59	42	30	57	42	27
Subject 9	%P	96	79	71	96	97	90	88	92	97	100	84
	B/N _x	625	292	213	1,394	872	523	462	859	1,074	938	424
	%C	44.8	20.9	15.3	100	63.4	37.5	33.1	61.6	77.0	67.3	41.6

* Symbols as in Table I.

TABLE V
Comparative influence of extreme concentrations of sodium, potassium, calcium, creatinine, citrate, and ammonium on phagocytosis of *Staphylococci* in lactated Ringer's solution*

Altered solute	Concentration	%P	B/N _x	%C	Na	K	Cl	Lactate	Ca
	mM/L				mM/L	mM/L	mM/L	mM/L	mM/L
Sodium	400	16	66	1.8	400	4	379	28	1.5
Potassium	130	84	1,260	34.7	130	130	235	28	1.5
Calcium	0	98	2,988 ₋₂	82.3	130	4	103	28	0
Calcium	15	98	3,963 ₋₁₂	109.2	130	4	136	28	15.0
Creatinine	31.8	94	1,959 ₋₅	54.0	130	4	109	28	1.5
Citrate	15.8	95	1,688 ₋₃	46.5	130	4	109	28	1.5
Ammonium	70	77	1,618 ₋₃	44.6	130	4	109	28	1.5
Serum		100	3,377 ₋₂₄	91.4		4	109	28	1.5
Lactated Ringer's solution†		100	3,629	100.0	130	4	109	28	1.5

* Symbols as in Table I.

† Control.

TABLE VI
Effect of pH on phagocytosis of *E. coli* in plasma-leukocyte suspensions*

Mixtures	%P	B/N _x	%C	pH ₁ †	pH ₂	0.06 N HCl
1†	94	622	138.0	7.3	7.2	0
2‡	86	462	103.0	7.5	6.9	0
3 (control)	95	450	100.0	7.6	7.0	0
4	82	532	118.0	6.2	6.2	.2
5	3	8	1.8	5.4	5.5	.3
6	3	4	.9	5.0	5.0	.4

* Symbols as in Table I; 0.8 ml plasma-leukocyte suspension diluted with 4.0 ml lactated Ringer's solution.

† pH₁ was determined when phagocytosis began and pH₂ when it was completed.

‡ Whole blood substituted for plasma-leukocyte suspension.

§ 0.4 ml plasma-leukocyte suspension instead of 0.8 ml.

be encountered in urine (Table V). The concentration of sodium, 400 mM per L, was close to the maximum observed in the renal medulla (1). The test solutions were examined simultaneously with leukocytes from the same sample of fresh human blood; hence, the results permit direct comparison of the different solutes. Although high concentrations of potassium, ammonium, creatinine, and citrate impaired phagocytosis of *Staphylococci* the inhibitory action of these solutes was mild compared to that of sodium. The

extreme concentrations of calcium failed to influence phagocytosis significantly (see Table V).

E. coli. Phagocytosis was sharply reduced between pH 5.4 and 6.2. The findings presented in Table VI were reproduced in three additional experiments. The neutrophils at pH 5.4 or less were shrunken and the nuclear lobes fused. Although the neutrophils exhibited good phagocytosis at pH 6.2, they seemed to be extruding cytoplasm and clumping in it. The *E. coli* were also clumped in the extruded cytoplasm.

TABLE VII
Effect of glucose on phagocytosis of *Staphylococci* and *E. coli**

Subject	Glucose	%P	B/N _x	%C
	%			
Phagocytosis of <i>Staphylococci</i>				
1	12.5	32	200	6.1
	7.4	90	1,220	36.9
	4.8	91	1,255	38.0
	2.5	99	1,594	48.2
	1.25	100	2,894 ₋₄	87.6
	†	100	3,304 ₋₂	100
2	10.0	62	431	21.7
	2.0	75	871	43.9
	.05	92	1,786 ₋₂	90.0
	†	90	1,984 ₋₁	100
3	8.0	73	790	32
	6.0	88	992 ₋₁	42
	2.0	97	2,000	81
	.05	92	2,113	86
	†	97	2,454 ₋₃	100
Phagocytosis of <i>E. coli</i>				
4	10	4	4	1
	5	98	497	66
	2	99	786	105
	†	98	749	100

* Symbols as in Table I.

† Control.

TABLE VIII
Phagocytosis of *Staphylococcus* in normal urine*

Subject	%P	B/N _x	B/N _o	%C	mOsm	Na	Urea	K	Cl	pH
						mM/L	mM/L	mM/L	mM/L	
1	54	509	2664	19.0	589	155	179	64	196	7.1
2	17	39	3767	1.0	>860†	170	605	96	210	6.5
3	40	644 ₋₁	2339	27.5	484	147	124	49	183	
4	3	8	2211	.4	>860	155	600	19	152	5.9

* Symbols as in Table I.

† Maximum value measurable on the osmometer.

F. Glucose. As shown in Table VII, inhibition of phagocytosis usually began above 2.0 per cent and became marked above 7 per cent.

II. Phagocytosis in urine

A. Normal and abnormal urines. The amount of phagocytosis occurring in urines ranged from 0 to 100 per cent of the controls as shown in Tables VIII, IX, and X. Phagocytosis was inhibited by urines with high osmolarity and high urea concentrations. In urines with osmolarity greater than 860 mOsm, or urea concentration greater than 400 mM per L, the phagocytic index (%C) never exceeded 8.6. The effects of the other factors studied are not clear. The infected urine of Subject 3 (see Table X), a diabetic with slight glycosuria, reduced phagocytosis to 1 per cent of control values, even though osmolarity and pH were not markedly abnormal. Depression of phagocytosis might be partly attributed to the low concentration of sodium. As shown in Table I, the concentration of 58 mM per L sodium in this subject's urine reduced phagocytosis about 50 per cent when titrated in Ringer's solution. It is also possible that bacterial toxins injured the leukocytes in this case.

B. Effect of dilution or dialysis of urine on phagocytosis. Urines were obtained from subjects before and after they drank water in order to determine if the inhibitory effect of urine on phagocytosis can be removed by dilution *in vivo*. In addition, the nondialyzable portions of concentrated urines were examined for their inhibitory effect on phagocytosis.

The results in Table XI demonstrated that ingestion of water can abolish the inhibitory effect of concentrated urine on phagocytosis, and they support the concept that high osmolarity and high concentrations of urea inhibit phagocytosis in urine. Again there was marked inhibition of phagocytosis at osmolarity greater than 860 mOsm and urea concentrations more than 400 mM per L. Loss of the inhibitory effect by dialysis of concentrated urine can also be explained by a reduction in tonicity, and clearly demonstrates that the inhibitors to phagocytosis are absent from the nondialyzable fraction. A rise in pH might have contributed to the improved phagocytosis both in dialyzed urines and in those obtained after water diuresis, but this possibility was not supported by the observation that the pH of the concentrated urines did not fall below 5.8, a level that

TABLE IX
Phagocytosis of *E. coli* in normal urine*

Subject†	%P	B/N _x	B/N _o	%C	mOsm	Na	Urea	K	Cl	pH
						mM/L	mM/L	mM/L	mM/L	
1a	11	31	455	6.8	>860	156	428	115	181	6.5
1b	14	42	455	9.2	540	70	340	30	43	6.3
1c	0	0	455	0.0	>860	95	567	74	64	6.0
1d	16	50	455	11.0	548	94	283	34	83	6.3
2	93	501	1,101 ₋₄	45.5	292	49	59.5	12.4	34.0	7.0
3a	2	7	1,181	.6	>860	163	388	54.0	128.0	5.8
3b	94	540	1,181	45.7	188	37	75	11.0	20.0	7.1

* Symbols as in Table I.

† Numbers refer to the human subject and letters to different samples from that subject.

TABLE X
Phagocytosis of *Staphylococci* in abnormal urine*

Subject's diagnosis	%P	B/N _x	B/N _o	%C	mOsm	Na	Urea	K	Cl	pH	Glucose
						mM/L	mM/L	mM/L	mM/L		
1. Chronic pyelonephritis; no bacteria or pyuria in this urine	26	87	1,118	8.5	759	214	263	70	202	7.4	0
2. Rt. staghorn calculus. <i>K. pneumoniae</i> cultured from urine	26	146	1,003	14.5	undetermined	90	143	36	70	6.0	0
3. Diabetes; exacerbation acute pyelonephritis; <i>P. mirabilis</i> and <i>E. coli</i> in pyuric urine	8	24	2,221	1.0	350	58	188	22	55	6.5	<1
4. Hyperthyroid; atonic bladder; <i>P. aeruginosa</i> cultured from pyuric urine	100	440	1,474	30.0	375	83	178	39	89	6.3	0

* Symbols as in Table I.

TABLE XI
Influence of water diuresis on the inhibitory effect of urine on phagocytosis of *E. coli* and *Staphylococci**

Urine	%P	B/N _x	B/N _o	%C	mOsm	Na	Urea	K	Cl	pH	
						mM/L	mM/L	mM/L	mM/L		
Subject 1, with <i>E. coli</i>											
1st	15	38	455	8.4	>860	151	467	5.9	104	6.3	
2nd	58	265	455	58.2	564	136	250	9.4	118	6.6	
after drinking water											
3rd	78	508	455	111.6	305	75	125	3.2		7.1	
after drinking water											
Subject 2, with <i>E. coli</i>											
1st	11	19	514	3.7	>860	268	416	5.4		6.0	
2nd	91	736	514	143.2	340	124	83	3.5		7.5	
after drinking water											
1st, non-dialyzable portion†	89	494	514	92.2	260					7.1	
Subject 3, with <i>Staphylococci</i>											
1st	3	8	2,211	.4	>860	155	600	19	152	5.9	
2nd	98	1,676	2,211	75.8	696	129	300	20	189	5.8	
after drinking water											
1st, non-dialyzable portion†	100	1,718	2,211	77.7	266						

* Symbols as in Table I.

† Dialyzed against running distilled water for 3 hours and then against lactated Ringer's solution for 1 hour.

was not markedly inhibitory, as shown in Table XI.

III. Statistical analysis

Calculation of correlation coefficients by the formula $r = \Sigma x^2y^2 / (\Sigma x^2 \Sigma y^2)^{1/2}$ of Pearson and Lee disclosed a significant inverse relationship ($p = .01$) between phagocytosis and concentrations of sodium, urea, and glucose, and between phagocytosis and osmolarity of normal urine.

DISCUSSION

Increasing concentrations of either sodium chloride, urea, or hydrogen ion progressively depressed phagocytosis until it was virtually abolished at a point well below the highest concentrations anticipated in the renal papilla. Moreover, sodium and urea in combination nearly abolished phagocytosis in concentrations that were only moderately inhibitory when these solutes were examined individually. This combined effect seemed to operate not only in artificial test solutions but also in those concentrated urines where phagocytosis was depressed to a greater extent than that anticipated from the effect of any solute alone. It is possible that the high concentration of these and other solutes in the kidney and urine also depressed phagocytosis *in vivo* in normal individuals. In diabetes, phagocytosis might be further depressed by severe glycosuria.

The mechanisms responsible for the antiphagocytic effect of these substances remain to be fully clarified, but good evidence suggests that high osmotic pressure is an important factor. A direct correlation could be made between the antiphagocytic properties of solutions and their osmolarities. The appearance of leukocytes also seemed to be related to tonicity. Thus in solutions made hypertonic with sodium, urea, and glucose, the leukocytes usually were contracted, while in hypotonic solutions they usually were swollen. The osmotic effectiveness of urea was also suggested by the observation that phagocytosis in 250 mM per L sodium was sharply depressed by the addition of only 120 mM per L urea, the amount required to produce an osmolarity shown to be critical for marked inhibition by sodium alone. The fact that urea has also been shown to be osmotically effective in preventing lysis of bacterial

protoplasts in urine might indicate that this solute does not freely penetrate the cell water of microbial cells or leukocytes (2).

A possible application of these findings to the problem of urinary infection was suggested by the improved phagocytosis noted in urine obtained from subjects who had drunk water. The inhibitory effect of their hypertonic morning urines was abolished when their urines approached isotonicity. Occasional urines in the isotonic range inhibited phagocytosis, but these were heavily infected with bacteria whose toxins could have injured the leukocytes.

Inhibition of phagocytosis also occurred in hypotonic solutions, where the neutrophils became swollen and distorted. Phagocytosis in markedly hypotonic urines was not examined because none was obtained for study, but it is reasonable to expect that excessive dilution of urine might injure leukocytes and reduce their activity.

Besides sodium and urea, hydrogen ion in high concentration also depressed phagocytosis sharply. This inhibitory effect occurred rather abruptly between pH 6.2 and 5.5. Because the pH of urine in distal tubules and bladder may fall well below this level, acidity may be an important factor in preventing phagocytosis in the urinary tract. The fact that complement is rapidly destroyed at pH less than 5.5 (3) and is normally involved in phagocytosis suggests that loss of this substance may contribute to the inhibition of phagocytosis (4).

These studies have attempted to delineate some of the factors that might depress phagocytosis in the urinary system, but others remain to be explored. Observations summarized nearly 30 years ago in the review by Mudd, McCutcheon, and Lucke (5) indicate that many variables in the physical and chemical environment may stimulate or depress phagocytosis. It must be emphasized, also, that the present experiments were done *in vitro* and that only limited inferences may be drawn to conditions that prevail *in vivo*. Nonetheless, it is becoming increasingly evident that a variety of chemical conditions peculiar to the kidney might lower resistance to infection. Evidence has been obtained in other studies that the high urea concentration of the renal medulla favors infection by *Proteus* organisms because of their powerful urease (6), that intrarenal formation of

ammonia inactivates the bactericidal system by destroying the fourth component of complement (7), and that hypertonicity allows protoplasts to develop and persist during antibiotic therapy (8). This evidence and that in the present study together may help explain why the kidney is uniquely susceptible to invasion by *E. coli*, *Proteus*, and other bacteria that possess little capacity to establish infection elsewhere.

SUMMARY

In order to elucidate the peculiar susceptibility of the kidney to infection by bacteria that seldom produce disease elsewhere, a study was made of phagocytosis in fluids containing solutes in the variable concentrations found only in the kidney and urinary tract.

Within the range found in the normal kidney and urine, high concentrations of sodium, urea, and hydrogen ion inhibited phagocytosis by human leukocytes. Phagocytosis was inhibited by glucose in concentrations found in the urinary tract in diabetes mellitus. Highly concentrated human urines

also depressed phagocytosis, but more dilute urines usually did not. Evidence has thus been obtained that phagocytosis might be impaired in the urinary system and that resistance to infection in the kidney might be thereby reduced.

REFERENCES

1. Lamdin, E. Mechanisms of urinary concentration and dilution, *A.M.A. Arch. intern. Med.* 1959, **103**, 644.
2. Braude, A. I., and Siemienski, J. Urinary Protoplasts. To be published.
3. Boyd, William C. *Fundamentals of Immunology*, 3rd ed. New York, Interscience, 1956, p. 373.
4. Berry, L. J., and Spies, T. D. Phagocytosis. *Medicine* 1949, **28**, 239.
5. Mudd, S., McCutcheon, M., and Lucké, B. Phagocytosis. *Physiol. Rev.* 1934, **14**, 210.
6. Braude, A. I., and Siemienski, J. Role of bacterial urease in experimental pyelonephritis. *J. Bact.* 1960, **80**, 171.
7. Beeson, P. B., and Rowley, D. The anticomplementary effect of kidney tissue. Its association with ammonia production. *J. exp. Med.* 1959, **110**, 685.
8. Braude, A. I., Siemienski, J., and Jacobs, I. Protoplast formation in human urine. *Trans. Ass. Amer. Phycns* 1961, **74**, 234.