

Rate of disappearance of labeled carbon dioxide from the lungs of humans during breath holding: a method for studying the dynamics of pulmonary CO₂ exchange

Richard W. Hyde, ... , William F. Raub, Robert E. Forster

J Clin Invest. 1968;47(7):1535-1552. <https://doi.org/10.1172/JCI105846>.

Research Article

The dynamics of CO₂ exchange in the lungs of man was studied by observing the rate of disappearance of a stable isotope of CO₂ (¹³CO₂) from the alveolar gas during breath holding. Over 50% of the inspired isotope disappeared within the first 3 sec followed by a moderately rapid logarithmic decline in which one-half of the remaining ¹³CO₂ disappeared every 10 sec.

The large initial disappearance of ¹³CO₂ indicated that alveolar ¹³CO₂ equilibrated in less than 3 sec with the CO₂ stored in the pulmonary tissues and capillary blood. The volume of CO₂ in the pulmonary tissues calculated from this initial disappearance was 200 ml or 0.33 ml of CO₂ per milliliter of pulmonary tissue volume.

The alveolar to end-capillary gradient for ¹³CO₂ was calculated by comparing the simultaneous disappearance rates of ¹³CO₂ and acetylene. At rest and during exercise this gradient for ¹³CO₂ was either very small or not discernible, and diffusing capacity for CO₂ (D_{LCO₂}) exceeded 200 ml/(min × mm Hg).

After the administration of a carbonic anhydrase inhibitor the rate of disappearance of ¹³CO₂ decreased markedly. D_{LCO₂} fell to 42 ml/(min × mm Hg) and at least 70% of the exchange of ¹³CO₂ with the CO₂ stores in the pulmonary tissues and blood was blocked by the inhibitor. These changes were attributed to impairment of exchange of [...]

Find the latest version:

<https://jci.me/105846/pdf>



Rate of Disappearance of Labeled Carbon Dioxide from the Lungs of Humans during Breath Holding: a Method for Studying the Dynamics of Pulmonary CO₂ Exchange

RICHARD W. HYDE, RICARDO J. M. PUY, WILLIAM F. RAUB, and ROBERT E. FORSTER

From the Department of Physiology, Division of Graduate Medicine, and the Department of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104

ABSTRACT The dynamics of CO₂ exchange in the lungs of man was studied by observing the rate of disappearance of a stable isotope of CO₂ (¹³CO₂) from the alveolar gas during breath holding. Over 50% of the inspired isotope disappeared within the first 3 sec followed by a moderately rapid logarithmic decline in which one-half of the remaining ¹³CO₂ disappeared every 10 sec.

The large initial disappearance of ¹³CO₂ indicated that alveolar ¹³CO₂ equilibrated in less than 3 sec with the CO₂ stored in the pulmonary tissues and capillary blood. The volume of CO₂ in the pulmonary tissues calculated from this initial disappearance was 200 ml or 0.33 ml of CO₂ per milliliter of pulmonary tissue volume.

The alveolar to end-capillary gradient for ¹³CO₂ was calculated by comparing the simultaneous disappearance rates of ¹³CO₂ and acetylene. At rest and during exercise this gradient for ¹³CO₂ was

A preliminary report of this work was presented at the Fall Meeting of the American Physiological Society, Buffalo, N. Y. 30 August 1962 (1).

Dr. Hyde's work was done in part during the tenure of a Daland Fellowship of the American Philosophical Society and an established investigatorship of the American Heart Association. Dr. Puy is a Fellow of the Consejo Nacional de Investigaciones Cientificas y Tecnicas, Buenos Aires, Argentina.

Received for publication 4 December 1967 and in revised form 14 March 1968.

either very small or not discernible, and diffusing capacity for CO₂ (DL_{CO₂}) exceeded 200 ml/(min × mm Hg).

After the administration of a carbonic anhydrase inhibitor the rate of disappearance of ¹³CO₂ decreased markedly. DL_{CO₂} fell to 42 ml/(min × mm Hg) and at least 70% of the exchange of ¹³CO₂ with the CO₂ stores in the pulmonary tissues and blood was blocked by the inhibitor. These changes were attributed to impairment of exchange of ¹³CO₂ with the bicarbonate in the pulmonary tissues and blood.

The pH of the pulmonary tissues (V_{tis}) was determined by a method based on the premise that the CO₂ space in the pulmonary tissues blocked by the inhibitor represented total bicarbonate content. At an alveolar P_{CO₂} of 40 mm Hg pH of V_{tis} equalled 6.97 ± 0.09.

INTRODUCTION

Carbon dioxide is present in the blood and tissues of the lungs in the form of physically dissolved CO₂, bicarbonate, carbonic acid, and protein complexes (2). Chinard, Enns, and Nolan studied the interaction of these forms of CO₂ and the alveolar gas in the dog by injecting radioactive bicarbonate (H¹⁴CO₃⁻) and carbon dioxide (¹⁴CO₂) into the pulmonary artery (3, 4). They concluded that the interconversion of the various forms of

CO₂ is so rapid that the expired CO₂ may be considered to come from one large common CO₂ pool. Similar conclusions were reached by Feisal, Sackner, and DuBois who injected bicarbonate into the pulmonary artery of the dog and measured the evolution of CO₂ in the alveoli with a sensitive body plethysmograph (5). These investigators as well as others have found that inhibition of the enzyme carbonic anhydrase modified the exchange of CO₂ between blood, pulmonary tissues, and alveolar gas sufficiently to produce a significant alveolar to end-capillary CO₂ gradient (A-c CO₂ gradient) (6-8).

By measuring the rate of uptake of a stable carbon dioxide isotope (¹³CO₂) from the alveolar gas during breath holding, we have developed a new technique for studying the interrelationships of the various forms of CO₂ and the A-c CO₂ gradient in the lungs of man. Measurements of the uptake of isotopic CO₂ have the advantage that they can provide the greatest possible A-c CO₂ gradient for a given total alveolar CO₂ tension, pulmonary capillary blood flow, and diffusing capacity. From the data it was possible to estimate the A-c CO₂ gradient and the diffusing capacity of the lungs for CO₂, and to determine the size of the CO₂ spaces in the lungs, the CO₂ dissociation curve of the lung tissues, the mean hydrogen ion concentration of the pulmonary tissues, and the relative contribution of bicarbonate to total CO₂ in the lungs.

METHODS

If a subject inspires a breath of a gas mixture containing CO₂ which has been enriched with the stable carbon isotope of mass 13 (¹³CO₂), the distribution of the isotope within the lungs is likely to be governed by factors similar to those reported to influence the disappearance of soluble inert gases such as nitrous oxide and acetylene (9). There should be a rapid disappearance of the isotope initially because of its movement into the pulmonary parenchymal tissue volume (V_{tis}) and pulmonary capillary blood volume (V_c), followed by a more gradual disappearance as it is carried away by the blood flowing through the pulmonary capillaries. If the highly insoluble inert gas, neon, is added to the inspired mixture, the effect of dilution of the isotope in the lung's residual air can be taken into account. If acetylene (C₂H₂), whose rate of disappearance during breath holding is a function of pulmonary capillary blood flow (Q̇_c), is included in the inspired mixture, it is possible to evaluate the influence of Q̇_c on the rate of disappearance of ¹³CO₂ (9). The CO₂ content of the pulmonary parenchymal tissue (V_{tis}) and V_c can be determined from the initial

rapid disappearance of ¹³CO₂. The alveolar to end-capillary gradient for CO₂ can be calculated by comparing the rates of disappearance of C₂H₂ and ¹³CO₂. Once this gradient is known, the diffusing capacity for CO₂ (DL_{CO2}) can be determined. The relative contribution of bicarbonate to total CO₂ transport can be evaluated by making measurements before and after the administration of a carbonic anhydrase inhibitor.

Experimental procedure. ¹³CO₂ was prepared by adding hydrochloric acid to barium carbonate enriched 10-fold or more with carbon atoms of mass 13.¹ The experiments were conducted in the following manner: a rapidly responding mass spectrometer was tuned to the mass 44 peak in order to measure ¹³CO₂, and the instrument's inlet tube was attached to a mouthpiece through which the subject first blew out to residual volume. He then began rebreathing for three to six breaths from an unheated rubber anesthesia bag containing approximately 2 liters of 7-12% CO₂, 0-12% oxygen, and balance nitrogen. In the early experiments the per cent of O₂ in the rebreathing bag and inspired mixture was 21%. Because under these conditions the O₂ saturation of the blood increased from about 75-98% on entering the pulmonary capillaries, the P_{CO2} would be expected to be slightly higher in the capillary blood than in the mixed venous blood (Haldane effect). In order to minimize this difference between mixed venous P_{CO2} and capillary blood P_{CO2} in the later experiments, we made the capillary O₂ saturation similar to the mixed venous level by using no O₂ in the rebreathing bag and by reducing the O₂ concentration in the inspired mixture to 5%.

The concentration and volume of CO₂ in the rebreathing bag was chosen so that after several breaths, the output of the mass spectrometer became constant indicating that the partial pressure of CO₂ (P_{CO2}) in the subject's alveolar volume (V_A) and the rebreathing bag were at the virtual mixed venous P_{CO2} (10) (Fig. 1). The subject then exhaled to his residual volume and maximally inspired a gas mixture containing a total P_{CO2} approximately equal to the virtual mixed venous P_{CO2}, 0.6-1.0% ¹³CO₂, 0.4% CO, 0.5% neon, 0.75% C₂H₂, 5 or 21% O₂, and balance nitrogen. After breath holding for about 10 sec, he forcefully expired. Note that the concentration of CO₂ in the rebreathing bag and the inspired mixture were chosen so that during breath holding the P_{CO2} in the blood entering the capillaries was similar to the P_{CO2} in V_A. As a result the total P_{CO2} and CO₂ content in the pulmonary capillaries and alveoli remained relatively constant during the breath holding period. The first liter of the expirate was discarded in order to wash out the respiratory dead space, and the remaining gas was collected for analysis. The procedure was repeated after intervals of an hour for breath holding periods of approximately 3, 7, and 14 sec. Breath holding periods longer than 14 sec were avoided because of the possibility of recirculation of the isotope.

Calculating rate of disappearance of labeled CO₂. The per cent of total CO₂ in the form ¹³CO₂ in the inspired and expired gases was measured on a mass spectrometer by recording the output of either mass 12 and mass 13 or mass

¹ Obtainable from the Isomet Corp., Palisades Park, N. J.




FIGURE 1 Tracing of a mass spectrometer record monitoring partial pressure of mass 44 ($P^{12}\text{CO}_2$) during measurement of rate of disappearance of $^{13}\text{CO}_2$ in subject RWH. Time scale commences when the subject began rebreathing from a 1.5 liter bag containing 12% CO_2 and balance air. $P^{12}\text{CO}_2$ became constant at 48.5 mm Hg and during the 14th sec he inspired the test gas mixture enriched with $^{13}\text{CO}_2$. He held his breath for 3 sec and then exhaled delivering an alveolar sample. Note that total PCO_2 is kept almost constant at 48 mm Hg, whereas $P^{12}\text{CO}_2$ falls to less than half of the value originally present in the inspired mixture.

44 and mass 45 (Table I). At the start of the experiment the natural occurrence of the isotope, defined as the concentration of the isotope divided by the total concentration of CO_2 in the specimen, was determined from a sample of the subject's expired air. Total $[\text{CO}_2]$, $[\text{C}_2\text{H}_2]$, $[\text{Neon}]$, and (carbon monoxide) of the gas samples were measured on a gas chromatograph. Alveolar PCO_2 during breath holding was determined from either the continuous mass spectrometer record of expired breath or by analyzing the contents of the rebreathing bag. Alveolar volume (V_A) during breath holding was measured by adding the inspired volume recorded on a spirometer to the residual volume which had been determined previously by the closed circuit helium method (12). The disappearance from the alveoli of C_2H_2 , carbon monoxide (CO), and $^{13}\text{CO}_2$ in excess of virtual mixed venous $^{13}\text{CO}_2$ ($^*\text{CO}_2$) were plotted on semi-logarithmic graph paper against time (Fig. 2). From the C_2H_2 disappearance, pulmonary capillary blood flow (\dot{Q}_c) and pulmonary parenchymal tissue volume (V_{tis}) were determined (9). Carbon monoxide diffusing capacity (DL_{CO}) was calculated from the disappearance of CO (13).

In practice it was found to be convenient to calculate the per cent disappearance of $^*\text{CO}_2$ by the following formula whose derivation is given in detail elsewhere (14, Appendix I):

$$\text{Fraction of } ^*\text{CO}_2 \text{ remaining} = \frac{(P_{\text{I}^{\text{neon}}})(P_{\text{E}^{\text{CO}_2}})}{(P_{\text{E}^{\text{neon}}})(P_{\text{I}^{\text{CO}_2}})} \times \left[\frac{P_{\text{E}^{13}\text{CO}_2} - \text{natural occurrence of } ^{13}\text{CO}_2}{P_{\text{E}^{\text{CO}_2}} - \text{natural occurrence of } ^{13}\text{CO}_2} \right] \quad (1)$$

$P_{\text{I}^{\text{neon}}}$ and $P_{\text{I}^{\text{CO}_2}}$ are the partial pressures of neon and CO_2 , respectively, in the test gas mixture which is to be inspired, $P_{\text{E}^{\text{neon}}}$ and $P_{\text{E}^{\text{CO}_2}}$ are the partial pressures of neon and CO_2 , respectively, in the expired gas sample, $P_{\text{I}^{13}\text{CO}_2}$ and $P_{\text{E}^{13}\text{CO}_2}$ are the partial pressures of $^{13}\text{CO}_2$ in the inspired test gas mixture and the expired gas mixture, and the natural occurrence of $^{13}\text{CO}_2$ is the fraction $[\text{I}^{13}\text{CO}_2]/[\text{total CO}_2]$ measured from the subject's expired air before inhaling gas mixtures enriched with $^{13}\text{CO}_2$.

Calculation of the CO_2 content of the lung's tissues (V_{tis}) from the rate of disappearance of labeled CO_2 . Theoretical analyses indicate that a soluble inert gas equilibrates extremely rapidly (in less than 1 sec) with the finer parenchymal tissues of the lung (9), and that the initial disappearance of such a gas can be used to calculate the pulmonary parenchymal tissue volume (V_{tis}). Experimentally the logarithm of alveolar inert gas concentration is plotted against time of breath holding and the curve extrapolated to time zero (see line for C_2H_2 in Fig. 2). Any rapid initial solution of inert gas in the lung tissue will be seen as a depression of the intercept with respect to the zero time axis, and will depend on the relative volumes of lung tissue and alveolar gas, as well as the partition coefficient of the inert gas. In the case of $^*\text{CO}_2$, while a small amount will physically dissolve in the lung tissues (V_{tis}) a larger amount will exchange with the bicarbonate contained in V_{is} and the pulmonary capillary blood volume (V_c) (3). An additional

TABLE I
Natural Occurrence of Stable Isotopes
of Carbon Dioxide*

Mass	Occurrence of mass as per cent of total CO_2	Molecular structure	Occurrence of molecular structure as per cent of total CO_2
44	98.415	$^{12}\text{C}^{16}\text{O}^{16}\text{O}$	98.415
45	1.177	$^{13}\text{C}^{16}\text{O}^{16}\text{O}$	1.103
		$^{12}\text{C}^{16}\text{O}^{17}\text{O}$	0.074
46	0.403	$^{13}\text{C}^{16}\text{O}^{17}\text{O}$	8.3×10^{-4}
		$^{12}\text{C}^{16}\text{O}^{18}\text{O}, ^{12}\text{C}^{17}\text{O}^{17}\text{O}$	
47	0.0045	$^{13}\text{C}^{16}\text{O}^{18}\text{O}, ^{13}\text{C}^{17}\text{O}^{17}\text{O}$	0.0045
		$^{12}\text{C}^{17}\text{O}^{18}\text{O}$	1.5×10^{-5}
48	4.2×10^{-5}	$^{13}\text{C}^{17}\text{O}^{18}\text{O}$	1.7×10^{-7}
		$^{12}\text{C}^{18}\text{O}^{18}\text{O}$	4.2×10^{-5}
49	4.6×10^{-7}	$^{13}\text{C}^{18}\text{O}^{18}\text{O}$	4.6×10^{-7}

* Calculated by law of probabilities from data of Nier (11) who reported occurrence of carbon-12 as 98.892% of total CO_2 , carbon-13 as 1.108% of total CO_2 , oxygen-16 as 99.758% of total O_2 , oxygen-17 as 0.0373% of total O_2 , and oxygen-18 as 0.2039% of total O_2 . In the earlier experiments analysis of the labeled CO_2 was performed by first absorbing the CO_2 in the expired gas in a precipitate free solution of barium and sodium hydroxide, and then regenerating the CO_2 by the addition of hydrochloric acid. This processing increased the CO_2 concentration to 80% or higher which gave sufficient output on the mass spectrometer to measure mass 13 (carbon-13 fraction of $^{13}\text{CO}_2$ and mass 12 (carbon-12 fraction of $^{12}\text{CO}_2$). For the later experiments a different mass spectrometer became available which had sufficient separation of masses so that mass 44 ($^{12}\text{C}^{16}\text{O}^{16}\text{O}$) and mass 45 ($^{13}\text{C}^{16}\text{O}^{16}\text{O}$ and $^{12}\text{C}^{16}\text{O}^{17}\text{O}$) could be measured directly obviating the need of concentrating the CO_2 in the gas samples.

amount of *CO₂ may exchange with CO₂ bound to hemoglobin as carbamino-hemoglobin or with other protein molecules (2). If it is assumed that the movement of *CO₂ into V_{tis} and Vc is complete in 3 sec (our shortest period of breath holding), then the amount of depression of the intercept at time zero of the curve for *CO₂ (Fig. 2) is a function of the amount of *CO₂ which is taken up by V_{tis} and Vc. The total amount of *CO₂ inspired must equal the amount of *CO₂ which is present in V_{tis}, Vc, and the alveolar volume just after inspiration and before a significant amount of *CO₂ is carried away by the pulmonary capillary blood flow ($\dot{Q}c$) or;

$$\frac{(VA)(P^{*CO_2}_{t-a})}{P_B - P_{H_2O}} = \frac{(VA)(P^{*CO_2}_{t-b})}{P_B - P_{H_2O}} + \frac{(Vc)(\alpha_b)(P^{*CO_2}_{t-b})}{760} + \frac{(V_{tis})(\alpha_{tis})(P^{*CO_2}_{t-b})}{760} \quad (2)$$

where VA equals the alveolar volume in ml STPD; P^{*CO₂}_{t-a} is the P^{*CO₂} in mm Hg in VA just after inspiration but before any P^{*CO₂} has moved into Vc and V_{tis}; Vc is the pulmonary capillary blood volume in ml; V_{tis} is the pulmonary parenchymal tissue volume in ml; P^{*CO₂}_{t-b} is the partial pressure of P^{*CO₂} in mm Hg in VA, Vc, and V_{tis} just after the movement of P^{*CO₂} into Vc and V_{tis}; α_b is the effective solubility coefficient for all forms of CO₂ carried in the blood (i.e. physically dissolved CO₂, H₂CO₃, CO₂ in bicarbonate, CO₂ in the form of carbamino-hemoglobin, and other CO₂-protein complexes) in ml of CO₂ STPD per ml of blood per standard atmosphere.² α_{tis} is the effective solubility coefficient for all forms of CO₂ present in V_{tis} in ml of CO₂ STPD per ml of V_{tis} per standard atmosphere, P_B is the barometric pressure in mm Hg, and P_{H₂O} is the vapor pressure of water at the subject's body temperature.

Note that the fraction P^{*CO₂}_{t-b} ÷ P^{*CO₂}_{t-a} equals the amount of *CO₂ present in the alveoli, determined by extrapolating the rate of disappearance of *CO₂ during breath holding back to time zero, divided by the amount of *CO₂ predicted to be present from the neon dilution (Fig. 2). This fraction therefore equals *CO₂ intercept in per cent ÷ 100 and equation 2 can be rewritten in the following manner:

$$\frac{760 VA}{P_B - P_{H_2O}} \left[\frac{100}{*CO_2 \text{ intercept in per cent}} - 1 \right] = V_{tis}(\alpha_{tis}) + Vc(\alpha_b) \quad (3)$$

Since V_{tis}(α_{tis}) + Vc(α_b) equals the CO₂ content in ml STPD of the lungs at a partial pressure of CO₂ of 760 mm Hg,

² α_b varies with the partial pressure of CO₂ present because of the alinearity of the CO₂ dissociation curve. Therefore, in the experiments in this paper α_b was always defined for the particular alveolar P_{CO₂} present during breath holding (P_{ACO₂}) and was calculated by the following formula:

$$\alpha_b = \frac{[CO_2] 760}{P_{ACO_2}}$$

where [CO₂] is the CO₂ content of the subject's blood at P_{ACO₂} in milliliter STPD per milliliter of blood. See text for technique of determining [CO₂].

the CO₂ content at the alveolar P_{CO₂} present during breath holding (P_{ACO₂}) can be calculated by multiplying the left-hand term of equation 3 by P_{CO₂} ÷ 760 or:

CO₂ content of (V_{tis} + Vc) at P_{ACO₂} =

$$\frac{(P_{ACO_2})(VA)}{P_B - P_{H_2O}} \left[\frac{100}{*CO_2 \text{ intercept in per cent}} - 1 \right] \quad (4)$$

If the CO₂ content of V_{tis} alone is desired, it can be determined by measuring Vc using carbon monoxide (15) and the CO₂ content of the subject's blood at P_{ACO₂} (see below), and then subtracting the product of these two terms from the CO₂ content of V_{tis} + Vc. In these experiments the CO₂ content of the pulmonary capillary blood during breath holding was determined indirectly from measurements of P_{CO₂} and P_{O₂} in the rebreathing bag or the expired sample. These values were then used to obtain the CO₂ content from a standard CO₂ dissociation curve (16). In two of the subjects the CO₂ content and P_{CO₂} of a blood specimen drawn from the antecubital vein were measured by the Astrup technique (17) and this value was then adjusted to the CO₂ content at P_{ACO₂} by using the nomograms published by Bartels and coworkers (18).

*Calculation of pulmonary capillary blood flow ($\dot{Q}c$) from the rate of disappearance of *CO₂.* If there is no alveolar to end-capillary gradient (A-c CO₂ gradient) for CO₂, the rate of disappearance of *CO₂ from the alveolar gas during breath holding will be flow limited. Then $\dot{Q}c$ calculated from the disappearance of *CO₂ should be equal to values obtained with an inert gas such as C₂H₂. As a means of verifying the presence or absence of a significant A-c CO₂ gradient, we compared in our subjects $\dot{Q}c$ measured by the C₂H₂ breath holding method with $\dot{Q}c$ determined from the rate of disappearance of *CO₂. $\dot{Q}c$ using *CO₂ was determined in the following manner: the instantaneous rate of change of the concentration of *CO₂ in the lungs after inspiration of the test gas and after equilibration of *CO₂ with lung tissue and capillary blood must equal the amount which is being removed by $\dot{Q}c$ or:

$$\frac{d}{dt} \left[\frac{PA^{*CO_2}}{760} \right] \left[VA \frac{760}{P_B - P_{H_2O}} + \alpha_{tis}(V_{tis}) + \alpha_b(Vc) \right] = (\dot{Q}c)(\alpha_b) \left[\frac{PA^{*CO_2}}{760} \right] \quad (5)$$

where $\dot{Q}c$ is in milliliter per minute and PA^{*CO₂} is the partial pressure of *CO₂ in the alveoli at any instant. The terms α_{tis}(V_{tis}) + α_b(Vc) can be eliminated by substitution (see equation 3). Integration over the interval of breath holding gives:

$$\dot{Q}c = \frac{VA \left[\frac{760}{P_B - P_{H_2O}} \right] \left[\frac{100}{CO_2 \text{ intercept in per cent}} \right]}{\alpha_b(t_{BH})} \times \ln \left[\frac{PA^{*CO_2}_{t-b}}{PA^{*CO_2}_t} \right] \quad (6)$$

where PA^{*CO₂} equals the PA^{*CO₂} in mm Hg at the end of breath holding and t_{BH} is the breath holding time in minutes.

Determination of the slope of the CO₂ dissociation curve of the pulmonary tissues. Techniques for the determination of the CO₂ dissociation curve of the lung tissues have included a plethysmographic method (19), direct measurement of the CO₂ content in vitro after equilibration with known concentrations of CO₂ (20), and methods based on the collection of alveolar gas samples after varying breath holding periods (21, 22). In this study a modification of the breath holding methods described by DuBois (21) and Fenn and Dejours (22) was used. The method is described in Appendix I.

Calculation of the diffusing capacity of lungs for CO₂—(DL_{CO₂}). While normally it is believed there is little or no end-capillary gradient for CO₂ between the alveolar gas, the plasma, and the red blood cells in the pulmonary end-capillary blood, Chinard and coworkers (3) and Soni, Feisal, and DuBois (23) have shown that after the administration of a carbonic anhydrase inhibitor, a significant gradient between alveolar PCO₂ and the PCO₂ in the end-capillary blood is present due to the slower conversion of bicarbonate to molecular CO₂. If a measureable gradient is produced, the diffusing capacity of the lungs for CO₂ (DL_{CO₂}) could be calculated from the rate of disappearance of *CO₂ during breath holding by a method identical with that recently described for O₂ using a stable O₂ isotope (14). Constantine, Craw, and Forster (24) have reported values for θ_{CO_2} , (defined as the rate at which blood takes up CO₂ expressed in ml of CO₂ STPD per minute per mm Hg per milliliter of blood) before and after the addition of a carbonic anhydrase inhibitor to the blood. These values for θ_{CO_2} , the calculated value of DL_{CO₂} (See above), and the diffusing capacity of the alveolar-capillary membrane (DM_{CO₂}) should be related to each other by the following equation (15):

$$\frac{1}{DL_{CO_2}} = \frac{1}{DM_{CO_2}} + \frac{1}{(V_c)(\theta_{CO_2})} \quad (7)$$

DL_{CO₂} and DM_{CO₂} are expressed in milliliter of CO₂ STPD per minute per mm Hg, and V_c is the pulmonary capillary blood volume in milliliters determined by measuring the carbon monoxide diffusing capacity at different alveolar O₂ tensions (15).

RESULTS

*Measurement of rate of disappearance of *CO₂ in five resting subjects.* Analysis of the gas samples collected after the shortest period of breath holding, which was about 2.5 sec, showed that over half of the *CO₂ had disappeared from the alveolar gas (Tables II and III, and Fig. 2). Thereafter the remaining *CO₂ decreased at approximately 10%/sec. Comparable disappearance rates have been reported by West and Dollery using radioactive CO₂ (¹⁴CO₂) (25). Pulmonary capillary blood flow calculated from the disappearance of C₂H₂ ($\dot{Q}_{C_2H_2}$) and the value obtained from the rate of disappearance of *CO₂ ($\dot{Q}^*_{CO_2}$),

TABLE II
Physical Characteristics of Experimental Subjects

Subject	Age	Sex	Height		Body surface area
			<i>yr</i>	<i>in.</i>	
RWH	32*	M	71	160	1.94
WFR	23	M	72	180	2.02
PBK	24	M	69	155	1.84
RJMP	30	M	68	170	1.90
MAF	27	F	67	118	1.62

* 36 yr old at time of exercise measurements.

differed by no more than 0.4 liter/min ($r = +0.99$, $P > 0.9$). Mean values for $\dot{Q}_{C_2H_2}$ and $\dot{Q}^*_{CO_2}$ were 7.24 liters/min and 7.23 liters/min, respectively. This close agreement indicates that on the average there was no significant alveolar to end-capillary gradient for *CO₂ because if a gradient was present, $\dot{Q}^*_{CO_2}$ would have been significantly smaller than $\dot{Q}_{C_2H_2}$ (Table III).

CO₂ content of lung tissue. The mean amount of CO₂ present in the lungs of the five subjects determined from the initial rapid loss of *CO₂ from the alveoli was 244 ml STPD (Table IV). Of this volume approximately 50 ml represented CO₂ in the capillary blood (V_c) and 194 ml represented CO₂ in the lung tissue (V_{tis}). The CO₂ concentration of the sum of V_{tis} and V_c was 0.36 ml STPD per ml and the CO₂ content of V_{tis} alone was 0.33 ml STPD per ml. If one assumes that the amount of physically dissolved CO₂ is the same as in water, then approximately 0.04 ml/ml is in the form of physically dissolved CO₂³ and 0.29 ml/ml must be present in other forms such as bicarbonate. The concentration of CO₂ in V_{tis} was about 35% less than its concentration in the pulmonary capillary blood, about equal to the values reported for muscle and brain (16, 26), and in good agreement with the measurements for lung tissue published by DuBois, Fenn, and Britt (20). They found that at an alveolar Pco₂ of 50 mm Hg dog lungs perfused with Ringer's solution contained approximately 0.35 ml of CO₂ per ml of tissue.

CO₂ dissociation curve of the lungs. The CO₂

³ At PMV_{CO₂} equal to 50 mm Hg the total physically dissolved CO₂ would be: $0.57 \times 50/760$ or 0.04-ml STPD per ml of V_{tis} where 0.57 is the solubility coefficient of CO₂ in water at 37°C in milliliter STPD per atmosphere.

TABLE III

Alveolar Volume (V_A), Disappearance Rates of C_2H_2 and $*CO_2$, Pulmonary Capillary Blood Flow (\dot{Q}_c), Alveolar PO_2 (PAO_2), Alveolar PCO_2 ($PACO_2$), and Diffusing Capacity for CO (DL_{CO}) and CO_2 (DL_{CO_2}) in Five Human Subjects

Subject	Body position	V_A ml STPD	% $*CO_2^*$ remaining at extrapolated zero time	% $C_2H_2^*$ remaining at extrapolated zero time	$K^*CO_2^\ddagger$	$K_{C_2H_2}^\ddagger$	$\dot{Q}^*CO_2^\S$	$\dot{Q}_{C_2H_2}^\S$	P_{ACO_2}	P_{AO_2}	DL_{CO}	DL_{CO_2}
					sec^{-1}	sec^{-1}	liter/min	liter/min				
A. At rest												
RWH	Sitting	5675	55.0	90.2	0.078	0.0123	6.42	6.66	50.0	130	33.1	223
WFR	Sitting	6080	60.8	89.4	0.104	0.0127	7.91	7.51	47.6	127	39.2	i
PBK	Sitting	5300	58.8	93.2	0.111	0.0165	7.79	8.12	48.0	131	33.5	224
RJMP	Supine	4100	56.7	90.5	0.128	0.0202	7.72	7.84	50.5	44	42.2	317
MAF	Supine	3960	59.7	90.2	0.115	0.0160	6.32	6.08	43.3	36	28.8	i
Mean		5025	58.2	90.7	0.107	0.0155	7.23	7.24	47.9	—	—	—
B. During exercise												
RWH	Supine	5230	55.7	90.6	0.149	0.025	12.0	12.6	54.1	32	60.0	263
C. After acetazolamide												
RWH	Sitting	5770	82.7	89.6	0.062	0.0195	3.01	10.9	40.5	137	40.5	42

* Calculated by extending the line of least mean squares drawn from the plots of the per cent of $*CO_2$ or C_2H_2 remaining in the alveolar gas samples back to the point on the time axis representing the start of breath holding (Figs. 2, 5, and 6).

‡ Calculated from the line of least mean squares drawn from the plots of per cent $*CO_2$ or C_2H_2 remaining at different breath holding times using the following formula: $*CO_{2t_0}/*CO_{2t}$ or $C_2H_{2t_0}/C_2H_{2t} = e^{Kt}$ where the zero and t subscripts are the per cent of the gas remaining at the start and the end of a time interval and t equals the time interval in seconds.

§ Pulmonary capillary blood flow determined from the rate of disappearance of $*CO_2$ (\dot{Q}^*CO_2) and C_2H_2 ($\dot{Q}_{C_2H_2}$).

|| Calculated DL_{CO_2} in this subject was an imaginary number because \dot{Q}^*CO_2 was greater than $\dot{Q}_{C_2H_2}$.

dissociation curve of the lungs ($V_{tis} + V_c$) is shown in Fig. 3. The middle point on the curve was determined at the alveolar P_{CO_2} present during




FIGURE 2 Graph showing the disappearance of C_2H_2 and $*CO_2$ from the alveolar gas during breath holding at an alveolar P_{CO_2} of approximately 50 mm Hg in subject RWH. The line of least squares for each gas was determined from the four points resulting from four separate breath holding periods. The extension of the lines from the 3 sec points to time zero is an extrapolation. The depressed intercepts at time zero result from the initial uptake of the gases by the pulmonary capillary blood (V_c) and pulmonary tissues (V_{tis}) (see text).

breath holding after inspiring $*CO_2$. Two additional points were obtained by multiple breath holding maneuvers after inspiring air or 12–15% CO_2 (Table V). Fig. 4 shows the CO_2 dissociation curve of V_{tis} alone. The data for this figure was determined by subtracting the volume of CO_2 calculated to be in V_c from the volume in $V_{tis} + V_c$. Above virtual mixed venous P_{CO_2} an increase in alveolar P_{CO_2} of 1 mm Hg resulted in an average movement of 1.5 ml of CO_2 into $V_{tis} + V_c$ or 1.2 ml into V_{tis} in the five subjects. Below virtual mixed venous P_{CO_2} a decrease in alveolar P_{CO_2} of 1 mm Hg was accompanied by a loss of 2.2 ml of CO_2 from $V_{tis} + V_c$ or 1.7 ml from V_{tis} (Tables VI and VII).

If the movement of CO_2 into and out of V_{tis} was limited to CO_2 in physical solution, only about 0.45 ml of CO_2 would be expected to leave or enter V_{tis} for each mm Hg change in alveolar P_{CO_2} .⁴ Since the mean figure for our subjects is two to

⁴ Assuming that the CO_2 solubility in V_{tis} is the same as in water (0.57-ml STPD per ml per atmosphere), the change in CO_2 content for each millimeter change in P_{CO_2} in a subject with V_{tis} equal to 600 ml would be: $(0.57 \times 600)/760$ or 0.45 ml/mm Hg.

TABLE IV

CO₂ Content of Pulmonary Tissue Volume (V_{tis}) and Pulmonary Capillary Blood Volume (V_c) in Five Human Subjects at Virtual Mixed Venous PCO₂

Subject	Alveolar PCO ₂	Volume of V _c + V _{tis}	CO ₂ in V _c + V _{tis}	CO ₂ /V _c + V _{tis}	Volume of V _c *	CO ₂ /V _c	Volume of V _{tis}	CO ₂ in V _{tis}	CO ₂ /V _{tis}
	mm Hg	ml	ml	ml/ml	ml	ml/ml	ml	ml	ml/ml
A. At rest									
RWH	50.0	865	326	0.39	112	0.53‡	753	255	0.34
WFR	47.6	900	262	0.29	110	0.52‡	790	205	0.26
PBK	48.0	542	250	0.46	78	0.52‡	464	209	0.45
RJMP	50.5	597	219	0.37	94	0.50§	503	172	0.34
MAF	43.3	604	161	0.27	71	0.42§	533	151	0.25
Mean:	47.9	702	244	0.36	93	0.50	608	198	0.33
B. During exercise									
RWH	54.1	760	318	0.42	—	0.53§	—	—	—
C. After acetazolamide									
RWH	40.5	942	66	0.07	—	0.48‡	—	—	—

* Determined from measurements of the carbon monoxide diffusing capacity (15).

‡ Estimated from mixed venous PCO₂ and PO₂ (see text).

§ Estimated from the CO₂ content, PCO₂, and PO₂ of a peripheral venous blood corrected to mixed venous PCO₂ and PO₂ with standard monograms (see text).

three times greater than this figure, CO₂ in forms such as bicarbonate or carbamates must make a major contribution to the steepness of the CO₂ dissociation curve.

The slope of the dissociation curve of V_{tis} observed in the five subjects of this study is similar to the values reported by Sackner, Feisal, and

DuBois using a plethysmographic technique (19). In their five subjects mean slope of V_{tis} was 1.32 ml of CO₂ per mm Hg compared to 1.41 ml of CO₂ per mm Hg in the present study. In addition, they made calculations from previous breath hold-




FIGURE 3 CO₂ dissociation curve of V_{tis} + V_c (pulmonary parenchyma tissue volume plus pulmonary capillary blood volume). The middle point on the curve for each subject was obtained by breath holding with ¹³CO₂. The upper and lower points were determined from analysis of alveolar gas samples after inhaling 12-15% CO₂ or 0% CO₂.

TABLE V

Changes in Volume of CO₂ in the Pulmonary Tissues (V_{tis}) and Pulmonary Capillary Blood (V_c) after the Inspiration of Air or 12-15% CO₂

Subject	RV	IV	Inspired PCO ₂	Change in alveolar PCO ₂ *	Change in volume of CO ₂ in V _{tis} + V _c
	ml	ml	mm Hg	mm Hg	ml
RWH	1860	4350	0	51-21	-45
	1630	4310	109	49-88	+37
WFR	1860	4840	0	48-20	-57
	1860	4840	99	48-79	+57
PBK	1450	4000	0	51-24	-74
	1450	4000	98	48-76	+49
RJMP	1000	3500	0	47-22	-64
	1000	3500	96	45-74	+61
MAF	880	3080	0	42-19	-50
	880	3080	100	43-78	+30

Abbreviations: RV, residual volume in ml STPD; IV, the inspired volume in ml STPD.

* The first figure is the alveolar PCO₂ in RV just before inhaling IV. It was determined by the rebreathing method. The second figure is P_{equil}CO₂ which is defined as the PCO₂ in the alveolar gas, V_{tis}, and V_c immediately after inspiration. See Appendix I for method of measurement.




FIGURE 4 CO_2 dissociation curve of the pulmonary parenchyma (V_{tis}). The data for these curves were obtained by subtracting from the total CO_2 in $V_{tis} + V_c$, the amount of CO_2 calculated to be in V_c . CO_2 contained in V_c was determined from the product of V_c measured by the carbon monoxide method (15) and the CO content of the capillary blood (see text).

ing data collected by DuBois and reported by Fenn and Dejours (22). Their technique, which is similar to the breath holding method described in this report, gave an average value of 1.63 ml/mm Hg in nine subjects for the slope at an alveolar P_{CO_2} slightly below virtual mixed venous P_{CO_2} . In our subjects the slope for V_{tis} below virtual mixed venous P_{CO_2} was 1.67 ml/mm Hg.

Measurements during exercise. In one subject the rate of uptake of $^*\text{CO}_2$ and C_2H_2 was determined in the supine position while exercising on a bicycle ergometer. Measurements were made during the 6th min of exercise at a work load of 90 w at which time pulse rate was 120 beats/min, O_2

consumption was 1250 ml/min, and the respiratory quotient was 0.91. Compared to resting measurements in the same subject the initial rapid disappearance of $^*\text{CO}_2$ and C_2H_2 showed little change, but the fall in the concentration of these gases during subsequent breath holding doubled (Tables III B and IV B and Fig. 5). \dot{Q}_c calculated from the rate of disappearance of the two gases was 12.0 and 12.6 liters/min for $^*\text{CO}_2$ and C_2H_2 , respectively. This similarity in blood flows suggested that even a doubling of the cardiac output is not associated with an alveolar to end-capillary CO_2 gradient of any significance.

Effect of body position. Because of the possibility that uneven distribution of ventilation and volume with respect to perfusion (uneven V_A/\dot{Q}_c and uneven V_A/Q_c) might influence the rate of disappearance of $^*\text{CO}_2$, measurements were made in both the sitting and supine positions as well as during exercise (Table III). Neither body position nor exercise were found to produce any consistent differences in the calculated CO_2 content of the lung tissues, relative disappearance rates of $^*\text{CO}_2$ and C_2H_2 , or other parameters. Therefore, the different degrees of uneven V_A/\dot{Q}_c and V_A/Q_c very likely produced by exercise and by varying body position did not appear to alter appreciably the factors determining the rate of disappearance of $^*\text{CO}_2$ from the alveoli.

Measurements after the administration of a carbonic anhydrase inhibitor (acetazolamide). One subject (RWH), over a 15 min interval, received an intravenous infusion of 500 ml of 5% glucose and water containing 7 g of sodium aceta-

TABLE VI
Slope of the CO_2 Dissociation Curve of the Pulmonary Parenchymal Tissues (V_{tis}) Plus the Pulmonary Capillary Blood (V_c)

Subject	ml/mm Hg	ml/mm Hg	ml/mm Hg	ml/mm Hg per m ² of sur- face area	ml/mm Hg per 100 ml of $V_{tis} + V_c$
	(20-50)*	(50-80)*	(20-80)*	(20-80)*	(20-80)*
RWH	1.52	0.95	1.20	0.62	0.14
WFR	2.06	1.84	1.95	0.97	0.22
PBK	2.79	1.75	2.23	1.21	0.41
RJMP	2.54	2.11	2.33	1.23	0.39
MAF	2.18	0.87	1.41	0.87	0.23
Mean	2.22	1.50	1.82	0.98	0.28

* Approximate lower and upper values of P_{CO_2} in mm Hg used for the calculation of the slope of the CO_2 dissociation curve.

TABLE VII
Slope of the CO₂ Dissociation Curve of the Pulmonary Parenchymal Tissues (V_{tis})

Subject	ml/mm Hg	ml/mm Hg	ml/mm Hg	ml/mm Hg per m ² of surface area	ml/mm Hg per 100 ml of V _{tis}
	(20-50)*	(20-80)*	(20-80)*	(20-80)*	(20-80)*
RWH	0.86	0.61	0.72	0.37	0.10
WFR	1.45	1.52	1.49	0.74	0.19
PBK	2.38	1.50	1.90	1.03	0.41
RJMP	2.01	1.79	1.91	1.01	0.38
MAF	1.65	0.64	1.05	0.64	0.21
Mean	1.67	1.21	1.41	0.76	0.26

* Approximate lower and upper values of Pco₂ in mm Hg used for the calculation of the slope of the CO₂ dissociation curve.

zolamide (100 mg/kg).⁵ At the time of the breath holding measurements 4 hr later end-tidal Pco₂ was 14 mm Hg and virtual mixed venous Pco₂ determined by the rebreathing method was 40.5 mm Hg. The subject complained of marked intoxication, difficulty in performing simple arithmetic calculations, severe headache, ataxia, and occasional transient episodes of red, blurred vision lasting 5-15 sec. This severe reaction is in striking contrast to the minor symptomatology reported by Janowitz, Dreiling, Rolbin, and Hollander who gave humans doses as high as 154 mg/kg (27). The difference in symptoms may be related to the rapid infusion of the drug within 15 min in the present study compared to the administration over a 1-8 hr period in the experiments reported by Janowitz and coworkers.

During intoxication with acetazolamide $\dot{Q}_{C_2H_2}$ was 10.9 liters/min and *CO₂ disappeared at a rate of 6.3%/sec so that \dot{Q}^*CO_2 was only 3.01 liters/min. The CO₂ stores in V_{tis} + Vc calculated from the initial disappearance of the isotope were only 69 ml (Tables III C and IV C and Fig. 6). Without enzyme inhibition at this $\dot{Q}_{C_2H_2}$ the subject would be expected to have a *CO₂ disappearance rate of 23%/sec and CO₂ stores in V_{tis} + Vc equal to 313 ml. Since the lower values observed after the administration of acetazolamide are most likely due to the decreased ability of *CO₂ to exchange with the bicarbonate in V_{tis} and Vc, the difference between the above observed and predicted values of CO₂ in V_{tis} + Vc must represent CO₂ in the form of bicarbonate. In this subject

the fraction of total CO₂ in the form of bicarbonate calculated by this method was 79% for V_{tis} and 73% for Vc (Table VIII). Because only the conversion of *CO₂ to bicarbonate inhibited by acetazolamide was taken into account by this calculation, these percentages are minimal values. The above findings support the view that the carbonic anhydrase in lung tissues and blood is necessary in order to have equilibration of the CO₂ stores in V_{tis}, Vc, and alveolar gas during the time it takes blood to traverse the pulmonary capillaries (3, 5).




FIGURE 5 Graph showing the rate of disappearance of C₂H₂ and *CO₂ from the lungs of subject RWH during exercise. Compared to resting values (Fig. 2) there was little change in the zero intercepts but subsequent disappearance was twice as fast.

⁵ Kindly supplied by Dr. R. N. Fallon, Lederle Laboratories as the sodium salt of Diamox acetazolamide.




FIGURE 6 Graph showing the rate of disappearance of C_2H_2 and $*CO_2$ from the lungs of subject RWH after the administration of the carbonic anhydrase inhibitor acetazolamide. The heavy dashed line represents the rate of disappearance of $*CO_2$ predicted on the basis that there was no alveolar to end-capillary gradient or alveolar to pulmonary tissue gradient for $*CO_2$. Since the experimental line had a higher value at zero time and a less steep slope, equilibration between CO_2 in blood, lung tissue, and alveolar gas was not present during enzyme inhibition.

Calculation of hydrogen ion concentration (H^+) in V_{tis} . If the concentration of bicarbonate ($-HCO_3$) and CO_2 in physical solution (αPco_2) in V_{tis} are known, its hydrogen ion concentration $[H^+]$ can be calculated from the relationship $pH = pK' + \log \left(\frac{[-HCO_3]}{\alpha Pco_2} \right)$. $[-HCO_3]$ and total CO_2 in V_{tis} was measured in subject RWH (see above) and in the other subjects $[-HCO_3]$ was assumed to be 80% of the total CO_2 in V_{tis} at the alveolar Pco_2 present during breath holding. αPco_2 was calculated from the formula $\alpha Pco_2 = (\text{alveolar } Pco_2) (V_{tis}) (0.57)/760$ where 0.57 is the Bunsen solubility coefficient for CO_2 in water at $37^\circ C$. pK' was assumed to equal 6.03 (28). If the amount of CO_2 which cannot be assigned to $-HCO_3$ or αPco_2 is assumed not to change with alterations in Pco_2 , $[H^+]$ of lung tissue for any CO_2 tensions above and below virtual mixed venous Pco_2 can be calculated.⁶ Fig. 7 shows the results for the five subjects. At Pco_2 of 25 mm Hg average pH was 7.14 (SD = 0.06), at Pco_2 of 40 mm Hg average pH was 6.92 (SD = 0.09), and

⁶ According to data reported by Roughton (2), CO_2 transported in the form of carbamino-hemoglobin does not show a great change with variation in Pco_2 . We have assumed that the CO_2 -protein complexes in lung tissue behave in the same manner.

TABLE VIII
 CO_2 Transport in Subject RWH at Virtual Mixed Venous $Pco_2 = 40.5$ mm Hg

	CO_2 content of $V_{tis}+Vc^*$		CO_2 content of Vc^*		CO_2 content of V_{tis}^*	
	ml	%	ml	%	ml	%
Total CO_2						
Control	313‡	100	54§	100	259	100
After acetazolamide	66	21	15¶	27	51	20
Bicarbonate**	247	79	39	73	208	80
CO_2 in physical solution‡‡	26	8	3	6	23	9
Undesignated CO_2 §§	40	13	12	21	28	11

* V_{tis} , pulmonary parenchymal tissue volume; Vc , pulmonary capillary blood volume.

‡ Control CO_2 content of $V_{tis}+Vc$ was determined from the initial rapid disappearance of $*CO_2$ during breath holding.

§ Control CO_2 content of Vc was determined from a measurement of Vc by the carbon monoxide method and an estimate of the CO_2 content of the pulmonary capillary blood (see text).

|| Apparent CO_2 content of $V_{tis}+Vc$ calculated from the initial rapid disappearance of $*CO_2$ after acetazolamide administration.

¶ The CO_2 content of Vc after acetazolamide administration was assumed to be reduced from control values in proportion to the decrease in the apparent Bunsen solubility coefficient for total CO_2 in the blood determined from the rate of disappearance of $*CO_2$ and C_2H_2 during breath holding.

** The bicarbonate fraction of the CO_2 content was considered to equal the difference in CO_2 content between measurements made before and after acetazolamide administration.

‡‡ Calculated assuming a solubility coefficient of 0.57 ml of CO_2 STPD per ml per atmosphere.

§§ Volume of CO_2 not assignable to bicarbonate or CO_2 in physical solution.

at Pco_2 of 70 mm Hg equalled 6.74 (SD = 0.11). The pH at Pco_2 of 25 and 40 mm Hg was similar to the values recently reported for intracellular muscle tissue (29), but the value at 70 mm Hg for muscle was 6.91 which is 0.17 pH units higher than the value determined for V_{tis} . This discrepancy is probably not significant in view of the




FIGURE 7 Relationship between PCO_2 and pH of the pulmonary parenchymal tissues (V_{tis}).

variation in the individual values, the fact that the calculations include both intra- and extracellular spaces, and that the change in P_{CO_2} in V_{tis} took place in a few seconds while, in the muscle experiments, the tissues were incubated at constant P_{CO_2} for 4–6 hr.

The diffusing capacity for CO_2 (DL_{CO_2}). In the presence of active carbonic anhydrase, the pulmonary capillary blood flows calculated from the rate of disappearance of *CO_2 and C_2H_2 were almost identical. This finding indicates that on the average there is an extremely small alveolar end-capillary gradient for *CO_2 so that DL_{CO_2} cannot be measured accurately. For example in two subjects (WFR and MAF) the solution for DL_{CO_2} produced an imaginary number because \dot{Q}^*CO_2 was actually greater than $\dot{Q}C_2H_2$. In the other subjects the value at rest was over 200 ml/(min \times mm Hg) but slight errors in the rate of disappearance of *CO_2 or C_2H_2 would produce large changes in DL_{CO_2} .

After the administration of acetazolamide, end-capillary P^*CO_2 calculated by previously described methods (14) was less than one-third of the alveolar P^*CO_2 and the resulting DL_{CO_2} was only 42 ml/(min \times mm Hg). Because of the large alveolar to end-capillary gradient DL_{CO_2} in this special circumstance can be measured with reasonable precision. For example, a 3% error in $\dot{Q}C_2H_2$ produces less than a 1% change in DL_{CO_2} . The reduction in DL_{CO_2} during carbonic anhydrase inhibition is most likely due to a fall in θ_{CO_2} , the diffusing capacity of the blood for CO_2 , which has been shown in vitro to decrease markedly after the administration of acetazolamide. According to this hypothesis, the P^*CO_2 in the blood in the capillary would most certainly have equilibrated with that in the alveolar gas long before the end of the capillary, but would not be in equilibrium with the blood bicarbonate, either within or without the red cell. Using equation 7 the value for θ_{CO_2} which would account for the DL_{CO_2} of 42 ml/(min \times mm Hg) in this subject can be calculated in the following manner: according to equation 7: $1/42 = 1/DM_{CO_2} + 1/(\theta \times 130)$ where 130 represents an estimate of V_c during carbonic anhydrase inhibition in this subject in ml (Tables III and IV). According to Table III DL_{CO_2} , which must be less than DM_{CO_2} , is at least five times greater than 42, if indeed it is not so large it cannot be measured

by this technique at all. Therefore, the term $1/DM_{CO_2}$ becomes so small it can be neglected, giving $\theta = 0.32$ ml/(min \times mm Hg \times ml). This in vivo value is considerably smaller than the in vitro value of 0.6 ml (min \times mm) reported by Constantine, Craw, and Forster (24).

This discrepancy may be related to the fact that a significant part of the initial reaction these authors measured during carbonic anhydrase inhibition was due to the formation of carbamate superimposed on the slower formation of bicarbonate. If θ_{CO_2} is calculated from their data collected after the first 100 msec of the reaction, at which time the carbamate reaction has likely gone to completion, θ_{CO_2} is only 0.24 ml/(min \times mm Hg) per ml, a figure in better agreement with the value measured in vivo. However it is likely that during carbonic anhydrase inhibition θ_{CO_2} varies along the length of the capillary. Large values would be expected at times when carbamate concentrations are changing and small values when only bicarbonate levels are varying. Because θ_{CO_2} may at times be a variable quantity, the present in vivo value as well as the reported in vitro values measured during carbonic anhydrase inhibition should be considered as approximations.

DISCUSSION

Estimate of size of DL_{CO_2} and the alveolar-capillary CO_2 gradients due to diffusion in normal subjects. In these experiments at the start of the capillary after inspiring $^{13}CO_2$, the alveolar $P^{13}CO_2$ was 5–10-fold greater than the $P^{13}CO_2$ in the virtual mixed venous blood. In contrast during resting ventilation the difference between unlabeled alveolar P_{CO_2} and mixed venous P_{CO_2} is only about 20%. Despite this ability of isotopic CO_2 to magnify any difference between alveolar and end-capillary P_{CO_2} , in our subjects no consistent gradient was detected and DL_{CO_2} could not be calculated accurately. Since we estimate that the analytical methods permit the detection of a difference of 5% between alveolar and end-capillary $P^{13}CO_2$, only a minimal value for DL_{CO_2} can be calculated based on the assumption of an alveolar to end-capillary gradient of this size. In the five subjects at rest this minimal value for DL_{CO_2} was 220 (sd = 30) ml/(min \times mm Hg). The minimal value for the diffusing capacity of the pulmonary membrane (DM_{CO_2}) determined from the minimal

value of DL_{CO_2} , a value of θ_{CO_2} equal to 5.1 ml/(min × mm Hg × ml) (24), and equation 7 was 464 (SD = 165) ml/(min × mm Hg). This value is approximately one-third the value for DM_{CO_2} predicted on the basis of measurements of the diffusing capacity of the pulmonary membrane for CO (DM_{CO}), the molecular weights of CO and *CO_2 , and the relative solubilities of CO and CO_2 in water.⁷

Because the technique for calculating a minimal value for DL_{CO_2} required that total alveolar P_{CO_2} (P_{ACO_2}) equal virtual mixed venous P_{CO_2} ($P_{MV_{CO_2}}$), this value of DL_{CO_2} does not directly describe the usual events that take place along the capillary during normal ventilation where P_{ACO_2} is about 7 mm Hg less than $P_{MV_{CO_2}}$. However, once an estimate of DL_{CO_2} has been obtained, the rate of change in capillary P_{CO_2} as blood traverses the pulmonary capillaries during normal physiological conditions can be calculated (Appendix II). This calculation is considerably simpler than is the case for O_2 because the CO_2 dissociation curve for blood is, for all practical purposes, linear over the range of values between $P_{MV_{CO_2}}$ and arterial P_{CO_2} , and θ_{CO_2} can be assumed to be constant over this interval. The slight inaccuracy in this calculation resulting from the difference in diffusion coefficient for CO_2 of mass 45 as against mass 44 is likely to be small compared to errors which may arise from uneven distribution of diffusing capacity and blood flow, the use of a minimal value for DL_{CO_2} , and the Haldane effect which, according to recent *in vivo* measurements, may modify the rate of change of capillary P_{CO_2} presumably because of variations in θ_{CO_2} along the capillary (30).

The unbroken curve in Fig. 8, curve A, was calculated using a minimal value for DL_{CO_2} for a normal subject at rest. This curve, therefore, represents the slowest rate of change in capillary P_{CO_2} compatible with the data. According to this minimal value P_{CO_2} would fall half way to the alveolar P_{CO_2} in 0.9 sec and 90% of the distance in 0.3 sec.

The dashed curve below curve A represents the change in P_{CO_2} calculated from a value of DL_{CO_2}

⁷ If it is assumed that DM_{CO} equals 60 ml/(min × mm Hg), DM_{CO_2} would equal $60 \times (0.592/0.0185)(\sqrt{28/45})$ or 1500 ml/(min × mm Hg). 28 and 45 are the mol wt of CO and *CO_2 , and 0.529 and 0.0185 are their respective solubilities in water.




FIGURE 8 Time course of the fall in capillary P_{CO_2} as blood traverses the pulmonary capillaries. Curve A is a plot for a normal subject at rest calculated using a minimal value for DL_{CO_2} derived from the rate of disappearance of *CO_2 from the alveolar gas. The broken curve below curve A was calculated using a value for DL_{CO_2} derived from the size of the carbon monoxide diffusing capacity (see text). Alveolar and capillary P_{CO_2} are essentially in equilibrium in 0.4 sec. Curve B is a plot for a normal subject after the administration of acetazolamide calculated on the basis that the inhibitor reduced the diffusing capacity for CO_2 from 250 to 40 ml/(min × mm Hg) and alveolar P_{CO_2} remained at 40 mm Hg. Under these conditions an alveolar to end-capillary gradient for CO_2 of 4 mm Hg would be present. Curve C describes the time course along the capillary during carbonic anhydrase inhibition predicted to take place in a subject with a pulmonary disease which had reduced the capillary blood volume and diffusing capacity of the pulmonary membrane to one-third of normal and in addition prevented him from decreasing his alveolar P_{CO_2} below 40 mm Hg. Under these conditions DL_{CO_2} would be only 10.8 ml/(min × mm Hg) and in order to have a sufficient alveolar-capillary CO_2 gradient for elimination of the body's CO_2 production, mixed venous P_{CO_2} would have to increase to 70.5 mm Hg. The alveolar to end-capillary CO_2 gradient would be 23.5 mm Hg. The following dimensions of the capillary bed and lungs representative for our subjects were used in these calculations, namely: alveolar P_{CO_2} = 40 mm Hg; CO_2 production = 288-ml STPD per minute; \dot{Q}_c = 7200 ml/min; slope of CO_2 dissociation curve = 0.0057 ml STPD per milliliter per mm Hg; minimal DL_{CO_2} = 250 ml/(min × mm Hg); DL_{CO_2} calculated from measurements of DL_{CO} = 435 ml/(min × mm Hg); and V_c = 100 ml. θ_{CO_2} was assumed normally to be 5.1 ml/(min × mm Hg) per ml and 0.35 ml/(min × mm Hg) per ml during carbonic anhydrase inhibition. Capillary transit time calculated from the relationship $V_c \div \dot{Q}_c$, equalled 0.83 sec in the normal subject and 0.28 sec in the hypothetical subject with pulmonary disease.

determined on the basis of the following two assumptions. First, according to the molecular weights of CO and *CO_2 and their relative solubilities in water, DM_{CO_2} is 25-fold greater than DM_{CO} . Second, V_c for CO_2 may be larger than V_c for CO but recent measurements of O_2 vs. ether exchanges between the pulmonary vessels and the alveolar gas imply that this difference can be at the most 20% (31). Using these assumptions according to equation 7: $1/DL_{CO_2} = 1/(25 \times 60) + 1/(5.1)(100 + 20)$ or $DL_{CO_2} = 435 \text{ ml}/(\text{min} \times \text{mm Hg})$ where 60 is a representative value for DM_{CO} in normal subjects (32), 5.1 is the value for θ_{CO_2} (24), and 100 is a representative figure for V_c in our subjects measured with the CO technique. According to this estimate of DL_{CO} P_{CO_2} would fall half way to the arterial P_{CO_2} in 0.06 sec and 90% of the distance in 0.18 sec. Since the mean capillary transit time for the pulmonary capillaries is approximately 0.8 sec, this data indicates that normally using either method of estimating DL_{CO_2} , P_{ACO_2} , and P_{CO_2} are in equilibrium by the time the blood reaches the end of the capillaries. However, if some of the capillaries have shorter transit times due to uneven distribution of blood flow, or if the mean transit time is decreased by severe exercise or lung disease, it is likely that an alveolar to end-capillary CO_2 gradient of approximately 1–2 mm Hg would result. Such a gradient would be difficult to detect with present methods of measuring alveolar and arterial P_{CO_2} and, moreover, would not cause hypercapnea because even a slight increase in alveolar ventilation can readily compensate for so small a rise in end-capillary P_{CO_2} .

After the administration of acetazolamide, carbon dioxide exchange along the capillary was dramatically altered and because a large alveolar to end-capillary *CO_2 gradient developed, DL_{CO_2} could be measured with reasonable precision (see above) (Fig. 9).

The subject compensated for the decrease in DL_{CO_2} by vigorous hyperventilation so that the alveolar P_{CO_2} fell to approximately 14 mm Hg and the gradient between $P_{MV_{CO_2}}$ and P_{ACO_2} increased from 7 to 26 mm Hg. The calculated decrease in the capillary P_{CO_2} as blood traverses the capillaries, assuming chemical equilibrium between the dissolved CO_2 and bicarbonate, was 12 mm Hg. The difference between alveolar P_{CO_2} and




FIGURE 9 Time course of the fall in capillary P_{CO_2} as blood traverses the pulmonary capillaries during carbonic anhydrase inhibition in subject RWH at which time his DL_{CO} had fallen to 42 ml/(min \times mm Hg). Note the large alveolar to end-capillary CO_2 gradient of 14 mm Hg. The P_{CO_2} of 40 mm Hg at the start of the capillary was measured with the rebreathing technique (10). Alveolar P_{CO_2} of 14 mm Hg is an approximation on the basis of an observed end-tidal P_{CO_2} varying between 11 and 15 mm Hg. Slope of the CO_2 dissociation curve was assumed to be 0.006-ml STPD per mm Hg (16), and V_c was assumed to have increased from the resting value of 112–130 ml on the basis of the measured increase in carbon monoxide diffusing capacity during carbonic anhydrase inhibition. Pulmonary capillary transit calculated from the relationship $V_c \div Q_c$ equalled 0.715 sec.

equilibrated end-capillary P_{CO_2} was 14 mm Hg. The hyperventilation was so intense that, despite the development of a much less efficient alveolar-capillary CO_2 exchange, virtual mixed venous P_{CO_2} fell from 50 to 40 mm Hg. This hyperventilation was probably secondary to intracellular hypercarbia and acidosis in the cells of the respiratory centers (33).

Figure 8 curve B shows the changes in capillary P_{CO_2} which would have taken place in a normal subject if acetazolamide administration had not been accompanied by a decrease in alveolar P_{CO_2} . In order to have a sufficient alveolar-capillary pressure gradient to eliminate the body's production of CO_2 , the arterial P_{CO_2} would have increased from 40 to 44 mm Hg and the end-capillary to alveolar CO_2 gradient would have been 4 mm Hg.

Effect of carbonic anhydrase inhibition on patients with compromised lung function. Carbonic anhydrase inhibitors have been advocated for use in patients with chronic obstructive lung disease in

order to improve arterial blood oxygenation and lower the arterial P_{CO_2} (34, 35). The rationale of this therapy has been disputed (36). In order to evaluate the effect of carbonic anhydrase inhibition in CO_2 elimination in the presence of severe pulmonary disease, we calculated the capillary P_{CO_2} for a hypothetical subject who was unable to lower his alveolar P_{CO_2} below 40 mm Hg, and who, in addition, had values for DM_{CO_2} and V_c one-third of normal (Figure 8 curve C). Pulmonary capillary blood flow does not usually decrease strikingly in pulmonary disease so that the capillary transit time equal to V_c/\dot{Q}_c is considerably reduced. Under these conditions during severe carbonic anhydrase inhibition a steady state for CO_2 elimination would not be reached until the end-capillary P_{CO_2} had risen from 40.3 to 63 mm Hg, a rise large enough to have significant physiological and clinical consequences. Since some patients with pulmonary disease cannot hyperventilate in order to compensate for the less efficient pulmonary elimination of CO_2 secondary to carbonic anhydrase inhibition, it is not surprising that occasionally symptoms suggestive of increased acidosis and clinical deterioration may accompany therapy with these drugs (35, case 10, 37). However, clinically much smaller doses of acetazolamide are administered than were used in this study so that the increase in the alveolar to end-capillary CO_2 gradient in patients is likely to be negligible or at least considerably smaller than the large gradient calculated to have developed in this hypothetical patient. While carbonic anhydrase inhibitors may cause hyperventilation and thereby lower the arterial and alveolar P_{CO_2} in many patients, even in normal subjects modest amounts of these agents are known to impair CO_2 elimination (38).

Validity of the measurement of hydrogen ion concentration in lung tissue. Intracellular electrodes, indicators, and the dissociation of weak bases and acids (39–41) have all been used to measure intracellular pH. In this study the lung pH was measured by the third technique using the bicarbonate and carbonic acid system. This method, which was first used by Warburg in 1922 (41) has given discordant values because of the difficulty in measuring intracellular bicarbonate levels. For example, in rat muscle the acid-labile CO_2 fraction is almost twice as large as the barium

soluble fraction, which suggests that significant amounts of intracellular CO_2 may be present in nonbicarbonate fractions such as CO_2 protein complexes (42). In the present study we measured the total CO_2 in the lungs that exchanges with alveolar gas in a second or less before and after carbonic anhydrase inhibition and assumed that the difference must be the total bicarbonate in the lungs.

This method has several limitations and sources of error. First, the method is only applicable to a tissue such as the lungs where very rapid exchange of labeled CO_2 facilitated by carbonic anhydrase takes place (4). Second, if carbonic anhydrase inhibition is incomplete, the bicarbonate fraction will be underestimated. Fortunately in this study, 80% of the CO_2 exchange was blocked by the inhibitor, and since 9% of the CO_2 exchange could be assigned to dissolved CO_2 , the error due to incomplete inhibition could not exceed 11% of the total CO_2 space. Third, carbonic anhydrase inhibition may produce a change in the bicarbonate and intracellular nonbicarbonate bound CO_2 . We have no way to evaluate this source of error except to point out that several hours after the administration of a carbonic anhydrase inhibitor, the CO_2 content of blood changes only slightly (7). Fourth, the toxic effects during carbonic anhydrase inhibition were severe enough in one human to make us reluctant to administer large doses of the inhibitor to others, and, therefore, we have data on only one subject. Fifth, evidence has been recently presented in vitro that carbonic anhydrase, in addition to catalyzing the conversion of CO_2 to bicarbonate, directly facilitates the transport of molecular CO_2 (43, 44). If this action of carbonic anhydrase is of significance in vivo, the CO_2 exchange space in the lungs blocked by carbonic anhydrase inhibition might be greater than the bicarbonate space. This would produce an underestimation of the hydrogen ion concentration. However, diffusion distances in the lung parenchyma are generally so small, that a carrier enzyme could not appreciably increase the size of the CO_2 space. Sixth, the concentration of CO_2 in the lungs that is neither bicarbonate nor dissolved CO_2 may vary widely among individuals and may be influenced by acetazolamide.

Since none of the above limitations to this

method of determining tissue pH appear insurmountable, we believe it deserves further evaluation and refinement in the experimental animal and isolated tissue preparations.

Dimensions and dynamics of CO₂ stores in the lungs of man. The data presented above support the findings of a number of investigators (3-5), that the various forms of carbon dioxide present in the lungs exchange so rapidly, that the alveolar Pco₂ can be considered to come from one common CO₂ pool. Representative figures for an adult human with a resting lung volume of 3000-ml STPD at an alveolar Pco₂ of 40 mm Hg are: 150 ml of CO₂ in the alveolar gas, 200 ml of CO₂ in the pulmonary tissue (V_{tis}), and 50 ml of CO₂ in the capillary blood (Vc). At least 70% of the CO₂ in Vc and V_{tis} is in the form of bicarbonate. In the subjects in this study a change in alveolar Pco₂ of 1 mm Hg was accompanied by the movement of about 2 ml of CO₂ into or out of V_{tis} and Vc.

Calculations made by DuBois, Britt, and Fenn (45) have demonstrated the considerable ability of the CO₂ stores in V_{tis} and Vc to buffer changes in alveolar Pco₂. They showed that this buffering action of V_{tis} and Vc reduces the predicted variation in alveolar Pco₂ during the respiratory cycle in resting man from about 3 to 2 mm Hg. The CO₂ stores in V_{tis} and Vc must also buffer oscillations in alveolar Pco₂ secondary to the pulsatile nature of the pulmonary capillary blood flow. We estimate that this change in our subjects is in the order of 0.13 mm Hg between diastole and systole. Without this buffering capacity of V_{tis} and Vc this figure would be 0.18 mm Hg. Bosman, Lee, and Marshall have also presented evidence for this buffering mechanism for CO₂ during the cardiac cycle from measurements of instantaneous carbon dioxide exchange in the lungs using a body plethysmograph (30).

Individual variation of the CO₂ content and shape of the CO₂ dissociation curve of lung tissue. The five subjects in this study showed considerable differences in the CO₂ content of their lung tissue and the slope of its CO₂ dissociation curve which could not be explained by error in the method or difference in body size. For example the CO₂ content of V_{tis} + Vc at an alveolar Pco₂ of 45 mm Hg varied from 104 ml/m² of surface area in subject MAF to 164 ml/m² in RWH. This is a

determination with an experimental error which we believe is less than 20% of the measurement. Other investigators using different techniques have also noted similar variations in these quantities (19, 20). Therefore, the pulmonary intracellular CO₂ concentration most likely has a wider range of normal values than observed in plasma.

Validity of calculating CO₂ content of pulmonary capillary blood from alveolar Pco₂ during breath holding. Recently several investigators have reported that virtual mixed venous Pco₂ obtained by rebreathing is greater than the mixed venous Pco₂ measured from samples of pulmonary arterial blood even when pulmonary arterial and alveolar Po₂ are the same (47, 48). The effect of this interesting finding on the pulmonary capillary CO₂ content which was required in this study for the calculation of \dot{Q}_{CO_2} and the alveolar end-capillary CO₂ gradient, has not been clarified at the present time. Fortunately the shape of the CO₂ dissociation curve for blood is such that changes in Pco₂ produce proportionally less changes in CO₂ content. For example a 10% elevation in the rebreathing Pco₂, which is the approximate magnitude suggested by these reports, will at most produce an overestimation of capillary CO₂ content of 4% and would give the false impression that alveolar P*co₂ was 4% greater than end-capillary P*co₂. This source of error is therefore small enough not to alter the conclusions reached in this study.

APPENDIX I

Method for the determination of the slope of the CO₂ dissociation curve of the pulmonary tissues. The subject rebreathed several breaths from a 2 liter bag containing 8% CO₂ in order to make alveolar Pco₂ equal to virtual mixed venous Pco₂ (PMVCO₂). He then maximally inspired a breath of room air, held his breath for 3 sec, and then delivered an alveolar gas sample. Pco₂ of the expired gas sample was determined from the output of the continuously recording mass spectrometer. Every 15 min the procedure was repeated until three to six determinations of Pco₂ had been made with time of breath holding varying from 2 to 14 sec. During breath holding CO₂ will leave V_{tis} and Vc, enter the alveoli, and a new equilibrium between Pco₂ in the alveolar gas, Vc and V_{tis} will be reached at a Pco₂ lower than PMVCO₂ (21, 22). This equilibration takes place in less than 1-2 sec and the amount of CO₂ entering the alveoli will be a function of the CO₂ dissociation curves of V_{tis} and Vc (5, 19). The CO₂ contained in the residual volume (RV), V_{tis}, and Vc before inspiration plus any CO₂ contained in the inspired gas will equal the CO₂ contained

in the alveolar volume, V_{tis} and V_c immediately after inspiration or

$$\frac{PMV_{CO_2}}{P_B - P_{H_2O}}(RV) + \text{CO}_2 \text{ content of } V_c + V_{tis} \text{ at } PMV_{CO_2} \\ + \frac{P_{inspCO_2}}{P_B - P_{H_2O}}(IV - DS) = \frac{P_{equilCO_2}}{P_B - P_{H_2O}}(RV + IV - DS) \\ + \text{CO}_2 \text{ Content of } V_c + V_{tis} \text{ at } P_{equilCO_2}$$

where RV and IV are respectively the residual volume and the inspired volume in ml STPD, $P_{equilCO_2}$ is the partial pressure of CO_2 in the alveoli just after inspiration in mm Hg, and P_{inspCO_2} is the partial pressure of CO_2 in mm Hg, if any, in IV and DS is the respiratory dead space in milliliters STPD. DS in milliliter BTPS was assumed to equal the subject's ideal weight in pounds (46). If $P_{equilCO_2}$ can be measured, equation 7 can be solved for the CO_2 content of $V_{tis} + V_c$ at $P_{equilCO_2}$.

$P_{equilCO_2}$ was determined in the following manner: the difference between the PCO_2 of the expired gas samples and mixed venous PCO_2 was plotted against time of breath holding on semilog paper and the line drawn through the points was extrapolated back to zero time. Plotting the points in this way should theoretically, and in practice did, produce a straight line function (21, 22) (Fig. 10). The PCO_2 at the extrapolated zero time under these conditions represented $P_{equilCO_2}$. The same procedure was repeated for each subject using an inspired gas containing 12–15% CO_2




FIGURE 10 Graph of data used to determine $P_{equilCO_2}$ from the rate of change of alveolar PCO_2 during breath holding after inhaling 12% CO_2 (upper line) and 0% CO_2 (lower line) in subject WFR. Alveolar gas samples were collected after breath holding periods varying from 2.4 to 11.4 sec. The difference between alveolar PCO_2 and PMV_{CO_2} was plotted against time of breath holding on semilog paper and a line drawn through the points was extrapolated back to the start of breath holding. In this subject after inspiring 12% CO_2 $P_{equilCO_2}$ minus the PMV_{CO_2} of 48 mm Hg was 31 mm Hg so that $P_{equilCO_2}$ equalled 48 plus 31 or 79 mm Hg. After inspiring air (lower line in the figure) the difference was 28 mm Hg so that $P_{equilCO_2}$ equalled 48 minus 28 or 20 mm Hg.

so that a point on the CO_2 dissociation curve of $V_{tis} + V_c$ could be determined at a PCO_2 greater than PMV_{CO_2} .

APPENDIX II

Method for determining the changes in intracapillary PCO_2 as blood flows through the pulmonary gas exchange vessels. If it is assumed that: (a) alveolar PCO_2 (PA_{CO_2}) is constant during the capillary transit time; (b) diffusing capacity for CO_2 (DL_{CO_2}) is evenly distributed along the capillary; (c) DL_{CO_2} and pulmonary capillary blood flow are evenly distributed; and (d) the diffusing capacity for CO_2 within the blood (θ_{CO_2}) is constant, then:

$$(P_{CCO_2} - PA_{CO_2})DL_{CO_2} = \frac{d}{dt} [CO_2]V_c \quad (1)$$

where P_{CCO_2} equals the intracapillary PCO_2 at any instant in mm Hg, $[CO_2]$ is the CO_2 content of 1 ml of blood in milliliter STPD and V_c is the pulmonary capillary blood volume in milliliter. DL_{CO_2} is expressed in ml/(min \times mm Hg). Since the CO_2 dissociation curve is almost linear over the interval between mixed venous PCO_2 and end-capillary PCO_2 , $[CO_2]$ can be said to equal $(K)P_{CCO_2} + K_1$ where K is the slope of the CO_2 dissociation curve in milliliter of CO_2 per milliliter of blood per mm Hg change in P_{CCO_2} and K_1 is a constant which will disappear during differentiation. Substituting this expression into equation 1 and integrating gives:

$$\frac{P_{CCO_2_0} - PA_{CO_2}}{P_{CCO_2_t} - PA_{CO_2}} = e^{\frac{t DL_{CO_2}}{(K)V_c(60)}} \quad (2)$$

where t equals the time along the capillary in seconds, $P_{CCO_2_0}$ equals the intracapillary PCO_2 at the beginning the capillary, and $P_{CCO_2_t}$ equals the P_{CCO_2} at time t .

ACKNOWLEDGMENTS

The authors are indebted to Doctor Arthur B. DuBois for critical reading of the manuscript.

This work was supported in part by research grant HE-10324 and HE-4108 from the National Heart Institute, National Institutes of Health, U. S. Public Health Service.

REFERENCES

- Hyde, R. W., and R. E. Forster, II. 1962. Rate of disappearance of a stable CO_2 isotope ($C^{13}O_2$) from the alveolar gas during breathholding. *Physiologist*. 5: 159. (Abstr.)
- Roughton, F. J. W. 1964. Transport of oxygen and carbon dioxide. In *Handbook of Physiology*. American Physiological Society, Washington, D. C. 1: 767.
- Chinard, F. P., T. Enns, and M. F. Nolan. 1960. Contributions of bicarbonate ion and of dissolved CO_2 to expired CO_2 in dogs. *Am. J. Physiol.* 198: 78.
- Chinard, F. P. 1966. The permeability characteristic of the pulmonary blood-gas barrier. In *Advances in Respiratory Physiology*. C. G. Caro, editor. The Williams & Wilkins Co., Baltimore. 106.

5. Feisal, K. A., M. A. Sackner, and A. B. DuBois. 1963. Comparison between the time available and the time required for CO₂ equilibration in the lung. *J. Clin. Invest.* **42**: 24.
6. Cain, S. M., and A. B. Otis. 1961. Carbon dioxide transport in anesthetized dogs during inhibition of carbonic anhydrase. *J. Appl. Physiol.* **16**: 1023.
7. Carter, E. T., and R. T. Clark, Jr. 1958. Effects of carbonic anhydrase inhibition during acute hypoxia. *J. Appl. Physiol.* **13**: 47.
8. Mithoefer, J. C. 1959. Inhibition of carbonic anhydrase: its effect on carbon dioxide elimination by the lungs. *J. Appl. Physiol.* **14**: 109.
9. Cander, L., and R. E. Forster. 1959. Determination of pulmonary parenchymal tissue volume and pulmonary capillary blood flow in man. *J. Appl. Physiol.* **14**: 541.
10. Hackney, J. D., C. H. Sears, and C. R. Collier. 1958. Estimation of arterial CO₂ tension by rebreathing technique. *J. Appl. Physiol.* **12**: 425.
11. Nier, A. O. 1960. A redetermination of the relative abundances of the isotopes of carbon, nitrogen, oxygen, argon, and potassium. *Physical Rev.* **77**: 789.
12. Bates, D. V., and R. V. Christie. 1950. Intrapulmonary mixing of helium in health and emphysema. *Clin. Sci.* **9**: 17.
13. Ogilvie, C. M., R. E. Forster, W. S. Blakemore, and J. W. Morton. 1957. A standardized breath holding technique for the clinical measurement of the diffusing capacity of the lung for carbon monoxide. *J. Clin. Invest.* **36**: 1.
14. Hyde, R. W., R. E. Forster, G. G. Power, J. Nairn, and R. Rynes. 1966. Measurement of O₂ diffusing capacity of the lungs with a stable O₂ isotope. *J. Clin. Invest.* **45**: 1178.
15. Roughton, F. J. W., and R. E. Forster. 1957. Relative importance of diffusion and chemical reaction rates in determining rate of exchange of gases in the human lung, with special reference to true diffusing capacity of pulmonary membrane and volume of blood in the lung capillaries. *J. Appl. Physiol.* **11**: 290.
16. Peters, J. P., and D. Van Slyke. 1933. Quantitative Clinical Chemistry. The Williams & Wilkins Co., Baltimore. 907.
17. Anderson, O. S., E. K. Engel, K. Jorgensen, and P. Astrup. 1960. A micro method for determination of pH, carbon dioxide tension, base excess, and standard bicarbonate in capillary blood. *Scand. J. Clin. Lab. Invest.* **12**: 172.
18. Bartels, H., E. Bucherl, C. W. Hertz, G. Rodewald, and M. Schwols. 1963. Methods in pulmonary physiology. Hafner Publishing Co., New York. 304.
19. Sackner, M. A., K. A. Feisal, and A. B. DuBois. 1964. Determination of tissue volume and carbon dioxide dissociation slope of the lungs in man. *J. Appl. Physiol.* **19**: 374.
20. DuBois, A. B., W. O. Fenn, and A. G. Britt. 1952. CO₂ dissociation curve of lung tissue. *J. Appl. Physiol.* **5**: 13.
21. DuBois, A. B. 1952. Alveolar CO₂ and O₂ during breath holding, expiration, and inspiration. *J. Appl. Physiol.* **5**: 1.
22. Fenn, W. O., and P. Dejours. 1954. Composition of alveolar air during breath holding with and without prior inhalation of oxygen and carbon dioxide. *J. Appl. Physiol.* **7**: 313.
23. Soni, J., K. A. Feisal, and A. B. DuBois. 1963. The rate of intrapulmonary blood gas exchange in living animals. *J. Clin. Invest.* **42**: 16.
24. Constantine, H. P., M. R. Craw, and R. E. Forster. 1965. Rate of the reaction of carbon dioxide with human red blood cells. *Am. J. Physiol.* **208**: 801.
25. West, J. B., and C. T. Dollery. 1960. Distribution of blood flow and ventilation-perfusion ratios in the lung, measured with radioactive CO₂. *J. Appl. Physiol.* **15**: 405.
26. Rahn, H. 1962. The gas stores of the body, with particular reference to carbon dioxide. In *Man's Dependence on the Earthly Atmosphere*. K. E. Schaeffer, editor. The Macmillan Company, New York. 297.
27. Janowitz, H. D., D. A. Dreiling, H. L. Rolbin, and F. Hollander. 1957. Inhibition of the formation of hydrochloric acid in the human stomach by diamox; the role of carbonic anhydrase in gastric secretion. *Gastroenterology.* **33**: 378.
28. Dill, D. B., C. Daly, and W. H. Forbes. 1937. The pK' of serum and red cells. *J. Biol. Chem.* **117**: 569.
29. Adler, S., A. Roy, and A. S. Relman. 1965. Intracellular acid-base regulations. I. The response of muscle cells to changes in CO₂ tension or extracellular bicarbonate concentration. *J. Clin. Invest.* **44**: 8.
30. Bosman, A. R., G. de J. Lee, and R. Marshall. 1965. The effect of pulsatile capillary blood flow upon gas exchange within the lung of man. *Clin. Sci.* **28**: 295.
31. Sackner, M. A., K. A. Feisal, and D. N. Karsch. 1964. Size of gas exchange vessels in the lung. *J. Clin. Invest.* **43**: 1847.
32. Hyde, R. W., R. Rynes, G. G. Power, and J. Nairn. 1967. Determination of distribution of diffusing capacity in relation to blood flow in the human lung. *J. Clin. Invest.* **46**: 463.
33. Mithoefer, J. C., and J. S. Davis. 1958. Inhibition of carbonic anhydrase: effect on tissue gas tensions in the rat. *Proc. Soc. Exptl. Biol. Med.* **98**: 797.
34. Naimark, A., D. M. Brodovsky, and R. M. Cherniack. 1960. The effect of a new carbonic anhydrase inhibitor (dichlorphenamide) in respiratory insufficiency. *Am. J. Med.* **28**: 368.
35. Christensen, P. J. 1962. The carbonic anhydrase inhibitor dichlorphenamide in chronic pulmonary emphysema. *Lancet.* **1**: 881.
36. Dorris, R., J. V. Olivia, and T. Rodman. 1964. Dichlorphenamide, a potent carbonic anhydrase inhibitor. *Am. J. Med.* **36**: 79.
37. McNicol, M. W., and N. B. Pride. 1961. Dichlorphenamide in chronic respiratory failure. *Lancet.* **1**: 906.

38. Stromme, J. H., and J. Fog. 1962. Effect of acetazolamide on respiratory gas exchange during hyperventilation in man. *J. Appl. Physiol.* 17: 6.
39. Carter, N. W., F. C. Rector, Jr., D. S. Campion, and D. W. Seldin. 1967. Measurement of intracellular pH of skeletal muscle with pH-sensitive glass microelectrodes. *J. Clin. Invest.* 46: 920.
40. Caldwell, P. C. 1956. Intracellular pH. *Intern. Rev. Cytol.* 5: 229.
41. Warburg, E. J. 1922. Theory of the Henderson-Hasselbalch equation. *Biochem. J.* 16: 153.
42. Conway, E. J., and P. J. Fearon. 1944. The acid-labile CO₂ in mammalian muscle and the pH of the muscle fibre. *J. Physiol.* 103: 274.
43. Longmuir, I. S., R. E. Forster, and C.-Y. Woo. 1966. Diffusion of carbon dioxide through thin layers of solution. *Nature.* 209: 393.
44. Enns, T. 1967. Facilitation by carbonic anhydrase of carbon dioxide transport. *Science.* 155: 44.
45. DuBois, A. B., A. G. Britt, and W. O. Fenn. 1952. Alveolar CO₂ during the respiratory cycle. *J. Appl. Physiol.* 4: 535.
46. Comroe, J. H., Jr., R. E. Forster, II, A. B. DuBois, W. A. Briscoe, and E. Carlsen. 1962. *The Lung*. Yearbook Medical Publishers Inc., Chicago. 2nd edition. 39.
47. Jones, N. L., E. J. M. Campbell, G. J. R. McHardy, B. E. Higgs, and M. Clode. 1967. The estimation of carbon dioxide pressure of mixed venous blood during exercise. *Clin. Sci.* 32: 311.
48. Gurtner, G. H., S. H. Song, and L. E. Farhi. 1967. Alveolar-to-mixed venous Pco₂ difference during rebreathing. *Physiologist.* 10: 190 (Abstr.)