

## Aggregation of $\gamma$ G3 proteins: relevance to the hyperviscosity syndrome

J. D. Capra, H. G. Kunkel

*J Clin Invest.* 1970;49(3):610-621. <https://doi.org/10.1172/JCI106272>.

### Research Article

Studies on the sera and isolated proteins from 14 patients with  $\gamma$ G3 multiple myeloma revealed a concentration- and temperature-dependent aggregation which was not encountered in 26 sera from patients with multiple myeloma involving other  $\gamma$ G subgroups. When the  $\gamma$ G3 myeloma sera were diluted and characterized by analytical ultracentrifugation, complex formation was minimal. However, when these sera were examined undiluted, marked complex formation was observed. Studies on the isolated proteins, their enzymatic fragments, as well as their heavy and light polypeptide chains localized the aggregating sites to the Fd fragment of the heavy chains. The findings were also documented by acrylamide-gel electrophoresis and capillary tube viscometry.

The hyperviscosity syndrome was observed in six patients: three with  $\gamma$ G3 myeloma and three with  $\gamma$ G1 myeloma. In the latter group extreme protein concentrations appeared essential for the development of the clinical symptoms. The  $\gamma$ G3 cases, however, because of the aggregation phenomenon, showed the syndrome at relatively low protein concentrations.

**Find the latest version:**

<https://jci.me/106272/pdf>



# Aggregation of $\gamma$ G3 Proteins: Relevance to the Hyperviscosity Syndrome

J. D. CAPRA and H. G. KUNKEL

From The Rockefeller University, New York 10021

**ABSTRACT** Studies on the sera and isolated proteins from 14 patients with  $\gamma$ G3 multiple myeloma revealed a concentration- and temperature-dependent aggregation which was not encountered in 26 sera from patients with multiple myeloma involving other  $\gamma$ G subgroups. When the  $\gamma$ G3 myeloma sera were diluted and characterized by analytical ultracentrifugation, complex formation was minimal. However, when these sera were examined undiluted, marked complex formation was observed. Studies on the isolated proteins, their enzymatic fragments, as well as their heavy and light polypeptide chains localized the aggregating sites to the Fd fragment of the heavy chains. The findings were also documented by acrylamide-gel electrophoresis and capillary tube viscometry.

The hyperviscosity syndrome was observed in six patients: three with  $\gamma$ G3 myeloma and three with  $\gamma$ G1 myeloma. In the latter group extreme protein concentrations appeared essential for the development of the clinical symptoms. The  $\gamma$ G3 cases, however, because of the aggregation phenomenon, showed the syndrome at relatively low protein concentrations.

## INTRODUCTION

Previous studies have documented the existence of at least four subgroups of  $\gamma$ G globulin based on antigenic differences in their heavy polypeptide chains (1, 2). While the  $\gamma$ G3 ( $V_i\gamma_{2c}$ ) subgroup accounts for only 8% of the normal  $\gamma$ G globulin and a similar percentage of  $\gamma$ G myeloma proteins (3), several lines of evidence suggest that this subgroup plays a significant role in human immunology. It is known that  $\gamma$ G3 proteins are catabolized at a faster rate than any other  $\gamma$ G globulins (4, 5) and the quantitative level of  $\gamma$ G3 in serum is influenced by the specific genetic makeup of the individual (3). Patients with hypogammaglobulinemia have been shown to have a disproportionate increase in  $\gamma$ G3 globulin (6). Cryoglobulins of the  $\gamma$ G type with rheumatoid factor

activity are predominately  $\gamma$ G3 proteins (7) and preliminary evidence suggests that  $\gamma$ G3 proteins are important in the binding of complement.<sup>1</sup>

An interesting physical property of  $\gamma$ G3 proteins of both normal sera and myeloma sera has been their partial adsorption near the well of origin in immunoelectrophoresis (3). Even isolated  $\gamma$ G3 proteins that have been purified frequently show a characteristic biphasic pattern in the agar system. In the present study this phenomena has been further investigated and a concentration and temperature-dependent aggregation of these proteins has been delineated. This aggregation apparently plays a major role in the hyperviscosity syndrome which occasionally accompanies multiple myeloma (8, 9) and may be the explanation for the rapid catabolism and special complement binding of  $\gamma$ G3 proteins.

Aggregates with S values of 9–15S are frequently encountered in  $\gamma$ A myeloma and are due to S-S bridges (10). Presumably these stable high molecular weight aggregates are responsible for the occasional occurrence of the hyperviscosity syndrome in  $\gamma$ A myeloma (11). Smith, Kochwa, and Wasserman (12) were the first to demonstrate aggregate formation in hyperviscosity syndrome in  $\gamma$ G myeloma. These aggregates were isolated and found to be acid dissociable indicating that the intermolecular bonds were noncovalent. Since their report, however, other workers have failed to document aggregate formation in the hyperviscosity syndrome in  $\gamma$ G myeloma despite analytical ultracentrifugation (13–15).

## METHODS

*Isolation, purification, and characterization of myeloma proteins.* Myeloma proteins were isolated by zone electrophoresis on polyvinyl copolymer (Pevikon). Further purification was accomplished by gel filtration on Sephadex G-200 equilibrated with 0.1 M tris(hydroxymethyl)amino-methane (Tris) and 0.5 M sodium chloride. The isolated proteins were concentrated by vacuum dialysis or by dialysis against polyethylene glycol. For determination of molecular weight by low speed analytical ultracentrifugation an addi-

Received for publication 27 August 1969.

<sup>1</sup> Müller-Eberhard. 1968. *Advan. Immunol.* 8: 1.

tional purification step was utilized. After Sephadex G-200 chromatography each protein was concentrated by vacuum dialysis in membranes which will exclude material of less than 70,000 molecular weight. Each preparation was then centrifuged at 35,000 rpm in a Spinco model L preparative ultracentrifuge for 2 hr at 4°C in a swinging bucket rotor. The top one-third was aspirated and used for molecular weight measurements by the field relaxation technique.  $\gamma$ G subgroup analysis and Gm typing were done as described previously (1, 16).

**Enzymatic cleavage of  $\gamma$ -globulin.** Fab and Fc fragments of myeloma proteins were prepared by papain digestion (17) using a protein:enzyme ratio of 50:1.  $\gamma$ G3 proteins were subjected to enzymatic digestion for 30 min at 37°C in the presence of 0.01 M cysteine and 0.002 M EDTA at pH 7.4.  $\gamma$ G1 proteins were digested for 16 hr. The reaction was stopped by dialysis against cold saline, and Fab and Fc were separated by zone electrophoresis on Pevikon. They were then concentrated by vacuum dialysis and further purified by gel filtration by Sephadex G-100 chromatography. Enzymatic fragments were tested for purity by immunoelectrophoresis and Ouchterlony analysis with appropriate antisera.

Pepsin digestion was accomplished in pH 4.0 acetate buffer with a protein:enzyme ratio of 50:1. Digestion proceeded at 37°C for 24 hr and was terminated by dialysis at 4°C against pH 7.2 phosphate-buffered saline (PBS). The 5S pepsin (Fab')<sub>2</sub> fragment was then isolated by Sephadex G-200 chromatography and purity was ascertained by ultracentrifugation, immunoelectrophoresis, and acrylamide-gel electrophoresis.

**Preparation of H and L chains.** Isolated myeloma proteins at 10 mg/ml were dialyzed into 0.5 M Tris buffer pH 8.6. The buffer was then made 0.1 M in  $\beta$ -mercaptoethanol and dialysis was continued overnight at 4°C. The buffer was then made 0.15 M in iodoacetamide and dialysis was continued for 24 hr at 4°C. Chain separation was accomplished by dialyzing against 1 M propionic acid at 4°C for 72 hr. Separation of H and L chains was performed by gel filtration on a Sephadex G-200 column in 1 M propionic acid and 0.001 M iodoacetamide. Chains were concentrated by dialysis against polyethylene glycol in 0.5 M propionic acid. For some experiments utilizing acrylamide-gel electrophoresis whole myeloma proteins were reduced and alkylated in small volumes and placed directly on the gels. Chain separation in these instances was accomplished by the gels themselves. For acid-urea gels, urea was added to the reduced and alkylated proteins as it was to all samples before placing them on the gels.

**Radioiodination.** 50 mg of isolated myeloma proteins G. J. and E. T. were separately dissolved in 10 cc of 0.05 M phosphate buffer pH 7.0. A modification of the chloramine T method of McConahey and Dixon (18) was used to label G. J. with <sup>125</sup>I and E. T. with <sup>125</sup>I. 5  $\mu$ g chloramine T per mg protein and 5  $\mu$ g sodium metabisulfite per mg protein was used. Over 97% of the radioactivity was precipitated with 10% trichloroacetic acid. Radioactivity was measured on a dual channel scintillation counter (Nuclear-Chicago Automatic Gamma Counter, model 4233). After radioiodination the labeled proteins were purified on a Sephadex G-200 column in order to remove higher molecular weight aggregates.

**Recombination of myeloma proteins.** Recombination and reoxidation of the labeled myeloma proteins was accomplished by a modification of the method of Olins and Edelman (19). Each of the radiolabeled and column-purified proteins was reduced with  $\beta$ -mercaptoethanol and dialyzed

against 1 M propionic acid for 72 hr. Chain separation was accomplished on a Sephadex G-200 column made 1 M in propionic acid and 0.01 M mercaptoethanol. Hybrid molecules were formed using H chains from a  $\gamma$ G3 myeloma protein labeled with <sup>125</sup>I (G. J.) and the light chains from a  $\gamma$ G1 myeloma protein labeled with <sup>125</sup>I (E. T.). Recombination was usually done on a milligram per milligram basis of H chain per L chain. Similar hybrids were made with a  $\gamma$ G1 heavy chain and the light chains derived from a  $\gamma$ G3 protein. After prolonged dialysis against acid and then neutral buffers containing 0.001 M mercaptoethanol, dialysis was continued for 72 hr against phosphate-buffered saline pH 7.2 with no mercaptoethanol. The recombined molecules were then concentrated slowly by dialysis against 3% polyethylene glycol (PEG) in 1 M propionic acid. They were then purified on Sephadex G-200 columns made 1 M in propionic acid. The  $\gamma$ G peak usually accounted for over 80% of the radioactivity. This was pooled and concentrated by dialysis against 3% PEG and then utilized in the acrylamide-gel electrophoresis experiments. In most experiments recombined molecules contained from 1.17 to 1.25 moles of H chain per L chain.

**Analytical ultracentrifugation.** A Spinco model E analytical ultracentrifuge equipped with schlieren optics was used for all studies. Unless otherwise noted, studies were performed in 12 mm double-sector cells at 20°C and 52,640 rpm. Plate measurements were made according to procedures described by Trautman (20). In correcting the sedimentation coefficients to the  $s_{20,w}$  value the partial specific volume was assumed to be 0.73. Studies were done on undiluted sera, sera diluted 1:1 with PBS and with purified myeloma proteins and their fragments in PBS. Sedimentation constants at infinite dilution were derived from plots of  $s_{20,w}$  vs. concentration.

Molecular weights were determined by the method of Kegeles and Sia (21) using the field relaxation procedure. Fluorochemical Fc 43 (Beckman/Spinco Division, Palo Alto, Calif.) was used in a capillary synthetic boundary cell. Molecular weights were determined in pH 7.2 PBS as well as 0.1 M acetate buffer pH 4.0. In each study the protein concentration was 8.0  $\pm$  0.2 mg/cc. Computation of the molecular weight at the cell bottom was made from the following equation:

$$M_w = \frac{RT \left. \frac{dc}{dr} \right|_b}{(1 - \bar{v}_p)r_b(A_1 + A_2)\omega^2}$$

where  $M_w$  is the weight average molecular weight at the cell bottom,  $\left. \frac{dc}{dr} \right|_b$  is the extrapolated deflection above the estimated baseline of the schlieren diagram at the lower meniscus,  $r_b$  is the absolute position in the cell and  $A_1$  and  $A_2$  the areas below the free peak and the peak near the cell bottom. These areas were measured by the trapezoid method.

**Acrylamide-gel electrophoresis.** Acid-acrylamide-gel electrophoresis was performed in an E-C Apparatus Corp. vertical gel electrophoresis cell. The basic procedure was the Canalco modification of the system of Williams and Reisfield (22). The acid gel buffer was KOH-citric acid pH 2.9 while the tank buffer was glycine-citrate pH 4.0. For urea-acid gels both the gel and the tank buffers contained 8 M urea. 7.5% acrylamide gels were constructed from a stock solution (5% bisacrylamide, 95% acrylamide) which contained ascorbic acid, ferrous sulfate, and 0.001 M iodo-

acetamide. Gels were polymerized by the addition of 5-10  $\mu$ l of 30% cold hydrogen peroxide (23). Acid gels were run for 4 hr at 300 v while acid-urea gels were run for 8-12 hr. The slabs were stained with amido black 0.25% in methanol:water:glacial acetic acid 5:5:1. Destaining was accomplished in an electrophoretic destainer (E-C Apparatus Corp.) in a similar solution without amido black.

**Tests for anti- $\gamma$ -globulin activity.** The Rh agglutination test was done according to Waller and Vaughan (24). Anti-CD serum Ripley (kindly supplied by Dr. Marion Waller, Richmond, Va.) was used for these studies. The latex fixation test was obtained from Hyland Laboratories, Los Angeles, Calif. (RA-Test). Both the Rh agglutination test and the latex fixation test were performed at 4° and 22°C.

In separate experiments Fab and (Fab')<sub>2</sub> from each protein were mixed with normal  $\gamma$ -globulin as well as isolated myeloma proteins from each  $\gamma$ G heavy-chain subclass and examined in the analytical ultracentrifuge for interactions. The area under the curve of Fab or (Fab')<sub>2</sub> was compared to the same material in a wedge cell without additional  $\gamma$ -globulin in order to assess for complex formation.

**Test for cryoglobulins.** Attempts were made to demonstrate significant amounts of cryoglobulins in each of the sera. Despite prolonged standing in the cold, dilution, and pH changes from 5 to 8.5 no cryoproteins were evident. In a final attempt each serum was spun at 35,000 rpm in an SW 39 Rotor of a Spinco model L ultracentrifuge for 18 hr at 4°C. No cryoprecipitate was evident in the bottom of any tube.

**Viscosity studies.** All solutions introduced into the viscometers were filtered repeatedly through a Millipore filter pad. The density of the solutions was measured with a 5 ml capillary vent pycnometer. Capillary viscosity measurements were done in a constant temperature bath utilizing an Ostwald Viscometer. Phosphate-buffered saline was the reference solvent in all studies. Measurements were performed at 4°, 20°, 22°, and 37°C and each determination was the average of three separate studies.

The specific viscosity was derived from the expression:  $\eta_{sp} = \eta_{rel} - 1$ . The intrinsic viscosity was determined by a plot of  $\eta_{sp}/c$  vs. C where C represents the protein concentration in grams per cubic centimeter. Studies were performed over a protein concentration range of 0.001-0.067 g/cc.

A modified version of the G. D. M. viscometer was used to determine the viscosity at low shear rates (25). The principle of this viscometer, which consists of coaxial cylinders separated by a small annulus in which the sample is located has been described (26, 27). Utilizing a water bath connected to a channel inside the rotor and a thermistor connected to a thermo regulator at the exit from the channel, sample temperature could be maintained at  $\pm 0.1^\circ\text{C}$ . Studies were done at 4°, 20°, and 37°C. Measurements were performed at five shear rates (52, 5.2, 0.52, 0.104, and 0.052  $\text{sec}^{-1}$ ).

## RESULTS

**Analytical ultracentrifugation of serum and isolated myeloma proteins.** When  $\gamma$ G3 myeloma sera were diluted 1:1 with phosphate-buffered saline and studied in the analytical ultracentrifuge they did not appear conspicuously different from the sera from patients with multiple myeloma involving other subgroups (Fig. 1, top). However, when studied undiluted, marked differences were obvious (Fig. 1, bottom). Seven  $\gamma$ G3 myeloma sera were studied and each showed a shelf of complexes arising from the 7S peak and moving to the bottom of the cell. That this was not an artifact of examining undiluted sera was shown by the fact that in the undiluted sera of several patients with multiple myeloma involving other subgroups no such complexes were noted (Fig. 1, E. T.). In addition, under these conditions marked convective disturbances were noted only




FIGURE 1 Analytical ultracentrifuge analysis of three sera from patients with  $\gamma$ G3 multiple myeloma (G. J., P. W., and A. L.) and serum E. T.  $\gamma$ G1. Each photograph was taken 64 min after reaching speed of 52,640 rpm. The shelf of aggregating material is only seen in the undiluted sera of the  $\gamma$ G3 cases.

in the undiluted sera of the patients with  $\gamma$ G3 myeloma. The amount of complexes seen was correlated with the amount of myeloma protein in the patient's serum and  $\gamma$ G3 myeloma sera with less than 15 mg/ml of myeloma protein did not readily demonstrate complex formation on analytical ultracentrifugation.

In order to determine the sedimentation constant for these  $\gamma$ G3 proteins three of the isolated myeloma proteins were sedimented at various concentrations and on S vs. C plot was constructed (Fig. 2). Protein G. J. had an extrapolating value of 6.72 at infinite dilution. Protein A. L. had a value of 6.81 and protein P. W. was 6.78. At higher concentrations each isolated myeloma protein was seen to aggregate and the S value reported in based on the analysis of the main peak alone (Fig. 2). The radius of gyration of each profile was calculated with the aid of a computer program. In each case, with increasing protein concentration there was accelerated skewing of the schlieren profile towards the base of the cell. Thus, the increase in the radius of gyration suggests that the heavier components seen at higher concentrations are the product of the higher concentration and not secondary to better visualization of these components. Each of the isolated proteins was dialyzed against acetate and phosphate buffers ranging from pH 4.0 to 8.0. Each solution was then adjusted to 20 mg/ml and studied in the analytical ultracentrifuge. Complex

formation was maximal at pH 6.8. This is illustrated in a later section for  $\gamma$ G3 fragments (Fig. 6).

Low-speed determinations of molecular weights were performed in an attempt to rule out an effect of the extreme pressures in the cell during high-speed centrifugation contributing to the complexing phenomenon. For these studies special care was taken that the isolated proteins did not contain stable aggregates which might have arisen from the isolation procedure (see Methods). As illustrated in Fig. 3 the weight average molecular weight at the bottom of the cell was considerably higher in the isolated protein from patient G. J. with  $\gamma$ G3 myeloma than the protein isolated from E. T. who had  $\gamma$ G1 multiple myeloma. In each case the molecular weight determined in pH 4.0 acetate buffer compared favorably with the accepted values for  $\gamma$ G globulin in the literature.

*Acrylamide-gel electrophoresis.* Acrylamide-gel electrophoresis proved to be a sensitive technique for demonstrating aggregation phenomena, apparently because of the concentrating effect that occurs before entrance into the gel. The concentration-dependent aggregation was seen even at low pH. Thus while at pH 4.0 no aggregation could be demonstrated in the analytical ultracentrifuge, aggregation was clearly seen in acrylamide gels at pH's as low as 2.9. Fig. 4 represents an acid-acrylamide-gel electrophoresis of seven sera from




FIGURE 2 The concentration dependence of sedimentation of G. J. isolated  $\gamma$ G3 protein. The extrapolated value of  $s_{20,w}$  (in Svedberg units) at infinite dilution was 6.72. The insert shows aggregation at higher concentration.




FIGURE 3 Apparent molecular weights of isolated myeloma proteins by the Field Relaxation Technique. In each experiment the sample was accelerated to 20,000 rpm for 4 min and decelerated at approximately  $1000 \text{ rpm min}^{-1}$  to 4000 rpm. Photographs were taken after 20 min at the lower speed.

patients with  $\gamma$ G3 myeloma and one serum from a patient with  $\gamma$ G1 myeloma, all with large protein peaks. It was clear that the  $\gamma$ -globulin in serum from the patients with  $\gamma$ G3 myeloma did not significantly enter the gel while the  $\gamma$ G from serum E. T. ( $\gamma$ G1 myeloma) did enter the gel. Fig. 5 represents the same experiment except the gel contained 8 M urea. Here the  $\gamma$ -globulin from all patients entered the gel and appeared virtually identical. Sera from four other patients with  $\gamma$ G3 myeloma were studied and behaved in an identical way with those shown in Figs. 4 and 5. Gm (b) or Gm (g) and kappa or lambda proteins behaved similarly. 25 sera from patients with multiple myeloma of the  $\gamma$ G1, 2, and 4 subgroups were similarly studied and in each instance the  $\gamma$ G paraprotein peak entered both acid-acrylamide-gel systems in a normal fashion.

In order to determine whether  $\gamma$ G3 proteins in normal serum also displayed a concentration-dependent aggregation, normal serum was run on acid-acrylamide-gel electrophoresis. Without staining, the origin was cut out as well as the position where the  $\gamma$ G normally mi-

grates. Each portion of the gel was then forced through an 18-gauge needle and agar plate analysis was performed with  $\gamma$ G subgroup-specific antisera. Only  $\gamma$ G3 proteins were detected at the origin and no  $\gamma$ G3 was found in the main  $\gamma$ G band.

*Molecular localization of aggregation site.* In order to determine which portion of the  $\gamma$ G3 molecule was responsible for this unusual property of aggregation, enzymatic fragments as well as H and L chains were studied. Fab and (Fab')<sub>2</sub> from these proteins failed to enter the acid-acrylamide gels while similar fragments from control  $\gamma$ G1 proteins entered normally. Studies in acid-urea gels did not distinguish the two groups, again indicating the noncovalent nature of the forces of aggregation. Fc fragments from  $\gamma$ G3 proteins entered the acid gels in a normal position. Analytical ultracentrifugation of A. L. (Fab')<sub>2</sub> is depicted in Fig. 6. Like several  $\gamma$ G3 whole proteins, aggregation was maximal around pH 6.8. Fc fragments from  $\gamma$ G3 proteins did not aggregate in the analytical ultracentrifuge.

Reduction and alkylation of the isolated proteins and controls revealed that on acid-urea gels both H and L chains entered the gels. On acid gels after reduction and




FIGURE 4 Acid-acrylamide-gel electrophoresis of seven  $\gamma$ G3 myeloma sera (1-7) and a  $\gamma$ G1 myeloma serum (slot 8). None of the  $\gamma$ G3 proteins are visualized because of failure to enter the gel.



alkylation the L chains of all  $\gamma$ G3 proteins entered normally but the H chains aggregated at the origin. All control H chain preparations from proteins of  $\gamma$ G1, 2, and 4 subgroups entered the acid gels under these conditions.

While these results strongly implicated the Fd portion of the H chain as the aggregating unit, studies were done with recombined and reoxidized molecules for confirmation. Hybrid molecules were formed between G. J. H. chains labeled with  $^{125}\text{I}$  and E. T. light-chains labeled with  $^{125}\text{I}$ . Also, G. J. light-chains labeled with  $^{125}\text{I}$  were recombined with E. T. H chain labeled with  $^{125}\text{I}$ . As seen in Fig. 7 (left) when the H chain in the hybrid molecule arose from the  $\gamma$ G3 protein, the entire recombined molecule would not enter the acid gel. However, when the H chain arose from a  $\gamma$ G1 protein, the entire hybrid entered the gel normally. On acid-urea gels the two hybrids could not be distinguished from each other as both entered normally. Finally on re-reduction only the H chains from the  $\gamma$ G3 protein accumulated at the origin. Thus the light chains seem to play no role in the aggregation phenomena whereas the H chain and specifically the Fd portion seems to be implicated.

*Tests for anti- $\gamma$ -globulin activity.* Whole sera as well as the isolated paraproteins were tested for rheumatoid factor activity by the Rh agglutination test and the latex fixation test. Serum A. L. had a titer of 1000 in the Rh agglutination test but the isolated  $\gamma$ G3 protein was devoid of rheumatoid factor activity. Pevikon block electrophoresis separated the rheumatoid factor in serum A. L. from the myeloma protein. All other sera in this study as well as the isolated proteins from most were




FIGURE 5 Same experiment as Fig. 4 except the gel was made 8 M in urea. The myeloma components are now apparent for the  $\gamma$ G3 proteins.




FIGURE 6 Effect of pH on aggregation of the pepsin  $(\text{Fab}')_2$  fragment of the  $\gamma$ G3 protein A. L.

devoid of rheumatoid factor activity both at 4°C and 20°C. In extensive experiments with the analytical ultracentrifuge neither pepsin  $(\text{Fab}')_2$  nor papain Fab fragments prepared from three of the  $\gamma$ G3 proteins could be demonstrated to combine with either autologous Fc or Fc prepared from pooled  $\gamma$ -globulin.

*Viscosity studies.* During the early stages of these studies two of the patients with  $\gamma$ G3 myeloma proteins developed the hyperviscosity syndrome. This initiated viscosity studies of their proteins, other  $\gamma$ G3 myeloma proteins, and myeloma proteins of other subgroups. The initial serum relative viscosity in the two cases was 14.6 and 10.0 at 22°C. The concentration-dependent aggregation of these proteins was presented earlier in the analytical ultracentrifuge studies (Fig. 1, G. J. and A. L. as well as P. W. without hyperviscosity syndrome).

The pronounced effect of temperature on the relative viscosity of  $\gamma$ G3 proteins is shown in Table I. All sera with  $\gamma$ G3 proteins demonstrated an increased relative viscosity at lower temperatures. This is evident whether the hyperviscosity syndrome is present or not. No patient with  $\gamma$ G1 myeloma or with macroglobulinemia demonstrated a significant increase in relative viscosity at lower temperatures. In  $\gamma$ G3 myeloma hyperviscosity becomes significant at much lower concentrations of myeloma protein. Thus with only 6.2 g/100 ml of  $\gamma$ G3 protein patient G. J. had a relative viscosity at 22°C of 14.6. Patient M. K. with  $\gamma$ G1 myeloma with 19.0 g of  $\gamma$ G1 protein had a serum relative viscosity of only 6.0 at 22°C. This distinction between the effect of protein concentration on relative viscosity in  $\gamma$ G3 vs.  $\gamma$ G1 myeloma is shown again in Fig. 8. The markedly different slopes indicate that per unit of concentration a  $\gamma$ G3 protein results in a far more viscous serum than a  $\gamma$ G1 protein.

Each serum studied in Table I was diluted with equal volumes of PBS and the relative viscosity determined. Table II depicts some of the results. The relative viscosity of NHS and the sera of patients with multiple




FIGURE 7 Acid-acrylamide-gel electrophoresis of radioiodinated recombined and reoxidized molecules. 7A shows that when the  $\gamma$ -chain of the hybrid molecule comes from the  $\gamma$ G3 protein, the molecule does not enter the gel. 7B indicates that when the  $\gamma$ -chain of the hybrid molecule comes from the  $\gamma$ G1 protein, the molecule enters the gel normally. 7C and 7D show that on re-reduction only the  $\gamma$ -chain from the  $\gamma$ G1 protein enters the gels. For further details see text.

myeloma of the  $\gamma$ G1, two or four subgroups without hyperviscosity syndrome, was reduced less than 10% by dilution. The sera of patients with Waldenström's macroglobulinemia and hyperviscosity syndrome was reduced less than 20% and those from patients with  $\gamma$ G1 myeloma and hyperviscosity syndrome was reduced approximately 50%. In contrast, the sera from patients with  $\gamma$ G3 myeloma and hyperviscosity syndrome was reduced 75% or greater.

Intrinsic viscosity was determined on some of the isolated proteins and was found to be 4.1 cc/g for G. J. and 4.6 cc/g for A. L. Thus at infinite dilution these proteins have intrinsic viscosity values which do not differ significantly from pooled gamma globulin.

Plasma viscosity determinations at low shear rates (Table III) showed no shear dependence. Thus the viscosity in centipoises did not change over a 1000-fold change in shear rate for any plasma. The clear temperature dependence is again noted for the  $\gamma$ G3 proteins but not for the  $\gamma$ G1 control. These studies were performed on plasma obtained via plasmapheresis and were thus diluted. This accounts for the lower relative viscosity values.

## DISCUSSION

The initial observations of concentration-dependent aggregation of  $\gamma$ G3 proteins were made on analytical ultracentrifugation of whole serum. The large shelf of

complexes arising from the 7S peak and in a continuum to the cell bottom suggested an unstable associating-dissociating system. The convective disturbances seen in Fig. 1 probably represent density inversions due to slow interconversion of monomers into polymers and vice versa. The sawtooth spikes are indicative of convective disturbances which lead to extremely sharp concentration gradients (28). The effect of pressure on aggregation phenomena are well described (29). At maximum angular rotation, pressures of over 250 atmospheres occur in ultracentrifuge cells (30) so the experiments at low speeds were designed to minimize the effect of pressure on the aggregation phenomena. The particular method of purifying the isolated myeloma proteins was designed to reduce all factors which might lead to aggregation thereby allowing the effect of concentration at the cell bottom to be the major determinant in the molecular weight measurement. The value of 305,000 for G. J represents a weight average molecular weight not of the original sample, but instead, is the apparent molecular weight of the material present at the lower meniscus at a given instant (21). Thus the demonstration of an apparent molecular weight of over 300,000 provides strong evidence that the aggregation phenomena is concentration and not pressure dependent.

Aggregation of  $\gamma$ G3 proteins was also documented by their failure to penetrate acrylamide gels in which proteins with molecular weights less than a million will



TABLE I  
Comparative Results of Protein and Viscosity Studies on the Serum of Patients with  $\gamma$ G3,  $\gamma$ G1,  
and Macroglobulin Peaks with and without the Hyperviscosity Syndrome

Patient	Type	Hyperviscosity syndrome	Total protein	Peak	Serum relative viscosity		
					4°C	22°C	37°C
Normal	—	—	7.1	—	1.5	1.4	1.4
G. J.	$\gamma$ G3 lambda	+	11.2	6.2	37.0	14.6	5.8
A. L.	$\gamma$ G3 lambda	+	11.0	5.8	16.4	10.0	4.0
W. P.	$\gamma$ G3 kappa	+	10.5	4.2	10.2	6.4	3.1
P. W.	$\gamma$ G3 kappa	—	10.0	3.6	7.7	4.2	2.2
C. V.	$\gamma$ G3 lambda	—	9.6	2.8	3.3	2.0	1.4
L. A.	$\gamma$ G3 kappa	—	7.4	2.0	2.0	1.7	1.4
M. K.	$\gamma$ G1 kappa	+	25.0	18.0	5.2	5.1	5.0
D. P.	$\gamma$ G1 kappa	+	24.1	17.2	4.9	4.9	4.8
C. F.	$\gamma$ G1 kappa	+	22.2	15.0	4.7	4.5	4.4
N. B.	$\gamma$ G1 kappa	—	19.0	12.0	3.6	3.5	3.5
R. B.	$\gamma$ G1 kappa	—	14.4	9.5	2.8	2.8	2.7
A. G.	$\gamma$ G1 lambda	—	12.6	7.5	2.2	2.1	2.1
E. T.	$\gamma$ G1 kappa	—	12.0	6.2	1.7	1.6	1.6
S. N.	$\gamma$ G1 kappa	—	11.9	6.2	1.6	1.5	1.5
A. G.	$\gamma$ M lambda	+	12.6	6.2	6.8	6.5	6.4
B. D.	$\gamma$ M kappa	+	12.0	7.0	8.2	8.2	8.1
S. W.	$\gamma$ M kappa	+	11.4	5.8	5.8	5.6	5.5

readily enter. The so-called "stacking" effect of the gel at the point of entry apparently produces a high concentration of the  $\gamma$ G3 proteins to effect their aggregation.

Studies on enzymatic fragments and reduction products localized the aggregating portion of the molecule to Fab and H chain. This strongly suggests that the  $\gamma$ G3 Fd fragment is responsible for this phenomenon. There




FIGURE 8 A graphic representation of some of the data in Table I.  $\gamma$ G1 myelomas fall on a line with a much lower slope than  $\gamma$ G3 proteins. The points represent the individual patients.

TABLE II  
Dilution Test for Concentration-Dependent Aggregation

Patient	Relative viscosity at 20°C of whole serum	Relative viscosity at 20°C of serum diluted with PBS	Reduction
NHS*	1.51	1.40	7%
Multiple myeloma without hyperviscosity syndrome†	2.06	1.90	8%
Waldenström's macroglobulinemia with hyperviscosity syndrome‡	6.60	5.40	18%
M. K.	6.01	3.12	52%
C. F.	5.07	2.61	51%
G. J.	13.00	2.81	88%
A. L.	8.04	2.02	75%
W. P.	7.52	1.88	75%

\* Average of five normals.

† Sera from 30 patients,  $\gamma$ G1,  $\gamma$ G2, and  $\gamma$ G4 multiple myeloma.

‡ Sera from three patients.

is a considerable body of data which suggests that subclass-specific antigens reside on both the Fc and the Fd portion of the H chain (16, 31). This is especially true in the  $\gamma$ G3 subgroup where the Fd antigens are particularly potent. It is likely that these same areas of the

molecule render it more susceptible to aggregating potential. It may be that a string of strongly hydrophobic residues mutually attract and lead to aggregation. Frangione and Milstein (32) report that the  $\gamma$ G3 subgroup is unique among the  $\gamma$ G globulins having five inter-heavy-chain disulphide bonds. One and perhaps two of these is in the Fd region. The role the additional half-cysteines play in this aggregation phenomena is not clear at the present time.

Recent studies on the catabolism of human myeloma proteins (4, 5) show that those of the  $\gamma$ G3 subclass are catabolized much faster than any other subgroup. Spiegelberg and Fishkin (4) also report that this difference is not accounted for by more recent studies on the Fc fragment. It is reasonable to assume on the basis of the present study that the different catabolic rate may be related to in vivo aggregation and subsequent destruction, which would be dependent on the Fd fragment.

The hyperviscosity syndrome is a constellation of signs and symptoms including tinnitus, mucous membrane bleeding, stupor, stroke, convulsions, blurred vision, dilated retinal veins, gastrointestinal hemorrhage, and congestive heart failure (33). Clinically the most common disease associated with this syndrome is Waldenström's macroglobulinemia (9). In multiple myeloma, however, the hyperviscosity syndrome is rare.

Bjørneboe and Jensen (13) described a patient with hyperviscosity syndrome and  $\gamma$ G multiple myeloma in which measurement of colloid osmotic pressure was normal and the plasma volume was only slightly increased. They postulated aggregation of the parapro-

TABLE III  
Plasma Viscosity at Low Shear Rates

Temperature	Shear rate	Viscosity in centipoises			Relative viscosity			
		G. J.	A. L.	E. T.	Water	G. J.	A. L.	E. T.
°C	sec <sup>-1</sup>							
37	52	2.47	2.98	1.82	0.6947	3.56	4.29	2.62
	5.2	2.47	2.98	1.82				
	0.52	2.47	2.98	1.82				
	0.104	2.47	2.98	1.82				
	0.052	2.47	2.98	1.92				
20	52	5.13	5.04	2.61	1.0050	5.10	5.01	2.60
	5.2	5.13	5.04	2.61				
	0.52	5.13	5.04	2.61				
	0.104	5.13	5.04	2.61				
	0.052	5.13	5.04	2.61				
4	52	13.4	11.2	3.92	1.5674	8.55	7.15	2.51
	5.2	13.4	11.2	3.92				
	0.52	13.4	11.2	3.92				
	0.104	13.4	11.2	3.92				
	0.052	13.4	11.2	3.92				

tein as a mechanism to explain these findings but were unable to demonstrate aggregation despite analytical ultracentrifugation studies. However, since they diluted the serum about 10-fold before centrifugation, concentration-dependent aggregation would not have been detected. Kopp, Beirne, and Burns (14) described a similar patient with  $\gamma$ G myeloma and hyperviscosity syndrome with a serum relative viscosity of 6.0 and again analytical ultracentrifugation did not demonstrate aggregate formation. However, the centrifuge photograph shown suggests that the sera had been diluted 5–10 times and concentration-dependent aggregates as described herein would not have been observed. They measured relative viscosity at 37°C and found it to be 4.8 while at 24°C it was 7.0. This increase in relative viscosity from 37°C to 24°C suggests that they were probably confronted with the same phenomenon reported here, i.e., concentration- and temperature-dependent aggregation. MacKenzie and Fudenberg (15) reported on two patients with hyperviscosity syndrome secondary to  $\gamma$ G myeloma in which no aggregates could be documented by analytical ultracentrifugation. The intrinsic viscosity of the isolated proteins was determined and it was normal in one patient while in the other it was reported to be twice normal. They concluded that hyperviscosity syndrome in  $\gamma$ G myeloma was due to the primary structure of the individual proteins, not their potential for aggregation. In view of the data presented in this report it is likely that in some of the above patients concentration- and temperature-dependent aggregation of  $\gamma$ G3 proteins lead to the hyperviscosity syndrome.

Two potentially useful simple tests for unstable aggregating  $\gamma$ G myelomas producing hyperviscosity syndrome are the serum relative viscosity determination at two different temperatures and the serum relative viscosity on whole serum and serum diluted 1:1. The temperature-viscosity index (TVI) was introduced by Waldenström (34).  $TVI = (100 \times \text{rel. visc. } 13^\circ\text{C}) / (\text{rel. visc. } 37^\circ)$ . Values greater than 120 were considered suggestive of macroglobulinemia. Somer (35) in his extensive study of viscosity in myeloma and macroglobulinemia reexamined this test. He found that of 16 cases of  $\gamma$ G myeloma only three had TVI 120 and in only two was the serum relative viscosity at 20°C 5.0. Only 5 of 15 sera from patients with Waldenström's macroglobulinemia had TVI values above 120. He concluded that the TVI was of somewhat limited usefulness clinically except in cases of cryoglobulinemia. It should be emphasized that in the present study cryoglobulins were not present in any of these sera. The temperature viscosity index can only be extrapolated from the present study as measurements were not done at 13°C. However, it is clear that none of the three patients with Waldenström's macroglobulinemia who served as con-

trols had a TVI > 120 (Table I). In addition, patients with multiple myeloma without hyperviscosity syndrome had a TVI within the normal range. However, the marked effect of temperature on viscosity is evident in the two patients with hyperviscosity syndrome with  $\gamma$ G3 multiple myeloma, for A. L., 276 and for G. J., 460. These values are higher than any value obtained by Somer (35) in his study of sera from 15 patients with macroglobulinemia, 7 with  $\gamma$ A myeloma and 16 with  $\gamma$ G myeloma. These studies would therefore suggest that viscosity determination at two different temperatures can be useful to distinguish this form of hyperviscosity syndrome from that caused by Waldenström's macroglobulinemia when analytical ultracentrifugation is not readily available.

Another clinically useful test utilizing the Ostwald viscometer is the dilution test. Steel (36) introduced the "interpolated 7% viscosity" based on a fixed protein concentration. He noted that by diluting the serum with elevated viscosity to a protein content of 7 g/100 ml the relative viscosity of myeloma sera is reduced to the normal range but in cases of macroglobulinemia the "7% viscosity" remains abnormal. The value of the test as reported here, however, is that protein concentration need not be measured as shown in Table II. A simple 1:1 dilution of the serum reduces the serum relative viscosity of these concentration-dependent aggregating myelomas to close to the normal range, while the relative viscosity of the sera of patients with Waldenström's macroglobulinemia is much less affected.

Capillary tube viscometers in general measure viscosity at relatively high shear rates. Measurements of flow velocity in post capillary venules suggest that the shear rate in this portion of the microcirculation is low (37). It was therefore desirable to study the effect of low shear rates on viscosity in this unique group of patients. The results (Table III) indicate that there is no shear dependence and suggest Newtonian behavior of the plasma. However, it has long been known that whole blood is non-Newtonian (38, 39) and that there is a tendency for the non-Newtonian behavior to increase as the viscosity level rises (27). Thus it appears that for maximum information whole blood instead of the diluted citrated plasma will have to be studied at low shear rates and such studies are now underway.

The  $\gamma$ G heavy-chain subgroups of myeloma proteins associated with the hyperviscosity syndrome shows a disproportionate number belonging to the  $\gamma$ G3 subgroup. Three in the present report are  $\gamma$ G3 proteins; however, the two patients of MacKenzie and Fudenberg (15) were of the  $\gamma$ G1 subgroup but a third patient provided to us by these authors was a  $\gamma$ G3 protein. Studies of the sera of patients with hyperviscosity syn-

drome with  $\gamma$ G1 myeloma indicate that: (a) the myeloma protein concentration in each case was 15 g or over (see Table I); (b) no aggregates could be documented by analytical ultracentrifugation; (c) all entered the acid-acrylamide gel; and (d) none demonstrated a significant increase in relative viscosity at lower temperatures. Although these studies are somewhat preliminary, it would appear that the phenomenon of hyperviscosity syndrome in  $\gamma$ G1 myeloma seems more fundamentally related to extreme levels of myeloma protein in the circulation.

The exact proportion of hyperviscosity syndrome cases that are due to  $\gamma$ G3 proteins remains uncertain because of the limited number of cases available. A total of nine cases of  $\gamma$ G myeloma with hyperviscosity syndrome have passed through the authors' laboratory including those described above; five of the nine proved to involve  $\gamma$ G3 proteins. Since only 7–10% of myeloma proteins belong to the  $\gamma$ G3 subgroup, this high proportion appears impressive. It seems probable that the concentration- and temperature-dependent aggregation described for these proteins played a significant role in the usual incidence. The evidence obtained in the present study indicates that the hyperviscosity syndrome develops when  $\gamma$ G3 proteins reach a concentration of approximately 4 g/100 ml. Other  $\gamma$ G globulins require considerably higher concentrations before hyperviscosity develops.

The finding of multiple myeloma with a  $\gamma$ G3 protein should alert the clinician to the possibility of hyperviscosity playing a role in the symptomatology at some point in the course of the disease particularly if the protein peak is above 4 g/100 ml. In addition, whenever  $\gamma$ G3 myeloma is diagnosed, reduction of the level of myeloma protein should be a definitive goal of therapy.

The preponderance of the  $\gamma$ G3 subgroup in this disorder recalls the group of  $\gamma$ G cryoglobulins with anti- $\gamma$ -globulin activity recently reported by Grey, Kohler, Terry, and Franklin (7). Of the five proteins in their series with anti- $\gamma$ G activity, four were  $\gamma$ G3 and one was  $\gamma$ G1. However, the proteins recorded in this present study were not cryoglobulins and lacked anti- $\gamma$ -globulin activity by several criteria. Thus the mechanism of aggregation of the two groups of proteins appears to be fundamentally different.

It appears, therefore, that there are at least three mechanisms whereby the hyperviscosity syndrome may develop in  $\gamma$ G myeloma. (a) The proteins may form relatively stable aggregates of 13–15S as in the cases of Smith et al. (12, 40); (b) extreme levels of myeloma proteins as our cases of  $\gamma$ G1 myeloma with hyperviscosity syndrome; and (c) unstable complex formation of  $\gamma$ G3 proteins.

## ACKNOWLEDGMENTS

We are pleased to acknowledge the able technical assistance of Miss Vicki Eisch whose careful work has made this endeavor a particular pleasure. We thank Doctors M. Grieco and T. Tamlyn of St. Luke's Hospital and Doctors G. Benninger and S. Kreps of Lenox Hill Hospital for their help in the diagnosis and management of some of these patients. We are also grateful to Doctors G. Penn, L. Cawley, M. Mannik, and W. Terry for providing certain sera. Our thanks also to Doctors S. Usami and S. Chien of the Department of Physiology, College of Physicians & Surgeons, Columbia University, who performed the viscosity studies at low shear rates.

This investigation was supported by U. S. Public Health Service Grant AM 09792. Dr. Capra is recipient of U. S. Public Health Service Special Fellowship 1-F3-CA-35, 190-01.

## REFERENCES

1. Grey, H. M., and H. G. Kunkel. 1964. H chain subgroups of myeloma proteins and normal 7S  $\gamma$ -globulin. *J. Exp. Med.* **120**: 253.
2. Terry, W. D., and J. L. Fahey. 1964. Subclasses of human  $\gamma$ 2 globulin based on differences in the heavy polypeptide chains. *Science (Washington)*. **146**: 400.
3. Yount, W. J., H. G. Kunkel, and S. D. Litwin. 1967. Studies of the Vi ( $\gamma$ 2c) subgroup of  $\gamma$  globulin. *J. Exp. Med.* **125**: 177.
4. Spiegelberg, H. L., B. G. Fishkin, and H. M. Grey. 1969. Catabolism of human  $\gamma$ G-immunoglobulins of different heavy chain subclasses. *J. Clin. Invest.* **47**: 2323.
5. Morell, A., W. D. Terry, and T. Waldmann. 1969. Relation between metabolic properties and serum concentration of IgG-subclasses in man. *Clin. Res.* **17**: 356.
6. Yount, W. J., R. Hong, M. Seligmann, and H. G. Kunkel. 1969. Structural gene defects and preponderance of  $\gamma$ G3 subgroup in patients with hypogammaglobulinemia. *J. Clin. Invest.* **48**: 92 a. (Abstr.)
7. Grey, H. M., P. F. Kohler, W. D. Terry, and E. C. Franklin. 1968. Human monoclonal  $\gamma$ G-cryoglobulins with anti- $\gamma$ -globulin activity. *J. Clin. Invest.* **47**: 1875.
8. Ritzmann, S. E., R. H. Thurm, W. E. Truax, and W. C. Levin. 1960. The syndrome of macroglobulinemia. *Arch. Intern. Med.* **105**: 959.
9. Fahey, J. L., W. F. Barth, and A. Solomon. 1965. Serum hyperviscosity syndrome. *J. Amer. Med. Ass.* **192**: 464.
10. Deutsch, H. F., and J. I. Morton. 1958. Human serum macroglobulins and dissociation units. I. Physicochemical properties. *J. Biol. Chem.* **231**: 1107.
11. Vaerman, J. P., L. B. Johnson, W. Mandy, and H. H. Fudenberg. 1965. Multiple myeloma with two paraprotein peaks. *J. Lab. Clin. Med.* **65**: 18.
12. Smith, E., S. Kochwa, and L. R. Wasserman. 1965. Aggregation of IgG globulin in vivo. *Amer. J. Med.* **39**: 35.
13. Bjørneboe, M., and K. B. Jensen. 1966. A case of myelomatosis with normal colloid osmotic pressure in spite of extremely high serum protein concentration. *Acta Med. Scand. Suppl.* **445**: 212.
14. Kopp, W. L., G. J. Beirne, and R. O. Burns. 1967. Hyperviscosity in multiple myeloma. *Amer. J. Med.* **43**: 141.
15. MacKenzie, M. R., and H. H. Fudenberg. 1967. Role of intrinsic viscosity in myeloma hyperviscosity syndrome. *Clin. Res.* **15**: 388.

16. Natvig, J. B., and H. G. Kunkel. 1968. Genetic markers of human immunoglobulins. *Ser. Haematol.* 1: 66.
17. Porter, R. R. 1959. The hydrolysis of rabbit  $\gamma$ -globulin and antibodies with crystalline papain. *Biochem. J.* 73: 119.
18. McConahey, P. J., and F. J. Dixon. 1966. A method of trace iodination of proteins for immunologic studies. *Intern. Arch. Allergy Appl. Immunol.* 29: 185.
19. Olins, D. E., and G. M. Edelman. 1964. Reconstitution of 7S molecules from light and heavy polypeptide chains of antibodies and  $\gamma$  globulins. *J. Exp. Med.* 119: 789.
20. Trautman, R. 1956. Operating and comparing procedures facilitating schlieren pattern analysis in analytical ultracentrifugation. *J. Phys. Chem.* 60: 1211.
21. Kegeles, G., and C. L. Sia. 1963. An ultracentrifugal field relaxation method for molecular weights. *Biochemistry.* 2: 906.
22. Williams, D. E., and R. A. Reisfeld. 1964. Disc electrophoresis in polyacrylamide gels. *Ann. N. Y. Acad. Sci.* 121: 373.
23. Jordan, E. M., and S. Raymond. 1969. Gel electrophoresis: a new catalyst for acid systems. *Anal. Biochem.* 27: 205.
24. Waller, M. V., and J. H. Vaughan. 1956. Use of anti Rh sera for demonstrating agglutination activating factor in rheumatoid arthritis. *Proc. Soc. Exp. Biol. Med.* 92: 198.
25. Chien, S., S. Usami, H. M. Taylor, J. L. Lundberg, and M. I. Gregersen. 1966. Effects of hematocrit and plasma proteins on human blood rheology at low shear rates. *J. Appl. Physiol.* 21: 81.
26. Merrill, E. W., E. R. Gilliland, G. Cokelet, H. Shin, A. Britten, and R. E. Wells. 1963. Rheology of blood and flow in the microcirculation. *J. Appl. Physiol.* 18: 255.
27. Whitmore, R. L. 1968. *Rheology of the Circulation.* Pergamon Press Ltd., Oxford.
28. Schachman, H. K. 1959. *Ultracentrifugation in Biochemistry.* Academic Press Inc., New York. 60.
29. Josephs, R., and W. F. Harrington. 1968. On the stability of myosin filaments. *Biochemistry.* 7: 2834.
30. Schachman, H. K., 1959. *Ultracentrifugation in Biochemistry.* Academic Press Inc., New York. 174.
31. Kunkel, H. G., J. B. Natvig, and F. G. Joslin. 1969. A "lepore" type of hybrid  $\gamma$ -globulin. *Proc. Nat. Acad. Sci. U. S. A.* 62: 144.
32. Frangione, B., and C. Milstein. 1968. Variation in the S-S bridges of immunoglobulin G: interchain disulfide bridges of  $\gamma$ G3 myeloma proteins. *J. Mol. Biol.* 33: 893.
33. Fahey, J. L. 1963. Serum protein disorders causing clinical symptoms in malignant neoplastic disease. *J. Chronic Dis.* 16: 703.
34. Waldenström, J. 1952. Abnormal proteins in myeloma. *In Advances in Internal Medicine*, Vol. 5. W. Dock and I. Snapper, editors. The Year Book Publishers Inc., Chicago.
35. Somer, T. 1966. The viscosity of blood, plasma and serum in dys- and paraproteinemias. *Acta Med. Scand. Suppl.* 456: 1.
36. Steel, A. E. 1963. The laboratory diagnosis of macroglobulinemia by viscometry. *Clin. Chim. Acta.* 8: 86.
37. Rees, S. B., R. A. Camerini-Cavalos, J. B. Calulfield, O. Lazano-Castaneda, A. Cervantes-Amenza, J. Taton, D. Pometta, G. P. Krauthammer, and A. Marble. 1963. The parthological physiology of microangiopathy in diabetes. *The Ciba Found. Symp. Etiology Diabetes Mellitus and its Complications*, 315.
38. Copley, A. L., and G. W. Scott Blair. 1960. Haemorheological method for the study of blood systems and of processes in blood coagulation. *In Flow Properties of Blood.* A. L. Copley and O. Stainsby, editors. Pergamon Press Ltd., Oxford. 412.
39. Charm, S., and G. S. Kurland. 1962. Tube flow behavior and shear stress/shear rate characteristics of canine blood. *Amer. J. Physiol.* 203: 417.
40. Kochwa, S., E. Smith, M. Brownell, and L. R. Wasserman. 1966. Aggregation of IgG globulin *in vivo*. II. Physicochemical properties of the isolated protein. *Biochemistry.* 5: 277.