

Effects of Fasting and Feeding on Protein Synthesis by the Rat Pancreas

J. A. Morisset, P. D. Webster

J Clin Invest. 1972;51(1):1-8. <https://doi.org/10.1172/JCI106779>.

Research Article

These experiments were designed to determine whether fasting and feeding were associated with differing rates of protein synthesis in the rat pancreas. It has been established that feeding, acetylcholine, or cholecystokinin-pancreozymin administration was associated with enhanced rates of digestive enzyme secretion; however, the literature is unclear as to effects of such stimulation on enzyme synthesis. Rats fed ad lib. or fasted 24, 48, or 72 hr were used for this study. Pancreases were removed and incubated in tissue culture medium with L-phenylalanine-¹⁴C, and incorporation into TCA-insoluble material as well as purified amylase was measured. Compared with fed controls, fasting 24, 48, and 72 hr was associated with 29%, 39%, and 35% decreases in incorporation of L-phenylalanine-¹⁴C into protein. Decreases of similar magnitudes were apparent whether the data were expressed in terms of protein or DNA. Pancreatic amylase isolated from rats fasted 48 hr contained 57% fewer counts of L-phenylalanine-¹⁴C than amylase isolated from fed rats. Moreover, rats fasted for 24 hr and given bethanechol chloride incorporated greater amounts of L-phenylalanine-¹⁴C into protein than fasted controls. Studies were performed to exclude changes in pool size of precursor (L-phenylalanine-¹⁴C) or product (amylase) in accounting for decreases associated with fasting. These studies demonstrate that fasting was associated with decreased rates of pancreatic amylase and protein synthesis in rats.

Find the latest version:

<https://jci.me/106779/pdf>



Effects of Fasting and Feeding on Protein Synthesis by the Rat Pancreas

J. A. MORISSET and P. D. WEBSTER

From the Section of Gastroenterology, Veterans Administration Hospital and Medical College of Georgia, Augusta, Georgia 30904

ABSTRACT These experiments were designed to determine whether fasting and feeding were associated with differing rates of protein synthesis in the rat pancreas. It has been established that feeding, acetylcholine, or cholecystokinin-pancreozymin administration was associated with enhanced rates of digestive enzyme secretion; however, the literature is unclear as to effects of such stimulation on enzyme synthesis. Rats fed ad lib. or fasted 24, 48, or 72 hr were used for this study. Pancreases were removed and incubated in tissue culture medium with L-phenylalanine-¹⁴C, and incorporation into TCA-insoluble material as well as purified amylase was measured. Compared with fed controls, fasting 24, 48, and 72 hr was associated with 29%, 39%, and 35% decreases in incorporation of L-phenylalanine-¹⁴C into protein. Decreases of similar magnitudes were apparent whether the data were expressed in terms of protein or DNA. Pancreatic amylase isolated from rats fasted 48 hr contained 57% fewer counts of L-phenylalanine-¹⁴C than amylase isolated from fed rats. Moreover, rats fasted for 24 hr and given bethanechol chloride incorporated greater amounts of L-phenylalanine-¹⁴C into protein than fasted controls. Studies were performed to exclude changes in pool size of precursor (L-phenylalanine-¹⁴C) or product (amylase) in accounting for decreases associated with fasting. These studies demonstrate that fasting was associated with decreased rates of pancreatic amylase and protein synthesis in rats.

INTRODUCTION

Current literature is unclear as to whether fasting or feeding is associated with changes in rates of protein synthesis in the mammalian pancreas. One group of investigators studying rats and pigeons found protein synthesis to be increased after pancreatic stimulation by

Received for publication 10 March 1971 and in revised form 26 July 1971.

feeding or administration of cholinergic drugs and pancreozymin (1-4). Another group of investigators found protein synthesis to be constant and not increased after pancreatic stimulation by feeding or administration of cholinergic drugs and pancreozymin (5-8). Resolution of this question seems basic to an understanding of acinar cell function, for if feeding is associated with increases in pancreatic enzyme synthesis, then it is reasonable to inquire as to the mechanisms whereby such increases are obtained. These experiments were performed to determine whether rates of protein synthesis in the rat pancreas were changed by fasting or feeding.

Pancreatic protein and amylase synthesis have been examined in rats which were fed, fasted, or fasted and given bethanechol. The results show that pancreases obtained from fed rats incorporated greater amounts of L-phenylalanine-¹⁴C into tissue proteins and amylase than pancreases from fasted rats. These differences were apparent whether results were expressed in terms of counts per minute/milligram protein or milligrams DNA. Moreover, pancreatic microsomes prepared from fed rats incorporated greater amounts of L-phenylalanine-¹⁴C into protein in vitro than microsomes prepared from fasted rats. Enhancement of protein synthesis by pancreases from fed rats could not be accounted for by changes involving pool size of either precursor or product. The results show that protein synthesis in the rat pancreas is variable and that it is increased after feeding or bethanechol administration. The precise mechanisms whereby these two different processes are controlled and integrated remains a problem for future investigation.

METHODS

Male Sprague-Dawley rats (250-300 g) maintained on Purina Rat Chow (Ralston Purina Co., St. Louis, Mo.) were used for all studies. Rats referred to as "fasted" were denied food for 24, 48, or 72 hr; rats referred to as "fed" had free access to food. All rats had free access to water.

The following materials were used: L-phenylalanine-¹⁴C (0.37 mCi/ μ mole) uniformly labeled (UL) (New England Nuclear Corp., Boston, Mass.; bethanechol chloride (Merck Sharp & Dohme, West Point, Pa.); materials for tissue culture media, NCTC-109, MEM, 100X, (Microbiological Associates, Inc., Bethesda, Md.); calf thymus deoxyribonucleic acid (DNA) and pancreatic alpha amylase (Worthington Biochemical Corp., Freehold, N. J.); hydroxymethylaminomethane (Tris) and orcinol (Fisher Scientific Company, Pittsburgh, Pa.); ultrafiltration cell model 402 and Diaflo Membranes PM-10 (Amicon Corp., Lexington, Massachusetts); diethylaminoethyl cellulose (DE-52) and carboxymethylcellulose (CM-52) (Scientific Division, Reeve Angel, Clifton, N. J.); Sephadex G-75 and G-100 (Pharmacia, Uppsala, Sweden); diphenylamine (Eastman Organic Chemical, Rochester, N. Y.); D-ribose (Nutritional Biochemical Corporations, Cleveland, Ohio); and bovine serum albumin (crystallized and lyophilized) (Sigma Chemical Co., St. Louis, Mo.).

Treatment of tissue. Rats were killed by decapitation, pancreases removed, fat and excess tissue trimmed, and the tissue weighed on a Roller-Smith balance and then placed in ice-cold freshly oxygenated Krebs-Ringer phosphate-buffered solution, pH 7.4. In vitro incubations were performed in 25-ml Erlenmeyer flasks placed in a shaking water bath (80 strokes/min) at 37°C under oxygen (100 mg pancreas/1 ml tissue culture medium). The tissue culture medium was prepared by mixing 50 ml of NCTC-109, 10 ml nonessential amino acids, 21.2 mg calcium chloride, 1.9 g glucose in 1 liter Krebs-Ringer phosphate-buffered (0.01 M, pH 7.4) solution. The concentration of L-phenylalanine in this medium was 48 μ M to which was added 0.02 μ Ci L-phenylalanine-¹⁴C. The incubation medium was oxygenated immediately before use.

Studies of L-phenylalanine-¹⁴C incorporation into tissue proteins. For incorporation studies, rats were either fed ad lib. or fasted 24, 48, or 72 hr. For studies examining effects of bethanechol chloride, rats fasted 48 hr were given either bethanechol chloride (2 mg/kg subcutaneous [s.c.]) or saline and killed 60 min later.

Studies of phenylalanine-¹⁴C incorporation into microsomal protein by rat pancreas. For these studies, rats were fed ad lib. or fasted 24, 48, and 72 hr. The animals were killed, and the pancreases removed and prepared as previously described. Whole pancreases were incubated at 37°C for 15 min in tissue culture medium (100 mg/1 ml medium per 0.1 μ Ci L-phenylalanine-¹⁴C). At completion of incubation, the supporting medium was aspirated, and tissue was washed quickly twice with ice-cold medium and then passed through a tissue press. The tissue was homogenized using a Teflon homogenizer (8 passes at 500 rpm) in 0.44 M sucrose (1 g tissue/10 ml sucrose). The homogenate was centrifuged 2500 rpm for 10 min (twice). The supernate was centrifuged at 11,000 rpm for 15 min (twice), and the resulting supernate was centrifuged at 43,000 rpm for 90 min in an L2-65 Beckman preparative centrifuge (50 rotor) (Beckman Instruments, Inc., Fullerton, Calif.). The supernate from the high-speed centrifugation was decanted, and the pellet was washed twice by swirling with 1 ml 0.44 M sucrose and resuspended in 2 ml of 2.1 N perchloric acid. The precipitate was washed twice with perchloric acid, once with 95% ethanol, and twice with ether-ethanol mixture. The precipitate was solubilized in 0.5 N potassium hydroxide; protein and RNA were measured as described.

Studies of L-phenylalanine-¹⁴C incorporation into amylase. Two separate experiments were performed; each experi-

ment used 10 rats, 5 fed ad lib. and 5 fasted 48 hr. The animals were killed, and pancreases were removed and incubated in tissue culture medium as described. Upon completion of in vitro incubation, pancreases of similar groups were pooled, supporting media aspirated, and tissue washed two times with ice-cold 0.02 M potassium phosphate buffer (pH 8.0). The pancreases were then suspended in 12 cc of 0.02 M potassium phosphate buffer (pH 8.0) and homogenized using a ground glass tissue grinder. Soybean trypsin inhibitor (50 mg) was added to the homogenate, and the homogenate was dialyzed overnight against 5 liters of 0.02 M potassium phosphate buffer (pH 8.0). The retentate was centrifuged again at 15,000 rpm for 15 min. The supernate was placed on a Sephadex G-75 column (45 cm \times 2.5 cm) coupled to a DEAE-52 column (30 cm \times 1.5 cm). Proteins were eluted with 0.02 M potassium phosphate buffer (pH 8.0). The eluent containing amylase activity was reduced to a 10 ml volume by ultrafiltration. The sample was then placed on a Sephadex G-100 column (45 cm \times 2.5 cm), and proteins were eluted with 0.02 M potassium phosphate buffer (pH 8.0). The eluent containing maximal amylase activity was concentrated and dialyzed by ultrafiltration against 450 ml 0.005 M potassium phosphate buffer (pH 6.2). The retentate (10 ml) was applied to a carboxymethyl-cellulose (CM-52) column (12 cm \times 1.5 cm). Proteins were eluted with 0.005 M potassium phosphate buffer (pH 6.2) and a linear gradient of NaCl (0.0–0.45 M). The eluent was concentrated by ultrafiltration; amylase activity, protein content, and radioactivity were then determined.

Studies of L-phenylalanine-¹⁴C incorporation by rat pancreatic microsomes. Six fasted (48 hr) or fed rats were used in each experiment. The pancreases were homogenized (10% w/v) in 0.44 M sucrose with a Teflon homogenizer (0.004 in. clearance, 8 passes, 500 rpm). The homogenate was centrifuged at 2500 rpm (10 min), supernate centrifuged at 11,000 rpm (15 min), and resultant supernate centrifuged at 65,000 rpm (30 min) using a Spinco L2-65B ultracentrifuge (Rotor 65). A common pooled supernatant fraction was used as a source for activating enzymes. The microsomal pellet was washed twice and resuspended in 0.25 M sucrose (0.5 ml/g tissue wet wt). Microsomes were incubated for 10 min at 37°C with an energy-generating system as described by Redman, Siekewitz, and Palade (9). Radioactivity and protein were measured. Results were expressed as counts per minute/milligram microsomal protein.

Studies of amino acid content. Two experiments were performed to measure free amino acid content of pancreases from rats fed or fasted 48 hr. Pancreases from three animals were pooled, homogenized in 1% picric acid, centrifuged, and excess picric acid removed from the supernate by passage through a Dowex 2-X10 column. The samples were lyophilized, buffered to pH 2.2, and stored at -20°C until time of assay. Amino acids were assayed by the technique of Spackman, Stein, and Moore using a Beckman 120-B amino acid autoanalyzer (10). These methods have been used previously (11).

Studies of L-phenylalanine-¹⁴C accumulation by rat pancreas. 15 fasted (48 hr) and 15 fed rats were used in each experiment; the pancreases were incubated in tissue culture medium (100 mg TWW/ml medium) containing 0.02 μ Ci/ml L-phenylalanine-¹⁴C for 5, 10, 20, 30, and 60 min. At completion of incubation, the tissue was removed from medium, blotted quickly on filter paper, and homogenized in 5% TCA. Radioactivity was measured in the media, precipitable (protein) and nonprecipitable (soluble)

TABLE I
Effects of Fasting and Feeding on L-Phenylalanine-¹⁴C Incorporation into Protein by Rat Pancreas

Fed or fasted	No. of rats	L-phenylalanine- ¹⁴ C	P value	% difference	L-phenylalanine- ¹⁴ C	P value	% difference
		<i>cpm/mg protein</i>			<i>cpm/ 100 μg DNA</i>		
Fed	6	17,661 ± 753			39,251 ± 1912		
Fasted 24 hr	6	14,483 ± 793	<0.01	18%	27,807 ± 1111	<0.01	29%
Fasted 48 hr	6	13,866 ± 1008	<0.02	21%	25,757 ± 1824	<0.001	34%
Fasted 72 hr	6	12,742 ± 743	<0.001	28%	25,671 ± 1285	<0.001	35%

Values are means ± SE. Fed or fasted rats were studied. Pancreases were incubated in 25-ml Erlenmeyer flasks containing tissue culture media (100 mg tissue/1 ml media) and L-phenylalanine-¹⁴C. Incubation was performed in a shaking water bath under oxygen at 37° C for 60 min. Reaction was stopped by addition of 1 volume of 10% TCA and incorporation into protein determined.

fractions of the tissue. These methods have been used previously (11).

Preparation of protein for counting. Precipitates were washed four times with vigorous stirring in 5 ml of 10% trichloroacetic acid containing unlabeled L-phenylalanine. The precipitate was then extracted with hot TCA, washed once with 95% ethanol and twice with ether-ethanol (3:1). At completion of the washing procedure, the precipitate was dissolved in 3 ml of 0.5 N KOH and heated at 90°C for 20 min. Samples of this solution were placed in glass counting vials along with 10 ml of scintillation mixture, and radioactivity was determined as previously described (11).

Protein was assayed by either the biuret (12) or Lowry, Rosebrough, Farr, and Randall method (13). Amylase activity was assayed by the Bernfeld method using lintner starch as substrate (14). A unit of amylase activity represents that amount which catalyzes the formation of 1 mg of maltose from starch in 3 min at 37°C. RNA and DNA were separated by modifications of methods described by Schneider and Schmidt and Tannhauser (15-17). RNA was assayed by the orcinol method using ribose as the standard; DNA was assayed by the diphenylamine method using calf thymus DNA as the standard (18).

RESULTS

Table I shows effects of feeding and fasting on L-phenylalanine-¹⁴C incorporation into tissue protein by rat pancreas incubated in vitro. Compared with fed rats, fasting for 24, 48, and 72 hr was associated with an 18%, 21%, and 28% decrease in L-phenylalanine-¹⁴C incorporation into protein per milligram protein. When the data were expressed as cpm in protein/100 μg DNA, fasting 24, 48, and 72 hr was associated with a 29%, 39%, and 35% decrease in incorporation. Differences between fed and fasted animals were statistically significant.

Fig. 1 shows rates of L-phenylalanine-¹⁴C incorporation into tissue proteins by pancreatic tissue obtained from rats fed ad lib. or fasted 48 hr. Incorporation, expressed as cpm/mg protein or cpm/100 μg DNA, was linear for periods of incubating ranging from 10 to 120 min. For all time periods studied, fed rats showed

greater incorporation of L-phenylalanine-¹⁴C into protein than fasted ones.

Table II shows effects of feeding and fasting on phenylalanine-¹⁴C incorporation into microsomal protein by whole pancreas incubated in vitro for 15 min. These experiments utilizing short periods of in vitro incubation as well as examining incorporation into microsomal proteins were performed to minimize effects of changes in pool size of product (protein), which might be associated with feeding or fasting. The table shows that pancreases from animals fasted 24, 48, and 72 hr incorporated fewer counts into protein when the data were expressed in terms of counts per minute incorporated into protein per milligram protein or milligram RNA. The increased RNA incorporation (+ 8%) into protein per milligram RNA found in the




FIGURE 1 Rate of L-phenylalanine-¹⁴C incorporation by pancreases from rats fed ad lib. or fasted 48 hr. Data expressed in terms of incorporation into protein per 100 μg DNA (top) or mg protein (bottom).

TABLE II
Effects of Fasting and Feeding on L-Phenylalanine-¹⁴C Incorporation into Microsomal Protein

Treatment	No. of exp.	L-phenylalanine- ¹⁴ C	P value	% difference	L-phenylalanine- ¹⁴ C	P value	% difference	mg protein mg RNA
		<i>cpm/mg protein</i>			<i>cpm/mg RNA</i>			
Fed ad lib.	3	10,563		—	90,073		—	9.7
Fasted 24 hr	4	8,445	NS	-20%	97,724	NS	+8%	11.3
Fasted 48 hr	4	5,945	<0.025	-44%	65,120	<0.050	-28%	11.0
Fasted 72 hr	3	4,334	<0.010	-59%	48,624	<0.050	-46%	10.1

Values are means. Fed or fasted rats studied. Pancreases were incubated for 15 min in tissue culture medium containing L-phenylalanine-¹⁴C. At completion of incubation, the medium was aspirated, the tissue homogenized, and a microsomal pellet prepared by centrifugation. Data expressed as cpm in protein/mg protein and cpm in protein/mg RNA. RNA expressed as RNA-ribose.

24 hr fasted groups while reproducible was without explanation. The ratio of microsomal protein to RNA (milligram protein/milligram RNA) ranged from 9.7 to 10.1 for the different treated groups. Recoveries of microsomal RNA compared with whole homogenate RNA were 21%, 24%, 26%, and 24% for fed ad lib., 24, 48, and 72 hr fasted groups, respectively.

Experiments in Table III demonstrate that pancreases obtained from fed rats incorporated greater amounts of L-phenylalanine-¹⁴C into amylase than pancreases from fasted rats. After a 48 hr fast, there was a 57% decrease in incorporation per milligram protein and a 41% decrease in incorporation per unit amylase activity. Amylase content as well as amounts of amylase applied to columns were approximately equal for fed and 48-hr fasted rats (28,000 and 29,000 U/g pancreas). Recoveries of amylase from total homogenate were similar for each group (60% and 62). The final preparation of amylase had an enzyme specific activity of 17,000 U/mg protein and gave only one band on gel electrophoresis.

TABLE III
Incorporation of L-Phenylalanine-¹⁴C into Pancreatic Amylase

	cpm mg protein	% difference	cpm unit amylase	% difference
Fed	40,178		32.9	
Fasted	17,273	-57%	19.4	-41%

Pancreases obtained from rats fed ad lib. or fasted 48 hr were used. The tissue was incubated in vitro under oxygen for 60 min at 37°C, homogenized, and amylase isolated using column chromatography. cpm/mg protein refers to radioactivity in amylase fraction; cpm/unit amylase refers to radioactivity in amylase fraction per unit amylase activity.

Table IV shows effects of fasting and feeding on L-phenylalanine-¹⁴C incorporation in vitro by rat pancreatic microsomes. For these studies, microsomes and supernate prepared from fed or fasted animals were incubated in vitro. Factors necessary for in vitro protein synthesis were isolated in either the cell sap fraction (supernatant portion) or the particulate fraction (microsomal portion). By incubating supernate and microsomes in a determined order, one can obtain an indication of whether changes in rates of protein synthesis result from changes in either the particulate or supernatant fraction. Microsomes and supernate prepared from fed animals incorporated greater amounts of L-phenylalanine-¹⁴C than microsomes and supernate prepared from fasted animals (44%). Microsomes from fed rats incubated with supernate from fasted rats incorporated 23% fewer counts than microsomes and supernate from fed rats. Likewise, microsomes and supernate from fasted rats incorporated fewer counts

TABLE IV
Effects of Fasting and Feeding on L-Phenylalanine-¹⁴C Incorporation In Vitro by Rat Pancreatic Microsomes

Microsomes	Supernate	cpm/mg microsomal protein	% difference	P value
Fed	+ Fed	560 ± 120		
Fed	+ Fasted	427 ± 157	+23.8%	<0.05
Fasted	+ Fasted	313 ± 28		
Fasted	+ Fed	411 ± 48	+23.8%	<0.01

Values are means ±SD (four experiments). Pancreatic microsomes prepared from rats fed or fasted 48 hr were incubated in vitro with pooled supernatant fractions; energy-generating system and incorporation into TCA-insoluble material were determined.

TABLE V

Relationship between Tissue Wet Weight, Protein, and DNA Content in Pancreas from Fed and Fasted Rats

	mg tissue wet weight	mg protein
	mg DNA	mg DNA
Fed	132.1 ± 66	14.7 ± 1.4
Fasted 24 hr	133.6 ± 58	24.0 ± 1.7
Fasted 48 hr	99.7 ± 50	13.3 ± 0.3
Fasted 72 hr	80.9 ± 36	10.7 ± 0.4

Values are means ± SE. Pancreas removed from fed or fasted rats, weighed on a Roller-Smith balance, minced, and homogenized. Protein and DNA determined as described in the Methods section.

than microsomes from fasted rats incubated with supernate from fed rats. This 23% greater incorporation in counts appeared to be due to ingredients present in the fed supernate. It seems in fed animals that enhancement of amino acid-¹⁴C incorporation into protein results from changes at both the microsomal as well as supernatant level. It is known that microsomes contain messenger RNA whereas cytoplasmic fractions contain activating enzyme, t-RNA, and transfer factors. Studies are being performed to determine which of these factors was responsible.

Table V shows relationships between tissue wet weight, protein, and DNA content of pancreas from fed and fasted rats. Tissue wet weight per milligram DNA was similar for fed and 24-hr fasted rats. Fasting 48 and 72 hr was associated with 24% and 39% decreases in tissue wet weight. There was an initial increase in protein content per milligram DNA with fasting 24 hr but a decrease thereafter (48 and 72 hr). Protein content per unit DNA was approximately equal in fed ad lib. and 48-hr fasted rats.

It was previously shown using pigeons that administration of analogues of the neurohormone, acetylcholine, was associated with enhancement of protein synthesis. If decreased protein synthesis in the pancreas after fasting were, in part, due to decreased neurohormonal stimulation of the pancreas, then administration of an analogue of the neurohormone, acetylcholine, should be associated with an increase in protein synthesis comparable to that observed in the fed animal. If, on the other hand, decreases in protein synthesis were due to "unknown dietary factors," then bethanechol chloride administration should not be associated with increased pancreatic protein synthesis. Fig. 2 shows rates of L-phenylalanine-¹⁴C incorporation into pancreatic tissue proteins by pancreases obtained from rats fasted 48 hr and then given either saline or bethanechol chloride. Bethanechol chloride administration to rats fasted 48




FIGURE 2 Rate of L-phenylalanine-¹⁴C incorporation by pancreas from rats fasted 48 hr given bethanechol chloride 0.2 mg/kg or saline s.c.

hr was clearly associated with a demonstrable increase in L-phenylalanine-¹⁴C incorporation whether the data were expressed in terms of protein or DNA. The difference in slopes of lines of the bethanechol-treated group most likely reflects secretory losses of protein.

The next series of experiments was designed to examine possibilities that changes in L-phenylalanine-¹⁴C incorporation observed with feeding or fasting might be due to changes in pool size or uptake of amino acids by the pancreatic acinar cell. Table VI shows content of L-phenylalanine per gram tissue wet weight in pancreases removed from fed or fasted (24 hr) rats. Phenylalanine content was slightly greater in pancreases of fasted than fed rats. These values agree with values previously obtained for pigeon pancreas. L-Phenylalanine content of the incubating medium was 4800 µmoles/ml or about 100 times greater than the concentration of

TABLE VI

L-Phenylalanine Content of Pancreases from Fed or Fasted Rats

	µmoles phenylalanine
	g tissue wet weight
Fed	58
Fasted 48 hr	60

Pancreases from four fed or fasted rats were pooled, minced, and homogenized in 1% picric acid, and amino acid content of the soluble portion was determined using amino acid autoanalyzer.




FIGURE 3 Pancreases from fed or fasted rats were incubated in vitro, and measurement of L-phenylalanine-¹⁴C in media, tissue water, and protein was performed.

free L-phenylalanine in tissue water. Thus, if it were shown that intracellular concentrations of amino acids were rapidly diluted by amino acids in the medium, it would be highly unlikely that such minor differences could account for differences in L-phenylalanine-¹⁴C incorporation by pancreases from fed or fasted rats.

Fig. 3 shows data obtained when pancreases from fed and fasted rats were incubated 5, 10, 20, 30, and 60 min in vitro, and amounts of L-phenylalanine-¹⁴C in

medium, tissue water, and protein were determined. After 60 min of in vitro incubation, the concentration of L-phenylalanine-¹⁴C in medium containing pancreases from fasted rats was greater than that of medium containing pancreases from fed rats. The concentration of L-phenylalanine-¹⁴C in tissue water of pancreases from fasted and fed rats was equal after 5 and 10 min of incubation, but after 20, 30, and 60 min of incubation, the concentration of label was greater in tissue water of fasted rats. Rates of incorporation of L-phenylalanine-¹⁴C into protein were greater in fed than fasted rats for all periods of incubation. Recovery studies for these experiments demonstrated accountability of up to 98% of label in either the media, tissue water, or protein. In summary, these studies show greater disappearance of label from the medium and greater incorporation into protein by pancreases obtained from fed rats. There was also rapid equilibration of amino acids in medium with that in tissue water.

DISCUSSION

It has been well established that feeding, sham feeding, administration of cholecystokinin-pancreozymin, or cholinergic drugs results in initiation of secretion of pancreatic digestive enzymes (19). However, effects of such stimulation on pancreatic protein synthesis has been unclear. In fact, two somewhat opposing points of view have described the relationship between synthesis and secretion in the pancreas. One view suggested that after secretion there was synthesis of new product. The second view suggested that protein synthesis was constant, that secretion was inconstant or variable, and that variations in enzyme content resulted only from changes in secretion.

Reconciliation of these differing points of view has been provided by recent studies comparing effects of in vivo and in vitro administration of bethanechol chloride, cholecystokinin-pancreozymin, and cyclic AMP (20). In these studies, it was shown that administration of these agents in vivo was associated with initiation of both secretory and synthetic processes, whereas, administration of these agents in vitro was associated only with initiation of secretion. Clearly, both points of view are correct as related to the type of experimental model under study by the investigator.

However, the question of whether feeding and fasting are associated with differing rates of pancreatic digestive enzyme synthesis has not been resolved. Poort and Kramer recently published experiments in which they were unable to show differences in rates of incorporation of a ¹⁴C-labeled amino acid into tissue proteins by pancreases obtained from fed or fasted rats. These investigators concluded that pancreatic protein synthesis in the mammalian pancreas was constant and that feed-

ing and fasting were not associated with changes in rates of pancreatic enzyme synthesis (8).

Their observations, utilizing rat pancreas, were at variance with experiments which had utilized pigeon pancreas. Pancreases from fed pigeons compared with pancreases from fasted pigeons demonstrated the following: greater incorporation of palmitate-¹⁴C into triglycerides and phospholipids, greater rates of oxidation of palmitate-¹⁴C to ¹⁴CO₂, greater rates of incorporation of uridine-³H into RNA, and greater rates of incorporation of L-phenylalanine-¹⁴C into proteins (21-23).

The present studies have examined rates of amino acid-¹⁴C incorporation into tissue proteins, microsomal proteins, and amylase by pancreases obtained from rats which were fed ad lib. or fasted 24, 48, and 72 hr. The data show that pancreases from fed rats incorporated greater amounts of L-phenylalanine-¹⁴C into tissue and microsomal protein than rats fasted 24, 48, or 72 hr. An enhancement of L-phenylalanine-¹⁴C incorporation was demonstrated whether the data were expressed in terms of milligram protein or microgram DNA. Moreover, administration of bethanechol chloride, an analogue of acetylcholine, to rats fasted 48 hr was associated with enhancement of L-phenylalanine-¹⁴C incorporation into protein at rates almost equal to those seen in fed animals. When pancreatic amylase was isolated from fed rats, there was greater specific radioactivity of L-phenylalanine-¹⁴C per milligram protein as well as per unit amylase. In addition, pancreatic microsomes isolated from fed rats and incubated in vitro with an energy-generating system incorporated greater amounts of label than microsomes from fasted rats. Moreover, experiments incubating cell sap and microsomes in random order suggested that increases associated with feeding resulted from changes at ribosomal as well as cytoplasmic levels of control.

A number of possible sources of error have been eliminated by design of these experiments. Variations in accumulation of amino acids, an indication of transport, by pancreases from fasted or fed animals could not account for changes in incorporation since accumulation of L-phenylalanine-¹⁴C at 5-20 min was almost identical for tissue obtained from either fasted or fed rats.

Variations in pool size or tissue content of amino acids between fasted and fed animals could not account for increased incorporation of L-phenylalanine-¹⁴C into protein since pool sizes of phenylalanine were almost equal in pancreases from fed or fasted rats. Moreover, by studying incorporation in vitro, intracellular pools of amino acids were rapidly equilibrated with much higher concentrations of amino acids in the incubating media. Hanking and Roberts have shown that intracellular concentrations of amino acids in tissue slices rapidly equilibrate with that of the supporting media (24). There

were almost linear increases in L-phenylalanine-¹⁴C incorporation despite the rapid changes in concentration of L-phenylalanine-¹⁴C in intracellular tissue water.

Another possible source of error might be due to secretory losses of pancreatic protein after feeding. In other words, could the apparent increase in incorporation result from decreases in milligrams protein of the tissue (a change in denominator rather than numerator)? However, protein content of pancreases obtained from animals fed ad lib. and animals fasted 48 hr was almost equal. In addition, decreases in rates of incorporation with fasting were apparent whether the data were expressed in terms of DNA or protein. Moreover, it should be pointed out that although the pancreas secretes large amounts of enzyme when expressed in terms of enzyme activity, such activity reflects only minor changes in protein content of the entire gland. For example, 2000 U of amylase activity represents 1-1.5 mg of protein. Secretory studies using rats have shown that maximal secretory rates obtained with in vivo pancreozymin or bethanechol elicit secretion of only 2-4 mg protein/hr (25). Such secretory losses of protein represent a small change compared with the total weight of the organ (2-4 mg protein vs. 750-1000 mg tissue wet wt).

In vitro incubation of pancreas and isolation of microsomes from rats demonstrated 25, 45, and 60% decreased incorporation of L-phenylalanine-¹⁴C into microsomal protein with 24, 48, and 72 hr of fasting. These experiments, utilizing a short incubation period (15 min) as well as isolation of subcellular sites of protein synthesis (ribosomes), were performed to minimize effects of changes in pool size of product (protein), which might alter results. The results of these experiments agree with results obtained when L-phenylalanine-¹⁴C incorporation into whole tissue protein was examined.

Incubation of microsomes prepared from fed and fasted rats with an energy-generating system demonstrated fasting was associated with decreased amino acid-¹⁴C incorporation into microsomal protein. The data suggest that such changes result from alterations involving both microsomal and cytoplasmic components. Differences in amino acid-activating enzymes, t-RNA, and transfer enzymes (pH 5 fraction) may account for changes at the cytoplasmic level. Differences at the microsomal level may reflect programming of the ribosomes with messenger RNA. Studies are underway to better define these changes.

In conclusion, these investigations strengthen the view that pancreatic protein synthesis is a variable process and that augmentation of secretion is frequently associated with augmentation of synthesis. The precise manner whereby such control is maintained and integrated is unknown. Moreover, the role that such com-

plex mechanisms of control may play in human disease remains undefined.

ACKNOWLEDGMENTS

We would like to acknowledge the technical assistance of Mrs. Marilyn LaSure and Mrs. Nancy Parks and the secretarial assistance of Mrs. Elizabeth Spainhour.

REFERENCES

1. Farber, E., and H. Sidransky. 1956. Changes in protein metabolism in the rat pancreas on stimulation. *J. Biol. Chem.* **222**: 236.
2. Fernandes, J. F., and L. C. V. Junqueira. 1955. Protein and ribonucleic acid turnover rates related to activity of digestive enzymes of pigeon pancreas. *Arch. Biochem.* **55**: 54.
3. Webster, P. D. 1969. Effect of stimulation on pancreatic amylase secretion and nuclear RNA synthesis. *Proc. Soc. Exp. Biol. Med.* **132**: 1072.
4. Webster, P. D., and M. P. Tyor. 1966. Effects of intravenous pancreozymin on amino acid incorporation in vitro by pancreatic tissue. *Amer. J. Physiol.* **211**: 157.
5. Hokin, L. E., and M. R. Hokin. 1961. The synthesis and secretion of digestive enzymes by pancreas tissue in vitro. In *The Exocrine Pancreas*. A. V. S. deReuck and M. P. Cameron, editors. Little, Brown and Company, Boston. 186.
6. Dickman, S. R., R. L. Holtzer, and G. Gazzinelli. 1962. Protein synthesis by beef pancreas slices. *Biochemistry.* **1**: 574.
7. Schucker, R., and L. E. Hokin. 1954. The synthesis and secretion of lipase and ribonuclease by pigeon pancreas slices. *J. Biol. Chem.* **210**: 551.
8. Poort, C., and M. F. Kramer. 1969. Effect of feeding on protein synthesis in mammalian pancreas. *Gastroenterology.* **57**: 689.
9. Redman, C. M., P. Siekewitz, and C. E. Palade. 1966. Synthesis and transfer of amylase in pigeon pancreatic microsome. *J. Biol. Chem.* **241**: 1150.
10. Spackman, D. H., W. H. Stein, and S. Moore. 1958. Automatic recording apparatus for use in the chromatography of amino acids. *Anal. Chem.* **30**: 1190.
11. Webster, P. D. 1968. Effect of methacholine on pancreatic amylase synthesis. *Gastroenterology.* **55**: 375.
12. Gornall, A. G., C. J. Bardawill, and M. M. David. 1949. Determination of serum proteins by means of the biuret reaction. *J. Biol. Chem.* **177**: 751.
13. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**: 265.
14. Bernfeld, P. 1955. Amylases α and β . *Methods Enzymol.* **1**: 149.
15. Schneider, W. C. 1957. Determination of nucleic acids in tissues by pentose analysis. *Methods Enzymol.* **3**: 680.
16. Schmidt, G., and S. J. Tammhauser. 1945. A method for the determination of deoxyribonucleic acid, ribonucleic acid, and phosphoproteins in animal tissues. *J. Biol. Chem.* **161**: 83.
17. Schneider, W. C. 1945. Phosphorus compounds in animal tissues. I. Extraction and estimation of desoxypentase nucleic acid and of pentose nucleic acid. *J. Biol. Chem.* **161**: 293.
18. Volkin, E., and W. E. Cohn. 1954. Estimation of nucleic acids. *Methods Biochem. Anal.* **1**: 287.
19. Dreiling, D. A., and H. D. Janowitz. 1961. The measurement of pancreatic secretory function. In *The Exocrine Pancreas*. A. V. S. deReuck and M. P. Cameron, editors. Little, Brown and Company, Boston. 225.
20. Morisset, J. A., and P. D. Webster. 1971. In vitro and in vivo effects of pancreozymin, urecholine, and cyclic AMP on rat pancreas. *Amer. J. Physiol.* **230**: 202.
21. Webster, P. D., and M. P. Tyor. 1966. Effect of fasting and feeding on lipid metabolism of pigeon pancreas. *Amer. J. Physiol.* **210**: 1076.
22. Webster, P. D., and M. P. Tyor. 1967. Effects of fasting and feeding on uridine-³H incorporation into RNA by pancreas slices. *Amer. J. Physiol.* **212**: 203.
23. Webster, P. D., and M. P. Tyor. 1966. Effect of intravenous pancreozymin on amino acid incorporation in vitro by pancreatic tissue. *Amer. J. Physiol.* **211**: 157.
24. Hanking, B. M., and S. Roberts. 1964. Influence of amino acid levels on protein synthesis in vitro. *Nature (London)*. **204**: 1194.
25. Leroy, J., J. A. Morisset, and P. D. Webster. 1971. Dose related response of pancreatic synthesis and secretion to cholecystokinin-pancreozymin. *J. Lab. Clin. Med.* **78**: 149.