

Chronic cobalamin inactivation impairs folate polyglutamate synthesis in the rat.

J Perry, ... , R Deacon, M Lumb

J Clin Invest. 1983;71(5):1183-1190. <https://doi.org/10.1172/JCI110867>.

Research Article

Nitrous oxide, by inactivating cobalamin in vivo, produces a suitable animal model for cobalamin 'deficiency.' The synthesis of folate polyglutamate with tetrahydrofolate as substrate is severely impaired in the N₂O-treated rat, but is normal with formyltetrahydrofolate as substrate. Methionine restores the capacity of the N₂O-treated rat to utilize tetrahydrofolate the minimum effective dose being 16 μmol. S-Adenosylmethionine was somewhat less effective than methionine but 5-methylthioadenosine, a product of S-adenosylmethionine metabolism, was significantly more effective than methionine in correcting the defect in folate polyglutamate synthesis. 5-Methylthioadenosine is metabolised to yield formate. It is suggested that these compounds have their effect in correcting folate polyglutamate synthesis by supplying formate for the formylation of tetrahydrofolate. Formyltetrahydrofolate, at least in the cobalamin-inactivated animal, is the required substrate for folate polyglutamate synthesis. Cobalamin is concerned with the maintenance of normal levels of methionine and this in turn is a major source of formate through S-adenosylmethionine and 5-methylthioadenosine.

Find the latest version:

<https://jci.me/110867/pdf>



Chronic Cobalamin Inactivation Impairs Folate Polyglutamate Synthesis in the Rat

JANET PERRY, I. CHANARIN, ROSEMARY DEACON, and M. LUMB, *Section of Haematology, Medical Research Council, Clinical Research Centre, Northwick Park Hospital, Harrow, Middlesex, United Kingdom*

ABSTRACT Nitrous oxide, by inactivating cobalamin in vivo, produces a suitable animal model for cobalamin 'deficiency.' The synthesis of folate polyglutamate with tetrahydrofolate as substrate is severely impaired in the N₂O-treated rat, but is normal with formyltetrahydrofolate as substrate. Methionine restores the capacity of the N₂O-treated rat to utilize tetrahydrofolate the minimum effective dose being 16 μmol. S-Adenosylmethionine was somewhat less effective than methionine but 5-methylthioadenosine, a product of S-adenosylmethionine metabolism, was significantly more effective than methionine in correcting the defect in folate polyglutamate synthesis. 5-Methylthioadenosine is metabolised to yield formate. It is suggested that these compounds have their effect in correcting folate polyglutamate synthesis by supplying formate for the formylation of tetrahydrofolate. Formyltetrahydrofolate, at least in the cobalamin-inactivated animal, is the required substrate for folate polyglutamate synthesis. Cobalamin is concerned with the maintenance of normal levels of methionine and this in turn is a major source of formate through S-adenosylmethionine and 5-methylthioadenosine.

INTRODUCTION

The anesthetic gas, nitrous oxide (N₂O), inactivates cobalamin (vitamin B₁₂) and has proved an extremely useful tool in investigating the biochemical effects of loss of cobalamin activity. On exposure to transition-metal complexes, of which cobalamin is one, N₂O, which is otherwise inert, is cleaved into nitrogen and oxygen (1). It thus acts as a two-electron oxidizing

agent and oxidizes cob[II]alamin to the inactive cob[III]alamin. In man too, exposure to N₂O inactivates cobalamin and, if exposure exceeds 6 h, megaloblastic hemopoiesis and marrow failure (2-4), which can be fatal (2, 5), develops. Prolonged intermittent exposure leads to a neuropathy essentially similar to that found in cobalamin deficiency (6).

Exposure of animals to N₂O leads to inactivation of the enzyme methionine synthetase (7), which has a methylcobalamin cofactor, but there is no increase in urinary methylmalonic acid excretion, which too requires cobalamin. However, long-term exposure of rats to N₂O is accompanied by a fall in hepatic methylmalonylCoA mutase activity (8).

The inactivation of cobalamin by N₂O also leads to major effects on folate metabolism. There is a rise in plasma folate on exposure to N₂O in both man (4) and rat (9) and a marked loss of folate from the tissues in rats (10, 11) due to excretion of 5-CH₃-H₄-pteroyl-glutamate (5-CH₃-H₄PteGlu) into the urine (12). Recovery of tissue folate on returning N₂O-exposed rats to air requires a dietary source of folate (13). There is an increased urinary formiminoglutamic acid excretion in the N₂O-treated rat (unpublished observations) but the activity of the enzyme, formiminotetrahydrofolate formiminotransferase, is unaffected (14). Exposure to N₂O also leads to impaired synthesis of thymidine, as assessed in the deoxyuridine suppression test, in both rats (15, 16) and man (3), and this impairment is corrected in part by the addition of cobalamin to the test system.

Exposure to N₂O also affects other enzymes concerned with folate metabolism. There is a rise in activity of 10-formyltetrahydrofolate synthetase (linking formate to H₄PteGlu), a fall in activity of 5,10-methylenetetrahydrofolate cyclohydrolase (10-CHO-H₄-PteGlu → 5,10-CH=H₄PteGlu) (14) and an increase in activity of thymidylate synthetase (17). Other enzymes, including serine transhydroxymethylase, 5,10-methylenetetrahydrofolate dehydrogenase, and 5-

Portions of these studies were presented at 18th Congress of the International Society for Hematology, Montreal, Canada, August 1980.

Received for publication 25 May 1982 and in revised form 6 December 1982.

10-methylenetetrahydrofolate reductase, are unaffected (14).

In untreated pernicious anemia there is a fall in the concentration of folate polyglutamate in erythrocytes without any change in the level of folate monoglutamate (18, 19). An impaired incorporation of PteGlu into hepatic folate polyglutamate has been demonstrated in sheep (20) and rats (21) with dietary-induced cobalamin deficiency. The N₂O-treated rat likewise showed an impairment in the synthesis of folate polyglutamate from parenteral PteGlu (15) as well as from H₄PteGlu (22). However, folate polyglutamate synthesis was normal when the N₂O-treated rats were given either 5-CHO-H₄PteGlu (folinic acid) or 10-CHO-H₄PteGlu as substrate (22). This suggested that formyltetrahydrofolates rather than tetrahydrofolate were the required substrates for folate polyglutamate synthesis in rats and that cobalamin was concerned with the provision of CHO-H₄PteGlu. This view was reinforced by the increased activity of formyltetrahydrofolate synthetase following exposure to N₂O (14).

The purpose of this study was to explore in greater detail the synthesis of folate polyglutamate in control and N₂O-treated rats and to determine how this could be related to the provision of 'formate' by testing the effect of compounds that might contribute a single carbon unit into the folate pool.

METHODS

Materials. L-Methionine, S-adenosyl-L-methionine (SAM)¹, 5'-methylthioadenosine (MTA), adenosine, L-serine, glycine, betaine, sarcosine, choline, and L-homocysteine thiolactone hydrochloride were obtained from Sigma, Poole, Dorset, U.K. [2-¹⁴C]PteGlu (Radiochemical Centre, Amersham, U.K.), sp act ~52 μCi/μmol was reduced to [2-¹⁴C]H₄PteGlu using sodium dithionite (23). Column chromatography of the product on DE52 cellulose showed the yield of H₄PteGlu to be between 82–85% of PteGlu. The remaining carbon-labeled fractions, which eluted in fractions 5–15, were not microbiologically active and appeared to be impurities in the [2-¹⁴C]PteGlu eluting from the column before folate compounds. The ¹⁴C-labeled fractions eluting in fractions 5–15 were concentrated, 0.5 μCi injected into rats, both air-breathing and N₂O-breathing, animals killed and livers removed at interval up to 24 h. No labeled activity was detectable in liver. We have shown elsewhere that these impurities are excreted into the urine (12). [2-¹⁴C]H₄PteGlu, made the day before use, was stored overnight at -30°C under nitrogen and was used without further purification. There was no deterioration of [2-¹⁴C]H₄PteGlu kept under these conditions for at least 9 d after preparation.

Animals. Male, 4–5-wk Sprague-Dawley rats, 80–120 g, were used in all experiments. For each compound to be tested three animals were placed for 18 h in a perspex chamber in which CO₂ and excess water vapour were removed, and through which was passed N₂O/O₂ (vol/vol 1:1). They were then injected intraperitoneally with [2-¹⁴C]H₄PteGlu

plus the test compound and returned immediately to the chamber. Control animals receiving [2-¹⁴C]H₄PteGlu alone were treated identically. A further group of control rats were left in air but received the same compounds. All animals were allowed water and standard commercial rat diet ad lib. 24 h following injection the animals were killed by exsanguination following intraperitoneal nembutal and the livers removed rapidly. A 24-h time period was chosen because folate polyglutamate reaches maximal levels at this time (51).

Preparation of doses. Unless stated otherwise the administered injections comprised 16 μmol of the compound under study, together with 1 μCi (0.02 μmol) of [2-¹⁴C]DL,H₄PteGlu in a total volume of 0.5 ml. Larger doses of L-methionine were administered in a 1.0 ml volume. 5-MTA was dissolved by heating in a water bath at 75°C for 15 min and syringes containing the measured dose were left at 37°C immediately before injection. Adenosine was warmed gently before administration, as was L-methionine at concentrations of 125 and 250 μmol.

Extraction of folates, column chromatography, and identification. Immediately on removal livers were weighed, diced with a sharp knife, and placed in 5 vol of 1% potassium ascorbate pH 7.0 in a boiling water bath for 5 min, to destroy conjugase enzyme activities. Thereafter the suspension was placed in an ice bath and homogenized. 0.2 ml of this homogenate was solubilized in NCS (Amersham Corp., Arlington Heights, IL) scintillation fluid (toluene 1 liter, PPO 6 g, POPOP 0.075 g) added and counted for radioactivity in an LKB-Wallace liquid scintillation counter (LKB Producter, Bromma, Sweden) with external standardization to determine total hepatic uptake of labeled folate. Homogenate equivalent to 2 g liver (10 ml) was diluted with an equal volume of 0.1 M potassium phosphate buffer pH 5.7 containing 0.1% ascorbic acid, autoclaved for 10 min at 10 psi and centrifuged. The clear supernatant was applied to a 40 × 0.9-cm column of DE52 cellulose (Whatman Ltd., Maidstone, Kent, U.K.). Columns were run at 4°C, and folates were eluted using a nonlinear gradient of potassium phosphate buffer pH 6.0 containing 0.2 M 2-mercaptoethanol. The gradient was made using 150 ml of 0.1 M phosphate in the mixing chamber and 500 ml 1.0 M phosphate in the reservoir. 5-ml fractions were collected and 1 ml of fractions 20–65 counted for radioactivity after addition of 5 ml of a micellar scintillation fluid (NE260, Nuclear Enterprises Lighthill Edinburgh, Scotland). The proportion of the radioactive dose converted into folate polyglutamate was expressed as a percentage of the total radioactive folate eluted from the column, using the formula dpm in polyglutamate fractions/dpm in fractions 20–65 × 100. Radioactivity in the folate fractions is shown in the chromatograms as a percentage of the intraperitoneal dose incorporated into a gram of liver i.e., (dpm/fraction ÷ 2)/(dpm/g liver) × 100.

Folates in the eluant fractions were identified by their position of elution from the column and by cochromatography with known folate compounds. The monoglutamate standards used were [C-³H]5-CHO-H₄PteGlu, prepared by the Radiochemical Centre, Amersham, from 5-CHO-H₄PteGlu (Lederle Laboratories, Pearl River, NY) by exchange with tritiated water in the presence of a platinum catalyst and purified as described by Nixon and Bertino (24), and 5[¹⁴C]CH₃-H₄PteGlu (Amersham Corp.). The elution position of polyglutamyl folates of chain lengths 4–6 were determined by comparison with those obtained from columns previously standardized with marker compounds of 5CH₃ [³H]H₄PteGlu_{4–6} made by incubation of *Lactobacillus casei* in a medium containing [³H]PteGlu (25). Column eluates were also assayed microbiologically with *L. casei*

¹ Abbreviations used in this paper: MTA, methylthioadenosine; SAM, S-adenosylmethionine.

(ATCC 7469) and *Pediococcus cerevisiae* (ATCC 8081). Polyglutamyl folates were made available to the assay organisms by deconjugation to monoglutamyl forms following incubation of 0.1 ml of each fraction with 0.05 ml of human plasma (which contains conjugase enzyme activity) in 1 ml 0.1% ascorbate pH 4.5.

RESULTS

Hepatic content of [^{14}C]H $_4$ PteGlu. The amount of labeled folate in the liver 24 h after 0.02 μmol parenteral H $_4$ PteGlu averaged $9.6 \pm 0.4\%$ (mean \pm SEM) in 15 air-breathing rats and was $2.0 \pm 0.01\%$ in 32 rats inhaling N $_2$ O. This difference was highly significant ($P = <0.001$). There was an improvement in hepatic uptake of H $_4$ PteGlu with increasing doses of intraperitoneal methionine (Fig. 1) but even with 250 μmol the uptake was 5.7% of the dose of H $_4$ PteGlu, which was still well below that in air-breathing controls.

The effect of an injection of 16 μmol of various additives on the hepatic content of labeled folate of rats inhaling N $_2$ O expressed as a percentage of the injected dose of [^{14}C]H $_4$ PteGlu is as follows: serine, mean \pm SEM, 4.4 ± 0.4 ; betaine 3.9 ± 0.5 ; sarcosine 2.1 ± 0.2 ; choline 1.4 ± 0.1 ; homocysteine 1.7 ± 0.3 ; glycine 3.7 ± 1.5 ; adenosine 1.1 ± 0.1 ; SAM 2.6 ± 0.1 and MTA 3.0 ± 0.6 . There were three rats in each group other than MTA where eight animals were studied. There were no significant differences.

Hepatic folate polyglutamate synthesis from [^{14}C]H $_4$ PteGlu. An average of $57.5 \pm 1.4\%$ of H $_4$ PteGlu in liver was converted into polyglutamate in 15 control animals (Table I). By contrast, there was no detectable

folate polyglutamate formation from H $_4$ PteGlu in 18 out of 32 rats given N $_2$ O (Fig. 2) and the mean polyglutamate synthesis in the whole group was $5.3 \pm 1.3\%$. The difference was highly significant ($P = <0.001$).

Effect of methionine on folate polyglutamate synthesis. The impairment in the synthesis of folate polyglutamate from H $_4$ PteGlu in N $_2$ O-treated rats was corrected by the administration of methionine (Fig. 3). Folate polyglutamates elute in the fractions collected after tube 40. 'Titration' of the dose of methionine required to produce maximum folate polyglutamate synthesis (Fig. 4) indicated that this was of the order of 16 μmol . There was no further increase in polyglutamates formed even with amounts of 250 μmol methionine. By contrast, this large dose of methionine did not fully correct the impaired hepatic uptake of folate (Fig. 1). On the basis of the studies set out in Fig. 4, 16 μmol was selected as the dose level at which the effect of compounds in restoring folate polyglutamate synthesis in N $_2$ O-treated animals was tested. At a dose of 16 μmol methionine $48.0 \pm 3.1\%$ of labeled hepatic folate was in the polyglutamate form in the N $_2$ O-treated rat (Table I). This was not significantly different from air breathing control rats but significantly different from N $_2$ O-treated animals not given methionine ($P = <0.001$).

Effect of SAM. When H $_4$ PteGlu was given to N $_2$ O-treated rats with 16 μmol SAM the mean amount of folate polyglutamate formed in three animals was $31.0 \pm 0.7\%$, significantly $<48.0 \pm 3.1\%$ formed in the presence of an equimolar dose of methionine




FIGURE 1 [^{14}C]H $_4$ PteGlu (0.02 μmol) was injected intraperitoneally to groups of three rats together with increasing doses of methionine. The animals were maintained under N $_2$ O/O $_2$ (1:1) and hepatic uptake of labeled folate (○) and the proportion converted into folate polyglutamate (●) determined.

TABLE I
 Synthesis of Folate Polyglutamate from [¹⁴C]H₄PteGlu
 (0.02 μmol). Effect of N₂O and Potential
 "Single Carbon" Donors (16 μmol)

Additive	Hepatic folate polyglutamate (percentage of labeled hepatic folate)	
	Air	N ₂ O
Nil	57.5±1.4 (15)	—
Nil	—	5.3±1.3 (32)
Methionine	70, 67	48.0±3.1 (3)
SAM	64, 62	31.0±0.7 (3)
Serine	ND	0.0 (3)
Betaine	66, 70	9.0±2.2 (3)
Sarcosine	57, 68	7.0±2.1 (3)
Choline	65, ND	15.0±1.4 (3)
Homocysteine	60	0.0 (3)
Glycine	61	0.0 (3)
Adenosine	65	0.0 (3)
5'MTA	68, 51	68.8±3.7 (8)

Data represent mean±SEM (number of rats). ND, not determined.

(*P* = <0.002). Increasing the dose of SAM to 32 μmol increased the amount of folate polyglutamate formed to the same level obtained with 16 μmol methionine. (Fig. 4).

Effect of other compounds. Serine, glycine, ho-

mocysteine, and adenosine had no effect on folate polyglutamate synthesis in the N₂O-treated rat. Choline and betaine produced 15 and 9% conversion of hepatic labeled folate into the polyglutamate form. Sarcosine, which yields formate directly by oxidation of its methyl group (26), yielded 7.0% folate polyglutamate formation.

Effect of 5'MTA. MTA was the most effective compound tested in overcoming the effect of N₂O in blocking folate polyglutamate synthesis (Fig. 5). The mean conversion in eight animals was 68.8±3.7% of the labeled folate in liver (Fig. 2, Table I). This value was significantly greater than the amount of folate polyglutamate formed in the presence of methionine or SAM (*P* = <0.001).

Effect of addition in control animals. Each addition was also given to 1-2 air breathing animals. There was no effect on folate polyglutamate formation, the result being the same as that obtained when H₄PteGlu was given alone.

DISCUSSION

Pteroylpolyglutamates have been the subject of two recent reviews (27, 28) but neither discuss the relation to cobalamin. In untreated pernicious anemia the fall in erythrocyte folate is due almost entirely to a fall in




FIGURE 2 The percentage of [¹⁴C]H₄PteGlu in liver converted into folate polyglutamate in N₂O-treated rats after an 0.02 μmol dose accompanied by 16 μmol of various compounds (●). Data in air-breathing rats (○) and in rats given only H₄PteGlu (second column) are also shown.




FIGURE 3 [^{14}C]H₄PteGlu (0.02 μmol) was given to two rats inhaling N₂O/O₂ (1:1) alone (O) and with 16 μmol methionine (●). The livers were removed and labeled folate analogues separated by chromatography on DE52 cellulose. The radioactivity in the eluate fractions are shown. The liver from the rat given H₄PteGlu alone failed to form any analogues containing longer glutamic acid chain folates (Glu 4-6) whereas the animal receiving methionine in addition formed folate polyglutamates in normal amount.

the predominant folate polyglutamate component (18, 19).

Impairment of folate polyglutamate synthesis was demonstrated in N₂O-treated mice (15) using PteGlu as substrate and in rats using H₄PteGlu and 5-CH₃-

H₄PteGlu as substrates (22). As a normal proportion of hepatic folate is converted into polyglutamate in the N₂O-treated rat when CHO-H₄PteGlu is given, it appears that there is no impairment in the enzyme pteroylpolyglutamic acid synthetase (22) but rather a failure to provide the appropriate folate substrate. The N₂O data suggest that formyltetrahydrofolates are the required substrates. Other studies of substrate requirements for folate polyglutamate synthesis in normal rat liver from air-breathing animals noted that H₄PteGlu was twice as active as 10-CHO-H₄PteGlu (29).

Further support for the suggestion that formylation of H₄PteGlu was impaired in the N₂O-treated rat arose from the increase in the activity of the enzyme formyltetrahydrofolate synthetase, which we interpreted as a response to lack of 'active formate'. Since the primary effect of N₂O is inactivation of methionine synthesis we looked for evidence for methionine being the source of active formate.

The observations reported in this paper suggest that the pathway from methionine to formate may be via SAM, decarboxylated SAM, 5'MTA, 5'methylthioribose to formate (Fig. 6). The last step in this cycle, namely metabolism of 1-phospho-5-methylthioribose to formate was demonstrated by Trackman and Abeles (30) in rat liver. 5'MTA was the most active of all the compounds we tested in overcoming the N₂O-induced impairment of folate polyglutamate synthesis. The lower activity of SAM as compared with methionine in restoring polyglutamate synthesis is probably related to the much lower hepatic uptake of SAM as opposed to methionine (31).




FIGURE 4 [^{14}C]H₄PteGlu (0.02 μmol) was given to rats inhaling N₂O/O₂ (1:1) with increasing doses of L-methionine (●) or SAM (○). The amount of folate polyglutamate formed from the labeled H₄PteGlu in liver was measured and expressed as a percentage of total uptake of substrate. Maximum values were reached with ~ 16 μmol methionine and 30 μmol SAM.




FIGURE 5 [^{14}C] H_4PteGlu ($0.02 \mu\text{mol}$) was given to two rats inhaling $\text{N}_2\text{O}/\text{O}_2$ (1:1) alone (○) and with $16 \mu\text{mol}$ 5'MTA (●). The livers were removed and labeled folate analogues separated by chromatography on DE52 cellulose. The radioactivity in the eluate fractions is shown. The liver from the rat given H_4PteGlu alone failed to form any analogues containing longer glutamic acid chain folates (Glu 4-6) whereas the rat given MTA converted 70% of substrate into folate polyglutamate.

Failure to form MTA results in a failure of cell division (32). The pathways shown in Fig. 6 leading to MTA and polyamine production are greatly accelerated by increased cell division (33). Limiting the amount of methionine or homocysteine causes a two-fold increase in SAM decarboxylase activity (34). Thus, these pathways seem to be essential to cell division and the reason may be the requirement for formate, which is an important single carbon unit donor.

Further evidence for a role of cobalamin in metabolism of MTA was reported by Sugimoto and Fukui (35). In studying the cobalamin requirements of *Ochromonas malhamensis* they found that methionine partially replaced the requirement for vitamin B_{12} , and the latter was replaced completely by a factor present in yeast, identified as 5'MTA.

The second aspect of N_2O toxicity brought out in these studies is impaired hepatic uptake (or retention) of folate analogues. This too is well documented in untreated pernicious anemia (36, 37). It is only poorly corrected by methionine in vivo (21, 38). It has been suggested that the reduced amount of labeled folate in liver of N_2O -treated animals is due to increased loss rather than impaired uptake (39). The amount of labeled folate in the liver of N_2O -treated animals is 20% of that in control animals at 24 h, whereas the fall in total folate was from a mean of 5.6 to $4.2 \mu\text{g/g}$ over the same period (10). An alternative hypothesis is that cobalamin is required for normal folate uptake by an action on the cell membrane and that this is impaired in the N_2O -treated rat.

The failure of the cobalamin-deficient or the N_2O -treated animal to utilize either PteGlu or H_4PteGlu normally (15, 20, 22, 40, 41) and the impaired utili-




FIGURE 6 Proposed pathways for formate formation from SAM.

zation of H₄PteGlu in untreated pernicious anemia (42, 43, 44) are not readily explained on the basis of the methylfolate trap hypothesis (45, 46), which requires that forms of folate other than 5-CH₃-H₄PteGlu be used in a normal manner in cobalamin deficiency. We have not been able to demonstrate significant accumulation of folates in the methyl form in the N₂O-treated animal (10, 11), that is, we have not been able to demonstrate methyl folate trapping.

A third possibility is that MTA is influencing folate function by its resynthesis into methionine, a pathway demonstrated in rat liver by Backlund and Smith (42). However, if MTA acts merely through its resynthesis to methionine it would be difficult to explain why it is significantly more effective than methionine in overcoming impaired folate polyglutamate synthesis, particularly as both MTA and methionine are taken up by liver at similar rates (31).

Methionine may act, too, either through SAM via its role in transmethylation, or as a source of formate through oxidation of the methyl group. Impaired transmethylation via SAM has been proposed as an explanation for the N₂O effects (48) and interference with transmethylation by cycloleucine produces changes in the central nervous system not dissimilar to those occurring in cobalamin neuropathy. The biochemical lesion has been postulated as impaired methylation of arginine in myelin basic protein (49). This has not been demonstrated, however, in the neuropathy occurring in some species on prolonged exposure to N₂O.

Choline, betaine, and sarcosine all show some activity in overcoming the defect studied in the N₂O-treated rat. No doubt this is due to their potential as single-carbon unit donors and their role in methyl group transfer. It is notable that serine and glycine, which are purported to be the principal source of single carbon units, are totally ineffective in this system, although the pathways by which they donate a carbon unit to the folate pool remains intact in the N₂O-treated rat. The carbon unit from serine is a methylene group whereas, we suggest, the more oxidized formate is required. The fall in the methenyltetrahydrofolate cyclohydrolase activity in response to N₂O may hamper oxidation of methylene units to formate.

A criticism that has been levelled at studies dependent on the use of H₄PteGlu is that this compound is unstable and hence degraded. H₄PteGlu is stable if stringent reducing conditions are maintained and this has been the practice in our studies. Its integrity has been confirmed by the absorption spectrum and by its availability for *P. cerevisiae* on microbiological assay.

A further difficulty with our hypothesis is that sodium formate given as a single 250- μ mol injection did not correct the impaired folate polyglutamate

synthesis in the N₂O-treated rat (unpublished observation). We continue to explore this. It may be that, like formaldehyde (50), "active-formate" is required under the particular circumstances present in the N₂O-treated rat.

REFERENCES

1. Banks, R. G. S., R. J. Henderson, and J. M. Pratt. 1968. Part III. Some reactions of nitrous oxide with transition-metal complexes. *J. Chem. Soc. (A)*: 2886-2889.
2. Lassen, H. C. A., E. Henriksen, F. Neukirch, and A. S. Kristensen. 1956. Treatment of tetanus. Severe bone marrow depression after prolonged nitrous-oxide anaesthesia. *Lancet* I: 527-530.
3. Amess, J. A. L., J. F. Burman, G. M. Rees, D. G. Nancekievill, and D. L. Mollin. 1978. Megaloblastic haemopoiesis in patients receiving nitrous oxide. *Lancet*. II: 339-342.
4. Skacel, P. O., A. M. Hewlett, J. D. Lewis, M. Lumb, J. F. Nunn, and I. Chanarin. 1982. Studies on the haemopoietic toxicity of nitrous oxide in man. *Br. J. Haematol.* In press.
5. Wilson, P., F. I. R. Martin, and P. M. Last. 1956. Bone-marrow depression in tetanus. Report of a fatal case. *Lancet*. II: 442-443.
6. Layzer, R. B. 1978. Myeloneuropathy after prolonged exposure to nitrous oxide. *Lancet*. II: 1227-1230.
7. Deacon, R., M. Lumb, J. Perry, I. Chanarin, B. Minty, M. J. Halsey, and J. F. Nunn. 1978. Selective inactivation of vitamin B₁₂ in rats by nitrous oxide. *Lancet*. II: 1023-1024.
8. Kondo, H., M. L. Osborne, J. F. Kolhouse, M. J. Binder, E. R. Podell, C. S. Utley, R. S. Abrams, and R. H. Allen. 1981. Nitrous oxide has multiple deleterious effects on cobalamin metabolism and causes decreases in activities of both mammalian cobalamin-dependent enzymes in rats. *J. Clin. Invest.* 67: 1270-1283.
9. Lumb, M., J. Perry, R. Deacon, and I. Chanarin. 1981. Changes in plasma folate levels in rats inhaling nitrous oxide. *Scand. J. Haematol.* 26: 61-64.
10. Lumb, M., R. Deacon, J. Perry, I. Chanarin, B. Minty, M. J. Halsey, and J. F. Nunn. 1980. The effect of nitrous oxide inactivation of vitamin B₁₂ on rat hepatic folate. Implications for the methylfolate-trap hypothesis. *Biochem. J.* 186: 933-936.
11. Lumb, M., J. Perry, R. Deacon, and I. Chanarin. 1981. Changes in tissue folates accompanying nitrous oxide-induced inactivation of vitamin B₁₂ in the rat. *Am. J. Clin. Nutr.* 34: 2412-2417.
12. Lumb, M., J. Perry, R. Deacon, and I. Chanarin. 1982. Urinary folate loss following inactivation of vitamin B₁₂ by nitrous oxide in rats. *Br. J. Haematol.* 51: 235-242.
13. Lumb, M., J. Perry, R. Deacon, and I. Chanarin. 1981. Recovery of tissue folates after inactivation of cobalamin by nitrous oxide. The significance of dietary folate. *Am. J. Clin. Nutr.* 34: 2418-2422.
14. Perry, J., R. Deacon, M. Lumb, and I. Chanarin. 1980. The effect of nitrous oxide-induced inactivation of vitamin B₁₂ on the activity of formyl-methenyl-methylenetetrahydrofolate synthetase transferase. *Biochem. Biophys. Res. Commun.* 97: 1329-1333.
15. Scott, J. M., B. Reed, B. McKenna, P. McGing, S. McCann, H. O'Sullivan, P. Wilson, and D. G. Weir. 1979. A study of the multiple changes induced *in vivo* in experimental animals by inactivation of vitamin B₁₂

- using nitrous oxide. In *Chemistry and Biology of Pteridines*. R. L. Kisliuk and G. M. Brown, editors. Elsevier/North Holland, New York. 335-340.
16. Deacon, R., I. Chanarin, J. Perry, and M. Lumb. 1980. Impaired deoxyuridine utilization in the B₁₂-inactivated rat and its correction by folate analogues. *Biochem. Biophys. Res. Commun.* **93**: 516-520.
 17. Deacon, R., J. Perry, M. Lumb, and I. Chanarin. 1981. The effect of nitrous oxide-induced inactivation of vitamin B₁₂ on thymidylate synthetase activity of rat bone marrow cells. *Biochem. Biophys. Res. Commun.* **102**: 215-218.
 18. Jeejeebhoy, K. N., S. M. Pathare, and J. M. Noronha. 1965. Observations on conjugated and unconjugated blood folate levels in megaloblastic anemia and the effects of vitamin B₁₂. *Blood*. **26**: 354-359.
 19. Chanarin, I., J. Perry, and M. Lumb. 1974. The biochemical lesion in vitamin B₁₂ deficiency in man. *Lancet*. **I**: 1251-1252.
 20. Gawthorne, J. M., and R. M. Smith. 1973. The synthesis of pteroylpolylglutamates by sheep liver enzymes *in vitro*. *Biochem. J.* **136**: 295-301.
 21. Gawthorne, J. M., and E. L. R. Stokstad. 1971. The effect of vitamin B₁₂ and methionine on folic acid uptake by rat liver. *Proc. Soc. Exp. Biol. Med.* **136**: 42-46.
 22. Perry, J., I. Chanarin, R. Deacon, and M. Lumb. 1979. The substrate for folate polyglutamate biosynthesis in the vitamin B₁₂-inactivated rat. *Biochem. Biophys. Res. Commun.* **29**: 678-684.
 23. Silverman, M., and J. M. Noronha. 1961. A method for the preparation of tetrahydrofolic acid. *Biochem. Biophys. Res. Commun.* **4**: 180-182.
 24. Nixon, P. J., and J. R. Bertino. 1971. Enzymic preparations of radiolabeled (+)-L-5-methyltetrahydrofolate and (+)-L-5-formyltetrahydrofolate. *Anal. Biochem.* **43**: 162-172.
 25. Buehring, K. U., Tamura, and E. L. R. Stokstad. 1974. Folate coenzymes of *Lactobacillus casei* and *Streptococcus faecalis*. *J. Biol. Chem.* **249**: 1081-1089.
 26. Frisell, W. R., and W. C. Sorrell. 1967. Oxidative phosphorylation in the metabolism of sarcosine and formaldehyde. *Biochim. Biophys. Acta.* **131**: 207-210.
 27. Kisliuk, R. 1981. Pteroylpolylglutamates. *Mol. Cell. Biochem.* **39**: 331-345.
 28. McGuire, J. J., and J. R. Bertino. 1981. Enzymatic synthesis and function of folylpolyglutamates. *Mol. Cell. Biochem.* **38**: 19-48.
 29. McGuire, J. J., P. Hsieh, J. K. Coward, and J. R. Bertino. 1980. Enzymatic synthesis of folylpolyglutamates. Characterization of the reaction and its products. *J. Biol. Chem.* **255**: 5776-5788.
 30. Trackman, P. C., and R. H. Abeles. 1981. The metabolism of 1-phospho-5-methylribose. *Biochem. Biophys. Res. Commun.* **103**: 1238-1244.
 31. Zappia, V., P. Galletti, and M. Porcelli. 1978. Uptake of adenosylmethionine and related sulfur compounds by isolated rat liver. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* **90**: 331-335.
 32. Toohey, J. I. 1978. Methylthioadenosine nucleoside phosphorylase deficiency in methylthio-dependent cancer cells. *Biochem. Biophys. Res. Commun.* **83**: 27-35.
 33. Tabor, C. W., and H. Tabor. 1976. 1,4-Diaminobutane (putrescine), spermidine, and spermine. *Annu. Rev. Biochem.* **45**: 225-306.
 34. Tisdale, M. J. 1981. Effect of methionine deprivation on S-adenosylmethionine decarboxylase of tumor cells. *Biochim. Biophys. Acta.* **675**: 366-372.
 35. Sugimoto, Y., and S. Fukui. 1976. Studies on biological functions of 5'-methylthioadenosine vitamin B₁₂-replacing effect for *Ochromonas malhamensis*. *J. Nutr. Sci. Vitaminol.* **22**: 89-100.
 36. Das, K. C., and A. V. Hoffbrand. 1970. Studies of folate uptake by phytohaemagglutinin-stimulated lymphocytes. *Br. J. Haematol.* **19**: 203-221.
 37. Tisman, G., and V. Herbert. 1973. B₁₂ dependence of cell uptake of serum folate: an explanation for high serum folate and cell folate depletion in B₁₂ deficiency. *Blood*. **41**: 465-469.
 38. McGing, P. G., and J. M. Scott. 1980. The role of methionine and vitamin B₁₂ in folate incorporation by rat liver. *Br. J. Nutr.* **43**: 235-237.
 39. McGing, P. G., and J. M. Scott. 1981. Evidence that the decreased liver folate status following vitamin B₁₂ inactivation in the mouse is due to increased loss rather than impaired uptake. *Biochim. Biophys. Acta.* **673**: 594-597.
 40. Smith, R. M., and W. S. Osborne-White. 1973. Folic acid metabolism in vitamin B₁₂-deficient sheep. Depletion of liver folates. *Biochem. J.* **136**: 279-293.
 41. Shane, B., J. E. Watson, and E. L. R. Stokstad. 1977. Uptake and metabolism of [³H]folate by normal and vitamin B₁₂- and methionine-deficient rats. *Biochim. Biophys. Acta.* **497**: 241-252.
 42. Deacon R., I. Chanarin, J. Perry, and M. Lumb. 1980. Marrow cell from patients with untreated pernicious anaemia cannot use tetrahydrofolate normally. *Br. J. Haematol.* **46**: 523-528.
 43. Deacon, R., I. Chanarin, J. Perry, and M. Lumb. 1982. A comparison of tetrahydrofolate and 5-formyltetrahydrofolate in correcting the impairment of thymidine synthesis in pernicious anaemia. *Scand. J. Haematol.* **28**: 289-292.
 44. Taheri, M. R., R. G. Wickremasinghe, B. J. Jackson, and A. V. Hoffbrand. 1982. The effect of folate analogues and vitamin B₁₂ on provision of thymine nucleotides for DNA synthesis in megaloblastic anemia. *Blood*. **59**: 634-640.
 45. Herbert, V., and R. Zalusky. 1962. Interrelation of vitamin B₁₂ and folic acid metabolism: folic acid clearance studies. *J. Clin. Invest.* **41**: 1263-1276.
 46. Noronha, J. M., and M. Silverman. 1962. On folic acid, vitamin B₁₂, methionine, and formiminoglutamic acid metabolism. In *Vitamin B₁₂ and Intrinsic factor*. H. C. Heinrich editor. Enke, Stuttgart. 728-736.
 47. Backlund, P. S., Jr., and R. A. Smith. 1981. Methionine synthesis from 5'-methylthioadenosine in rat liver. *J. Biol. Chem.* **256**: 1533-1535.
 48. Scott, J. M., J. J. Dinn, P. Wilson, and D. G. Weir. 1981. Pathogenesis of subacute combined degeneration: a result of methyl group deficiency. *Lancet*. **II**: 334-337.
 49. Crang, A. J. and W. Jacobson. 1980. The methylation *in vitro* of myelin basic protein by arginine methylase from mouse spinal cord. *Biochem. Soc. Trans.* **8**: 619-620.
 50. Frisell, W. R., and V. M. Randolph. 1973. Effect of adenine nucleotides on the oxidation of N-methyl groups in liver mitochondria. *Biochim. Biophys. Acta.* **292**: 360-365.
 51. Leslie, G. B., and C. M. Baugh. 1974. The uptake of pteroyl[¹⁴C]glutamic acid into rat liver and its incorporation into the natural pteroyl-γ-glutamates of that organ. *Biochemistry*. **13**: 4957-4964.