

Supplemental Figure 1. EC-Del1 mice have low inflammatory profile in aorta. mRNA levels of aortic fibrotic (A-C), pro- (E-M) and anti-inflammatory (N) markers. mRNA levels of Del-1 (O) (n=6-8 mice per group for all experiments). Data represent mean \pm SEM. * P <0.05, ** P <0.01 analyzed by two-way ANOVA with Bonferroni post hoc test to adjust for multiple comparisons.

Supplemental Figure 2. EC-Del1 mice have low inflammatory profile in heart. mRNA levels of cardiac fibrotic (A-B), hypertrophic (C-F), pro- (G-M) and anti-inflammatory (N) markers. mRNA levels of Del-1 (O) (n=6-8 mice per group for all experiments). Data represent mean \pm SEM. * P <0.05, ** P <0.01 analyzed by two-way ANOVA with Bonferroni post hoc test to adjust for multiple comparisons.


Supplemental Figure 3. DEL-1-Fc treated mice have low inflammatory profile in aorta. mRNA levels of aortic fibrotic (A-C), pro- (D-K) and anti-inflammatory (L) markers (n=6 mice per group). Data represent mean \pm SEM. * P <0.05, ** P <0.01 analyzed by one-way ANOVA with Bonferroni post hoc test to adjust for multiple comparisons.

Supplemental Figure 4. DEL-1-Fc treated mice have low inflammatory profile in aorta. mRNA or protein levels of cardiac fibrotic (A-B), hypertrophic (C-F), pro- (G-K) and anti-inflammatory (L) markers (n=6 mice per group). Data represent mean \pm SEM. * P <0.05, ** P <0.01 analyzed by one-way ANOVA with Bonferroni post hoc test to adjust for multiple comparisons.


Supplemental Figure 5. DEL-1-Fc, but not DEL-1-RGE-Fc treated mice have low inflammatory profile in aorta. mRNA or protein levels of aortic fibrotic (A-C), pro- (D-K) and anti-inflammatory (L) markers (n=6 mice per group). Data are means \pm SE. * P <0.05, ** P <0.01 analyzed by one-way ANOVA with Bonferroni post hoc test to adjust for multiple comparisons.

Supplemental Figure 6. DEL-1-Fc, but not DEL-1-RGE-Fc treated mice have low inflammatory profile in heart. mRNA or protein levels of cardiac fibrotic (A-B), hypertrophic (C-F), pro- (G-K) and anti-inflammatory (L) markers (n=6 mice per group). Data are means \pm SE. * P <0.05, ** P <0.01 analyzed by one-way ANOVA with Bonferroni post hoc test to adjust for multiple comparisons.


Supplemental Figure 1




Supplemental Figure 2




Supplemental Figure 3



Supplemental Figure 4



Supplemental Figure 5



Supplemental Figure 6

