

Bacterial vaginosis associates with dysfunctional T cells and altered soluble immune factors in the cervicovaginal tract

Finn MacLean, ... , Jairam R. Lingappa, Jennifer M. Lund

J Clin Invest. 2025. <https://doi.org/10.1172/JCI184609>.

Clinical Research and Public Health

In-Press Preview

AIDS/HIV

Immunology

Infectious disease

Background: Bacterial vaginosis (BV) is a dysbiosis of the vaginal microbiome that is prevalent among reproductive-age females worldwide. Adverse health outcomes associated with BV include an increased risk of sexually-acquired HIV, yet the immunological mechanisms underlying this association are not well understood. **Methods:** To investigate BV-driven changes to cervicovaginal tract (CVT) and circulating T cell phenotypes, Kinga Study participants with or without BV provided vaginal tract (VT) and ectocervical (CX) tissue biopsies and PBMC samples. **Results:** High-parameter flow cytometry revealed an increased frequency of cervical conventional CD4⁺ T cells (Tconv) expressing CCR5. However, we found no difference in number of CD3⁺CD4⁺CCR5⁺ cells in the CX or VT of BV⁺ versus BV⁻ individuals, suggesting that BV-driven increased HIV susceptibility may not be solely attributed to increased CVT HIV target cell abundance. Flow cytometry also revealed that individuals with BV have an increased frequency of dysfunctional CX and VT CD39⁺ Tconv and CX tissue-resident CD69⁺CD103⁺ Tconv, reported to be implicated in HIV acquisition risk and replication. Many soluble immune factor differences in the CVT further support that BV elicits diverse and complex CVT immune alterations. **Conclusion:** Our comprehensive analysis expands on potential immunological mechanisms that may underlie the adverse health outcomes associated with BV including increased HIV susceptibility.

Find the latest version:

<https://jci.me/184609/pdf>



**Bacterial vaginosis associates with dysfunctional T cells and altered soluble immune factors
in the cervicovaginal tract**

Finn MacLean¹, Adino Tesfahun Tsegaye², Jessica B. Graham¹, Jessica L. Swarts¹, Sarah C. Vick¹, Nicole Potchen¹, Irene Cruz Talavera¹, Lakshmi Warriar¹, Julien Dubrulle³, Lena K. Schroeder³, Ayumi Saito², Corinne Mar², Katherine K. Thomas², Matthias Mack⁴, Michelle C. Sabo⁵, Bhavna H. Chohan^{2,6}, Kenneth Ngunjiri^{2,7}, Nelly Mugo^{2,8}, Jairam R. Lingappa^{2,5,9*}, Jennifer M. Lund^{1,2*} for the Kinga Study Team

¹Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center, Seattle, USA

²Department of Global Health, University of Washington, Seattle, USA

³Cellular Imaging Shared Resource, Fred Hutchinson Cancer Research Center, Seattle, USA

⁴Department of Internal Medicine-Nephrology, University Hospital Regensburg, Regensburg, Germany

⁵Department of Medicine, University of Washington, Seattle, USA

⁶Center for Virus Research, Kenya Medical Research Institute, Nairobi, Kenya

⁷School of Public Health, Jomo Kenyatta University of Agriculture and Technology, Nairobi, Kenya

⁸Center for Clinical Research, Kenya Medical Research Institute, Nairobi, Kenya

⁹Department of Pediatrics, University of Washington, Seattle, USA

*Contributed equally

Corresponding Authors:

Jennifer M. Lund

1100 Fairview Ave N., E5-110

Seattle, WA 98109

+1-206-667-2217

jlund@fredhutch.org

Jairam R. Lingappa

908 Jefferson St, Box 359927

Seattle, WA 98104

+1-206-520-3822

lingappa@uw.edu

The authors have declared that no conflict of interest exists.

Abstract

Background: Bacterial vaginosis (BV) is a dysbiosis of the vaginal microbiome that is prevalent among reproductive-age females worldwide. Adverse health outcomes associated with BV include an increased risk of sexually-acquired HIV, yet the immunological mechanisms underlying this association are not well understood.

Methods: To investigate BV-driven changes to cervicovaginal tract (CVT) and circulating T cell phenotypes, Kinga Study participants with or without BV provided vaginal tract (VT) and ectocervical (CX) tissue biopsies and PBMC samples.

Results: High-parameter flow cytometry revealed an increased frequency of cervical conventional CD4⁺ T cells (Tconv) expressing CCR5. However, we found no difference in number of CD3⁺CD4⁺CCR5⁺ cells in the CX or VT of BV⁺ versus BV⁻ individuals, suggesting that BV-driven increased HIV susceptibility may not be solely attributed to increased CVT HIV target cell abundance. Flow cytometry also revealed that individuals with BV have an increased frequency of dysfunctional CX and VT CD39⁺ Tconv and CX tissue-resident CD69⁺CD103⁺ Tconv, reported to be implicated in HIV acquisition risk and replication. Many soluble immune factor differences in the CVT further support that BV elicits diverse and complex CVT immune alterations.

Conclusion: Our comprehensive analysis expands on potential immunological mechanisms that may underlie the adverse health outcomes associated with BV including increased HIV susceptibility.

Trial Registration: Clinicaltrials.gov ID# NCT03701802

Funding: This work was supported by the following grants from the National Institutes of Health: R01 AI131914 (to JML and JRL), R01 AI141435 (to JML), and R01 AI129715 (to JRL).

Introduction

Bacterial vaginosis (BV) is characterized by a vaginal dysbiosis in which the normally *Lactobacillus*-dominated microbiome is replaced by one with higher bacterial diversity, including increased concentrations of anaerobic bacteria (1). BV results in symptoms of vaginal inflammation, discharge, and discomfort, and is estimated to affect 23-29% of reproductive-age females worldwide (2), making BV the most common cause of vaginal symptoms leading patients to seek medical care (3). BV is also associated with several adverse medical outcomes (1, 2, 4-11) including preterm delivery (12), chorioamnionitis (13), endometritis (14), and an increased risk of HIV acquisition (5, 11). The issue of HIV acquisition risk is of particular concern in parts of the world where both BV and HIV are prevalent, such as in sub-Saharan Africa (15). Although many studies have investigated immune system alterations associated with BV (1, 5-7, 10, 11, 16-19), the link between cervicovaginal tract (CVT) T cells, which may mediate the relationships between BV and adverse health outcomes, is not well understood. Here, we sought to better understand the impact of BV on CVT mucosal and systemic immune responses in Kenyans that may alter HIV susceptibility.

In the CVT, innate and adaptive immune cells line the epithelium and lamina propria to provide immunological surveillance (20). Notably, T cell populations in the CVT maintain distinct phenotypes from those circulating in the blood (21, 22). These include tissue-resident memory T cells (T_{RM}) bearing CD69 with or without CD103 (23). CD69 is a marker of activation (24) that has a unique role in maintaining tissue residency by binding S1PR1 and preventing egress from the tissue (25-27). CD103 further promotes tissue residency by binding to E-cadherin (28). T_{RM} are retained in the mucosal tissue and have the ability to readily respond to pathogens *in situ* (21, 29-31). Upon recognition of their cognate antigen, T_{RM} use their “sensing

and alarm function” to produce pro-inflammatory cytokines and chemokines that rapidly recruit and activate other immune cells, thereby potentiating the tissue immune response while also executing their specific effector and cytotoxic functions (32-35).

Given that in the context of heterosexual transmission, genital mucosal tissue is where host exposure to HIV most commonly first occurs, activated CD4⁺ T cells expressing the HIV coreceptor CCR5 in the CVT have been thought to be prime targets for initial HIV infection (36). Studies using a mouse model demonstrated that the introduction of BV-associated bacteria into the vagina led to an increased quantity of CVT CD4⁺ T cells expressing the activation marker CD44 and HIV coreceptor CCR5 (7), suggesting dysbiosis of normal vaginal flora may lead to an increase of HIV target cells. In humans, studies of immune responses in the context of BV have characterized immune cells collected from the CVT lumen via minimally invasive sampling techniques such as cervicovaginal lavage, vaginal swabs, cervical swabs, or cytobrushes. These studies have identified an increase in CCR5⁺CD4⁺ T cells in the CVT lumen of BV+ individuals (37-40), without addressing whether this observation is true in the CVT deeper tissue layers. In addition to the detection of HIV target cells in the CVT lumen of individuals with BV, studies have shown a connection between BV and the presence of pro-inflammatory cytokines in the CVT, most reliably IL-1 β (7, 37, 41, 42), adding further support to the idea that proinflammatory mechanisms underlie the link between BV and increased HIV susceptibility. However, the relationship between BV and an increase in CD4⁺ T cell activation is inconsistently observed (43, 44), suggesting more complex mechanisms may be contributing to increased HIV susceptibility in individuals with BV. Further characterizing BV-driven immune modulations in the unique immune environment of the CVT is paramount to understanding mechanisms underlying increased HIV susceptibility in individuals with BV.

To comprehensively evaluate immune alterations associated with BV, we used a recent advance in tissue cryopreservation (45) combined with high-parameter, high-throughput flow cytometry to compare immune cells isolated from CVT tissue biopsies and blood from individuals with versus without BV. Given the dense T cell populations within CVT tissue (46), cells isolated from CVT mucosal biopsies may better represent the mucosal tissue immune environment than those isolated from the CVT lumen by cytobrush or other methods. Additionally, improved flow cytometry technology allowed us to broadly evaluate immune cells in the context of BV for phenotypic markers not previously characterized.

Here, we report detailed T cell characteristics of vaginal tract (VT) and ectocervix (CX) mucosal tissues and PBMC samples from the same participants, comparing individuals with to those without BV. Additionally, we analyzed CVT fluid and serum for local and systemic cytokines and chemokines in individuals with versus without BV. While we did not observe an overall increase in $CD3^+CD4^+CCR5^+$ HIV target cells in the CVT of BV+ individuals, we did observe CX and VT T cells with dysfunctional phenotypes and altered expression of soluble mediators in CVT fluid from BV+ individuals. Our findings highlight different mechanisms that may underlie increased HIV susceptibility associated with BV.

Results

Study population characteristics

204 participants met the criteria to be included in the BV flow cytometry and/or soluble immune factor analysis (Table 1). 46 participants contributed at least one sample when they were BV+ by a Nugent score of 7-10 (34 at study enrollment and 12 at study exit). The BV- comparator group comprised N=158 participants with normal flora (Nugent score 0-3) who provided at least one sample at enrollment. Overall, our study population was young (55.9% were <30 years of age), sexually active (median of 8 unprotected sex acts in the prior month), with 40.8% using hormonal contraceptives, 39.9% Herpes Simplex Virus 2 (HSV-2) seropositive, and 17.6% had a sexual partner living with HIV (Table 1). 187 participants provided a CX sample (BV- N=149, BV+ N=38), 190 participants provided a VT sample (BV- N=149, BV+ N=41), and 185 participants provided a PBMC sample (BV- N=143, BV+ N=42) for flow cytometry analysis. 193 participants provided a serum sample (BV- N=149, and BV+ N=44) and 174 participants provided a CVT fluid sample (BV- N= 136, BV+ N=38) for soluble immune factor analysis. A subset of individuals who were HSV-2 seronegative and HIV unexposed individuals provided a CX and/or VT tissue sample for immunofluorescent imaging at their enrollment visit (CX BV- N=44, BV+ N=11; VT BV- N=55, BV+ N=12).

Bacterial vaginosis did not alter the balance of T cell frequency in the cervicovaginal tract tissues

To investigate the CVT mucosal tissue and circulating T cell population in detail, we performed high-parameter flow cytometry on cells isolated from CX and VT tissue biopsies cryopreserved using a previously published method (45), as well as PBMC samples. PBMC

comparisons were *a priori* adjusted for hormonal contraceptive use, and CX and VT comparisons were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and semen exposure to reduce the effects of potential confounding variables on the analysis of BV-driven T cell alterations. We found that the majority of CD45⁺ cells in the VT and CX are CD3⁺ T cells (median > 75%; Figure 1A, Supplemental Figure 1, Supplemental Table 1), demonstrating the importance of T cells in immunosurveillance of the CVT. We assessed whether BV was associated with a shift in the proportion of CD25⁻CD127^{+/-} conventional CD4⁺ T cells (Tconv) as a fraction of CD45⁺ cells in all three specimen types and found no significant difference comparing BV- versus BV+ samples (Figure 1B). The proportion of CD8⁺ T cells was analyzed as a fraction of CD45⁺ cells (Figure 1C), and no significant differences were detected comparing each sample type from BV- versus BV+ individuals. The fraction of Tconv cells of total CD3⁺ T cells (Figure 1D) and CD8⁺ of total CD3⁺ T cells (Figure 1E) were also not significantly altered by BV for any of the three sample types analyzed, indicating that BV may not be associated with a shift in the major T cell subsets.

Conventional CD4⁺ T cells displayed increased markers of activation and tissue residency in the cervix of individuals with bacterial vaginosis

We hypothesized that BV may be associated with phenotypic alterations among T cell subsets to promote adverse health outcomes. Previous studies have limited the characterization of cervicovaginal T cells to markers of inflammation, including the HIV co-receptor CCR5. Herein, we include markers of activation previously described to be enriched on CD4⁺ T cells of the CVT in the context of persistent BV, including CCR5, HLA-DR, and CD38 (44). However, we additionally sought to broaden our characterization of T cell phenotypic alterations associated

with BV. We included Treg lineage markers (CD25, CD127, FoxP3) (47), a Th1 lineage marker (T-bet) (48), Th17 lineage markers (CD161, CCR6) (49, 50), markers of tissue residency (CD69 and CD103) (23), markers of inhibition (CD101 and CTLA-4) (51, 52), dysfunction (CD39) (53), progenitor potential (TCF-1) (54), exhaustion (PD-1) (55), an additional chemokine receptor used for trafficking and associated with activation (CXCR3) (56), cytotoxic function (Granzyme B) (57), and memory subsets (CCR7 and CD45RA) (58) to more broadly evaluate the phenotypic differences of CVT and circulating T cells in those with versus without BV.

We evaluated whether BV is associated with activated Tconv phenotypes that may contribute to increased HIV susceptibility through a previously hypothesized mechanism – CCR5 expression. We found that CX CD4⁺ Tconv cells displayed a higher frequency of the HIV coreceptor CCR5 in BV⁺ compared to BV⁻ individuals (median frequency 59% BV⁻ vs 75% BV⁺, $p_{\text{adj}} = 0.021$; Supplemental Table 1) as detected by flow cytometry and calculated as a fraction of total Tconv cells; minimal differences were observed when VT or PBMC samples from BV⁻ versus BV⁺ individuals were compared (Figure 2A). We also observed a significantly increased frequency of the activation marker HLA-DR on CX Tconv in those with compared to without BV (median frequency 22% BV⁻ vs 30% BV⁺, $p_{\text{adj}} = 0.012$; Supplemental Table 1), and minimal differences were observed comparing HLA-DR expression on VT and PBMC Tconv cells from BV⁻ versus BV⁺ individuals (Figure 2B). Together, these results demonstrate that BV does promote local proinflammatory Tconv activation and that Tconv activation is elevated in the CX mucosa but not in the VT or circulation.

We hypothesized that T_{RM} would play a critical role in immune modulations associated with BV, so we evaluated CD103⁺CD69⁺ co-expression on CD4⁺ Tconv T cells as canonical T_{RM} markers (24, 46). We found that the T_{RM} phenotype was more frequently observed on CX Tconv

cells in BV+ individuals (median frequency 14% BV- vs 35% BV+, $p_{\text{adj}} = 0.001$; Supplemental Table 1) and found no significant differences when comparing the frequency of CD103⁺CD69⁺ Tconv cells isolated from VT or PBMC samples from BV- versus BV+ individuals (Figure 2C). These results support that BV is associated with an increased proportion of tissue-resident Tconv in the CX but not in the VT. Not unexpectedly, CD4⁺ T_{RM} were not present in the circulation.

The total density of T cells and HIV target cells in the cervix and vagina was not altered by bacterial vaginosis

To assess whether BV is associated with an increased number or density of CCR5⁺ HIV target T cells in the CVT mucosa, we analyzed VT and CX tissue sections by immunofluorescent staining and microscopy. H&E staining was used to evaluate tissue quality and integrity and visualize the epithelium and lamina propria compartments (representative image Figure 3A). Immunofluorescent staining and cellular imaging were used to visualize CD3⁺, CD3⁺CD4⁺, and CD3⁺CD4⁺CCR5⁺ cells in each tissue section (representative image Figure 3B). In contrast to the increased frequency of CCR5⁺ Tconv that we observed as a fraction of total Tconv cells in the CX by flow cytometry (Figure 2A), we found no significant differences in the overall density of HIV target T cells in the CX (Figure 3C, median density 228.3 cells/mm² BV- vs 219.6 cells/mm² BV+, $p = 0.764$; Supplemental Table 2) or VT (Figure 3D, median density 227.0 cells/mm² BV- vs 336.2 cells/mm² BV+, $p = 0.787$; Supplemental Table 2). We also observed no difference when analyzing epithelium and lamina propria tissue compartments individually (Supplemental Figure 2, Supplemental Table 2), suggesting that BV may not significantly alter the overall abundance of HIV target cells within the deeper CX or VT mucosal tissue layers.

Conventional CD4⁺ T cells in the cervix showed signs of dysfunction in individuals with bacterial vaginosis

To more broadly characterize how BV impacts mucosal or systemic T cells that may elucidate alternative immunological mechanisms for adverse outcomes associated with BV, we evaluated Tconv cells for additional markers that could indicate altered functional capacity. We found CD39 expression significantly increased on Tconv cells in CX samples from BV⁺ versus BV⁻ individuals (median frequency 11% BV⁻ vs 23% BV⁺, $p_{\text{adj}} = 0.005$; Supplemental Table 1) and in VT samples from BV⁺ versus BV⁻ individuals (median frequency 15% BV⁻ vs 23% BV⁺, $p_{\text{adj}} = 0.039$; Supplemental Table 1) with minimal differences observed in PBMC samples (Figure 4A). CD39 is a marker of metabolic stress (53) that may contribute to Tconv dysfunction in the CX and VT of BV⁺ individuals. CD101 expression was also significantly increased on Tconv cells in CX samples from BV⁺ versus BV⁻ individuals (median frequency 19% BV⁻ vs 38% BV⁺, $p_{\text{adj}} = 0.002$; Supplemental Table 1), with minimal differences observed in VT and PBMC samples (Figure 4B). CD101 is another marker associated with tissue residency (23), though it has also been shown to contribute to an inhibitory phenotype (51) that may impede the proliferation of Tconv cells in BV⁺ individuals. Finally, we found a significant decrease in the frequency of TCF-1⁺ Tconv in the CX of BV⁺ compared to BV⁻ individuals (median frequency 44% BV⁻ vs 24% BV⁺, $p_{\text{adj}} = 0.008$; Supplemental Table 1), with no significant differences in the VT or PBMC (Figure 4C). TCF-1 is a stabilizing transcription factor implicated in the regulation of progenitor potential and promotion of T cell fate specification (54, 59-64). These results suggest that Tconv cells in the CX of BV⁺ individuals may have a reduced capacity to undergo differentiation or serve as proliferative progenitor cells, which, taken together, could suggest that these cells have reduced capacity to serve as potent effector T cells.

Th17 cells exhibited increased markers of activation and tissue residency in cervical samples from individuals with bacterial vaginosis

Lineage-specific Th17 Tconv (Th17) cells have been shown to play a role in immune responses to extracellular bacteria (65-67). To investigate the impact of BV on Th17 cells, we first analyzed the frequency of Th17 cells (defined as CD161⁺CCR6⁺ Tconv cells (49); Supplemental Figure 1) among total Tconv cells and found that the frequency of Th17 cells as a proportion of total Tconv cells was not significantly increased in any of the tissue types comparing BV⁺ and BV⁻ individuals (Figure 5A). However, the frequency of Th17 cells expressing the proinflammatory activation marker HLA-DR was significantly increased in CX during BV (median 40% BV⁻ vs 55% BV⁺, $p_{\text{adj}} = 0.024$, Supplemental Table 1) with a trend towards an increase in the VT during BV (median 38% BV⁻ vs 54% BV⁺, $p_{\text{adj}} = 0.067$, Supplemental Table 1) and minimal differences in PBMCs from BV⁻ versus BV⁺ individuals (Figure 5B).

We also measured the frequency of markers of tissue-residency (CD69⁺CD103⁺) on Th17 cells and found them significantly increased in the CX of BV⁺ compared to BV⁻ individuals (median 31% BV⁻ vs 54% BV⁺, $p_{\text{adj}} = 0.002$, Supplemental Table 1) (Figure 5C). In addition, the frequency of Th17 cells expressing CD101, consistent with a T_{RM} phenotype, was significantly increased in the CX of BV⁺ compared to BV⁻ individuals (median 39% BV⁻ vs 60% BV⁺, $p_{\text{adj}} = 0.006$, Supplemental Table 1) (Figure 5D).

Finally, we found that the frequency of Th17 cells expressing the activation marker and HIV co-receptor CCR5 was not significantly different between BV⁻ versus BV⁺ individuals, although notably, the median frequency of Th17 cells expressing CCR5 in BV⁺ individuals was

90% in CX and 93% in VT samples (Supplemental Table 1) (Figure 5E). Therefore, CVT Th17 cells predominantly express the HIV coreceptor CCR5, coinciding with previous reports that Th17 cells are viable targets for HIV infection (68). These results demonstrate that BV is associated with CX Th17 activation and that the increased tissue-resident or CD101-expressing CX Th17 cells may contribute to a mechanism for increased HIV susceptibility in individuals with BV.

Cervical CD8⁺ T cells displayed a dysfunctional phenotype in individuals with bacterial vaginosis

CD8⁺ T cells are thought to play an important role in pathogen clearance in the CVT, including against BV-associated organisms. To comprehensively evaluate how BV impacts CD8⁺ T cell phenotypes in the CVT mucosa and circulation, we analyzed CX, VT, and PBMC samples for the expression of effector markers (Supplemental Figure 1 and Supplemental Table 1). We found that the frequency of CD39⁺CD8⁺ T cells was significantly increased in CX samples from BV⁺ versus BV⁻ individuals (median frequency 7% BV⁻ vs 10% BV⁺, $p_{\text{adj}} = 0.023$; Supplemental Table 1), and moderately but insignificantly increased in the VT of BV⁺ individuals (median frequency 6% BV⁻ vs 10% BV⁺, $p_{\text{adj}} = 0.068$; Supplemental Table 1, Figure 6A). We also observed a moderate though insignificant decrease in Granzyme B expression, a marker of cytotoxicity, on CD8⁺ T cells in the CX of BV⁺ versus BV⁻ individuals (median frequency 25% BV⁻ vs 22% BV⁺, $p_{\text{adj}} = 0.094$; Supplemental Table 1) with marginal differences in the VT and PBMC samples from those with versus without BV (Figure 6B). Additionally, we observed a moderate but insignificant decrease in the frequency of T-bet⁺ CD8⁺ T cells in the CX of BV⁺ versus BV⁻ individuals (median frequency 18% BV⁻ vs 12% BV⁺, $p_{\text{adj}} = 0.064$;

Supplemental Table 1) with marginal differences in the VT and PBMC samples from those with versus without BV (Figure 6C). Altogether, these findings suggest a trend towards decreased CX CD8⁺ T cell activation in addition to an increased dysfunctional CD39⁺ CD8⁺ T cell phenotype observed in the CX of BV+ individuals.

Bacterial vaginosis was associated with reduced chemokine concentrations and increased inflammatory cytokine concentrations in cervicovaginal fluid

To investigate soluble immune factors in individuals with BV versus those without BV, we analyzed 71 cytokines and chemokines in serum and CVT fluid collected via menstrual cup (Softcup). In CVT fluid samples, 61 cytokines and chemokines met the criteria for continuous analysis (Figure 7A), while 10 cytokines and chemokines were analyzed as a dichotomous outcome (detected/not detected; Supplemental Figure 3A). In the serum, 52 different cytokines and chemokines met the criteria for continuous analysis (Figure 7B), 17 cytokines and chemokines were analyzed using the dichotomous outcome (Supplemental Figure 3B), and 2 cytokines were detectable in all individuals but had greater than 20% of samples out of range high so were excluded from the analysis (Supplemental Table 3). We observed significantly higher concentrations of 23 soluble immune factors including IL-1 α in CVT fluid collected from participants with versus without BV (Figure 7A). Additionally, we observed significantly lower concentrations of 12 soluble immune factors, including MIG/CXCL9 and IP-10 (CXCL10) (Figure 7A) in participants with BV versus those without BV. In our analysis of serum cytokines, fewer significant observations were observed than in the CVT fluid. In total, we observed a significant increase of 1 and a significant reduction of 5 quantifiable soluble immune factors in participants with versus without BV (Figure 7B). In the CVT fluid samples, of the 10 soluble

immune factors analyzed for their dichotomous outcome, 3 were detected significantly more frequently, and 1 cytokine was detected significantly less frequently in BV+ individuals (Supplemental Figure 3A). The dichotomous outcome analysis revealed an additional 2 other cytokines observed significantly less frequently in the serum of BV+ individuals (Supplemental Figure 3B). Overall, differentially expressed and detected soluble immune factors were primarily observed in CVT fluid samples, with fewer differences observed in serum samples, highlighting that BV has the greatest effects on the local CVT, as opposed to the circulating immune environment.

Discussion

In context of the global HIV pandemic, T cells serve as a double-edged sword: mediating anti-viral functions required to fight the virus while also acting as potential targets for HIV infection and replication (69, 70). The findings presented here add further detail that may underlie the effect of BV on HIV susceptibility by highlighting a more complex scenario that involves deleterious effects on T cell antiviral functions in addition to modulations in subsets that may act as potential HIV target cells.

Our flow cytometry analysis revealed an increase in the frequency of activation markers expressed on cervical Tconv, including a higher frequency of CCR5⁺ among total Tconv cells (Figure 2A), which supports prior studies that have identified increased levels of total CD3⁺CD4⁺CCR5⁺ T cells in the CVT lumen as a potential source of HIV target cells (7, 39, 40). However, our analysis of CD3⁺CD4⁺CCR5⁺ HIV target cell tissue density by immunofluorescent staining and cellular imaging of tissue sections indicated a relatively equivalent abundance of CD3⁺CD4⁺CCR5⁺ HIV target cells among BV- and BV+ participants in CX and VT tissue evaluated overall (Figure 3, C and D) and when the epithelium and lamina propria compartments were analyzed separately (Supplemental Figure 2). This supports the idea that BV does not have a profound effect on the overall abundance of HIV target cells in the cervicovaginal mucosa. Our conclusions are limited by the fact that the results are based in distinct technologies (flow cytometry versus microscopy), and that flow cytometry involves the evaluation of more samples and therefore has increased power to detect differences compared to microscopy. Nevertheless, our results underscore that the abundance of CD3⁺CD4⁺CCR5⁺ HIV target cells within the deeper CX and VT tissue layers in BV+ individuals is unlikely to solely account for increased

HIV susceptibility among those with BV, and this motivates the possibility that other factors contribute to this adverse outcome.

Our application of high-parameter, high-throughput flow cytometry on cells isolated from VT and CX tissue biopsies and PBMC samples documents that BV elicits complex changes to CVT T cells that may contribute to the inhibition of antiviral T cell function. We observed an increased frequency of CD4⁺ Tconv cells expressing CD39 in the CX and VT of BV⁺ individuals (Figure 4, A and B). The potential dysfunction of this T cell phenotype is suggested by prior studies showing CD39⁺ T cells have a decreased response to vaccines and an increased likelihood of undergoing apoptosis (53), raising the possibility that in the context of BV, cervical and vaginal CD4⁺ Tconv may exhibit increased intrinsic immunoregulation leading to reduced effector potential. This could lead to increased HIV susceptibility through an inability to mount an effective antiviral response upon exposure to the virus.

We also found that CD101⁺ Tconv cells, and specifically CD101⁺ Th17 cells, are more frequently observed in the CX of BV⁺ individuals (Figure 4B and 5D). Less is known about the function of CD101⁺ T cells in comparison to the other phenotypes examined in this study, but CD101⁺ T cells have been associated with decreased expansion after adoptive transfer in mice (51), suggesting CD101⁺ Tconv cells are inhibited and less likely to undergo proliferation. Additionally, CD101 is associated with exhausted T cell phenotypes that result from chronic activation and can promote immune dysfunction (71-73). Thus, the increased abundance of CD101⁺ Tconv and Th17 cells in the CX of BV⁺ individuals may also contribute to a dysfunctional Tconv immune response upon secondary exposure to HIV in the CX. Given the often recurrent and persistent nature of BV (44), chronic immune activation from repeated antigen exposure caused by BV-associated microbiota may lead to the progression of these

observed dysfunctional phenotypes that could promote HIV susceptibility by limiting antiviral T cell function, expansion, and differentiation.

We also observed a significant reduction in the proportion of TCF-1⁺ CD4⁺ Tconv in the CX of BV+ individuals (Figure 4C). TCF-1 is necessary for follicular T cell differentiation (59-61) and has been shown to play a role in Th17 responses by inhibiting IL-17 production (64). Fewer progenitor CD4⁺ T cells implies that more T cells are likely to be terminally differentiated in the CX, which may further limit lineage-specific differentiation upon secondary exposure to an infectious agent.

In addition to CD4⁺ T cells expressing CCR5, it has been shown that CD69⁺CD4⁺ T cells are associated with increased susceptibility to HIV infection in the CVT mucosal tissue (74). This suggests that the increased frequency of T_{RM} in the CX of BV+ individuals (Figure 2C) may mediate increased HIV acquisition in the context of BV. In particular, we found the CX had an increased frequency of Th17 T_{RM} (Figure 5C), which highly expresses CCR5 (Figure 5E). The possibility that these cells are a target for HIV acquisition in BV+ individuals is supported by a prior report that Th17 cells serve as primary target cells during vaginal SIV infection (75). In addition to the role CD4⁺ T_{RM} in CVT mucosa play in HIV susceptibility, they also act as reservoirs for HIV viral replication, which can occur during early HIV infection (76). The increased frequency of Tconv T_{RM}, and in particular Th17 T_{RM}, may not only contribute to susceptibility to primary HIV infection, but may also promote viral replication in the CX of BV+ individuals.

Activated CD8⁺ T cells are thought to be important in controlling HIV replication after initial infection (77, 78). The increased frequency of mucosal CD39⁺ CD8⁺ T cells observed in individuals with BV may also augment CVT immune dysfunction. Trends toward decreased CX

Granzyme B⁺ or T-bet⁺ CD8⁺ T cell activation (Figure 6, B and C) may relate to the increased frequency of dysfunctional CD4⁺ Tconv cells observed (Figures 4 and 5), since functional CD4⁺ Tconv cells are known to enhance CD8⁺ T cell effector function (79), and in the context of BV, CD4⁺ Tconv cells appear to be diminished in function. The interactions, or possibly lack thereof, between CD8⁺ T cells and dysfunctional Tconv cells in the context of BV, may also contribute to a dysfunctional CD8⁺ T cell response during BV and increased HIV susceptibility in BV+ individuals by inhibiting an effective CD8⁺ T cell response after exposure to HIV.

Our analysis of soluble immune factor detection in BV+ versus BV- individuals further demonstrates the diverse immune modulations that BV can promote in the CVT. We observed significant increases in several pro-inflammatory cytokines in BV+ CVT fluid samples, including IL-1 α , as previously reported by other studies (80-82). We also observed significantly increased IL-17A, IFN- γ , IL-21, IL-23, and IL-12p70 in the CVT fluid of BV+ individuals (Figure 7A), all of which can each enhance Th17-mediated inflammation (49, 66, 83-85). The activation of Th17 cells may then lead to the formation of Th17 T_{RM}, which were observed more frequently in the CX of BV+ individuals by flow cytometry (Figure 5C). Additionally, increased M-CSF in the CVT fluid of BV+ individuals (Figure 7A), produced by macrophages, has been shown to promote Th17 differentiation of CD4⁺ memory T cells (86), which may also contribute to the expanded Th17 T_{RM} pool in the CX of BV+ individuals. On the other hand, LIF and IL-17E/IL-25, observed to be increased in the CVT fluid of BV+ individuals (Figure 7A), inhibit inflammatory responses, including the inhibition of Th17-mediated immune responses (87, 88). LIF and IL-17E/IL-25 may be produced in response to the activation of Th17 cells to limit overall Th17 cell differentiation and proliferation. The reduction of TCF-1⁺ progenitor Tconv cells concurrent with the production of LIF and IL-17E/IL-25 in the CVT could limit the

differentiation of progenitor T cells to Th17 cells, which could explain why we do not observe a significant overall expansion in frequency of Th17 cells in BV+ individuals (Figure 5A).

While we observe an increased concentration of many proinflammatory cytokines in BV+ CVT fluid samples, we also observe the reduction of many T cell recruiting chemokines. Our results support previous studies that have also observed reductions of IP-10 (CXCL10) and MIG/CXCL9 in the CVT of BV+ individuals (82, 89-91). We hypothesized that RANTES (CCL5) would be increased in BV+ individuals (37) to promote the recruitment of CCR5⁺ HIV target cells to the CVT. However, our observed reduction of RANTES in the CVT fluid of BV+ individuals has also been reported in a prior study that evaluated CVT fluid collected by menstrual cup (90). While RANTES does act as a primary ligand for CCR5, which can promote the recruitment of CCR5⁺ cells, it also reduces cell-surface expression of CCR5 by binding and internalizing the CCR5 receptor (92). Increased CVT RANTES has been associated with increased HIV resistance (93), possibly by competing with HIV for CCR5 binding (94). Reduced RANTES in BV+ individuals may limit the number of CCR5⁺ HIV target cells that are recruited to the CVT. Simultaneously, this could promote stabilization of CCR5 expression on existing T cells in the CVT by reducing ligand availability and binding to internalize residually expressed CCR5, a protein preferentially observed on CVT T_{conv} cells versus circulating T_{conv} cells (Figure 2A). The relationship between CCR5 expression and CVT RANTES production in BV+ individuals is complex and how these factors impact HIV susceptibility over time will require more detailed longitudinal observational studies. The reduction of IP-10, MIG/CXCL9, RANTES, and several other proinflammatory chemokines in the CVT fluid of BV+ individuals supports the notion of impaired T cell recruitment to the CVT, which adds further evidence to dysregulated CVT immune responses in BV+ individuals. This may also contribute to adverse

health outcomes, including increased HIV susceptibility by inhibiting antiviral T cell recruitment to the CVT in BV+ individuals.

In addition to our *a priori* hypothesis of an increase in RANTES in the CVT fluid of BV+ individuals, we also hypothesized that IL-6, IL-8, and IL-1 β would each be increased in BV+ individuals based on a previous meta-analysis (37). We observed no significant difference in IL-6, IL-8, or IL-1 β in CVT samples from BV+ versus BV- individuals. However, CVT IL-1 β was trending towards an increase in BV+ (adjusted log mean difference BV+ - BV- = 0.24; p_{adj} = 0.188; Figure 7A, Supplemental Table 3). While IL-1 β is frequently described as increased in CVT fluid from BV+ individuals, this is not a universal observation (95, 96), nor is IL-6 or IL-8 uniformly increased across cohorts of BV+ individuals (97, 98). The factors that mediate altered expression of soluble immune mediators during BV have not been elucidated, highlighting the complexity and variability of BV-driven CVT immune responses.

A majority of the significant phenotypic alterations observed in this study occurred when comparing CX samples from individuals with versus without BV. This indicates that while BV is a dysbiosis of vaginal flora, the immune cells in the CX may be more greatly impacted by exposure to BV-associated organisms than the VT. One reason for this may be that the unique CX microbiota (99) may be more sensitive to changes in flora than the VT, or that the immune cells of the VT may be more resistant to BV-driven phenotypic alterations than those in the CX. The primary cellular targets for intravaginal infection of SIV in rhesus macaques are in the CX lamina propria (100), suggesting that the CX may be the primary site of sexual transmission of HIV in humans. Therefore, the more distinct immune alterations observed in the CX of BV+ individuals may be of particular concern for the increased risk of HIV transmission and

replication in BV+ individuals. Comparing immune modulations in the CX versus VT among those with or without BV warrants further exploration.

Our study has several limitations: first, we focused on characterizing diverse T cell markers and a large panel of soluble immune mediators in the context of BV infection, leaving the relationship of BV to other components of the immune response (monocytes/macrophages, granulocytes, and other immune cell types) unevaluated. One justification for limiting our evaluation to characteristics of CD3⁺ T cells is that these constitute the predominant lymphocyte population in the CVT mucosa (Figure 1A). A second limitation was that we collected 3-mm tissue biopsies that often captured low amounts of rare T cell subsets such as Tregs. We were underpowered to evaluate CX Treg phenotypes (BV+ N= 2) or VT Treg phenotypes (BV+ N= 4) (Supporting Data File). This also reduced our ability to effectively evaluate the role of other rare subsets in immune responses to BV. Third, an alternative hypothesis for adverse health outcomes associated with BV is that BV drives epithelial damage which, in the context of HIV susceptibility, reduces the effectiveness of the epithelial barrier in providing a physical barrier to protect against HIV transmission. We attempted to quantify tissue damage (data not shown) but were limited by sample quality for blinded pathology scoring. A fourth limitation to our study was that tissue biopsies were variable in size, so we cannot compare cell numbers in tissue by flow cytometry. Future studies will include weighing tissue biopsies before cryopreservation to normalize the number of cells analyzed by flow cytometry per tissue biopsy by weight. A final limitation of our analysis is that it was performed as an exploratory, hypothesis-generating analysis, and as such, used nominal p-values adjusted for confounders but not discounted for multiple comparisons (138 immune cell subsets comparisons, and 71 soluble mediator

comparisons). Thus, our findings should be confirmed through studies employing hypothesis-driven testing.

In summary, our study performed the most comprehensive evaluation of CVT tissue T cell subsets associated with BV to date. This data shows BV-driven immune alterations have the most profound impact on CX T cells and changes at this site may be most responsible for increased HIV susceptibility in BV+ individuals. While we observe increased CX Tconv activation including increased frequency of CCR5 detection on CX Tconv cells by flow cytometry, our analysis of CX and VT tissue sections by immunofluorescent microscopy shows that BV does not have a profound effect on overall CCR5⁺ HIV target cell density. Our high-parameter, high-throughput flow cytometry analysis revealed that BV drives diverse phenotypic alterations that extend beyond CD4⁺ Tconv activation in the CVT. Specifically, we identify an increased frequency of dysfunctional T cell subsets, including CD39⁺ Tconv and CD39⁺ CD8⁺ T cells, and a decreased frequency of progenitor TCF-1⁺ Tconv in the CX of BV+ individuals that could alter host response and contribute to increased HIV susceptibility by limiting the antiviral capabilities of CX T cells. Furthermore, we found the enrichment of Tconv T_{RM}, and specifically the Th17 T_{RM} subset, in the CX of BV+ individuals that may also serve as targets for HIV infection and replication. Confirmation of these findings and elaboration on molecular mechanisms may identify novel targets for immune interventions to reduce the risk of adverse health outcomes associated with BV, including increased risk of HIV infection.

Methods

Sex as a biological variable

This study focuses on cervicovaginal immune alterations in the context of BV. Therefore, samples from individuals who were identified as female at birth are exclusively used herein.

Participants, samples, and data collection

Samples and data for this analysis came from the Kinga Study which enrolled a total of 406 heterosexual Kenyan couples from Oct. 2018 through Dec. 2019 to evaluate how exposure to sexually transmitted infectious agents alters genital mucosal immune responses. Among these couples, 110 were HIV serodifferent (defined as a person living with HIV (PLWH) and their heterosexual HIV-exposed partner), with the remaining 298 couples involving partners who were both HIV-seronegative at enrollment. HIV serodifferent couples were excluded if, prior to enrollment, the PLWH had initiated antiretroviral therapy (ART) with resulting suppressed HIV viral load, or the HIV-exposed partner had initiated tenofovir-based pre-exposure prophylaxis (PrEP). ART and PrEP were provided to enrolled PLWH or HIV-exposed partners, respectively, upon enrollment.

Seven sample types were requested from all Kinga Study participants at enrollment and 6-month follow-up visits for evaluation of immune responses: CVT fluid via menstrual cup (Softcup[®]); two 3-mm VT tissue biopsies with one cryopreserved for immunofluorescent imaging, and one cryopreserved for flow cytometry; two 3-mm CX tissue biopsies processed in parallel to the VT biopsies; fractionated PBMCs; and serum were all collected (Note: genital samples and CVT fluid collection were deferred if participants were actively menstruating and participants were encouraged to return to clinic when not menstruating for collection of all

sample types at the same timepoint). Swabs for BV testing were collected prior to biopsies, and BV was assessed by Nugent score with 0-3 defined as normal flora, 4-6 as intermediate flora, and 7-10 classified as BV (101). Additional demographic, epidemiologic, clinical, and sexual behavior data (including self-reported frequency of vaginal sex, condom use, and hormonal contraceptive use) were collected at all visits. Participants were screened for HIV and HSV-2 infection. Additional details related to clinical laboratory testing are described in the Supplemental Methods.

To identify samples for the current cross-sectional analysis of the effects of BV on immune responses, we focused on 245 participants living without HIV who were identified as female at birth. These consisted of all 44 participants who may have been exposed to HIV by their enrolled heterosexual sexual partners, and 201 whose enrolled heterosexual sexual partner was without HIV and whose data could support a variety of analyses, including the analysis of BV mediated immune responses. Immunofluorescent imaging was performed on enrollment samples from a subset who were HSV-2 seronegative and had HIV-uninfected partners. All immunologic testing was performed blinded to participant exposure data.

Analysis of tissue samples by high-parameter flow cytometry

Sample collection: Biopsies were collected using baby Tischler forceps biopsies either at the lateral vaginal wall or the ectocervical os, placed in a cryovial containing fetal bovine serum at 4°C, and transported to the lab. In the lab, dimethyl sulfoxide (DMSO) was added to the cryovial to a final concentration of 10%, cryopreserved overnight at -80°C, and transferred to liquid nitrogen for long-term storage as described previously by Hughes *et al* (45). PBMCs were isolated using standard protocols.

Additional details related to sample collection, tissue processing for flow cytometry, and flow cytometry protocols and analyses are described in the Supplemental Methods and Supplemental Table 4.

Analysis of tissue samples by immunofluorescent microscopy

Fresh frozen CX and VT biopsies were sectioned and used for H&E staining and immunofluorescent (IF) staining. Tissue sections were imaged and images were acquired and analyzed. Additional details related to sample collection, tissue staining, and tissue section image analysis are described in the Supplemental Methods.

Cytokine and chemokine sample collection and processing

CVT fluid was collected via menstrual cup, and serum was collected in an SST vacutainer. Levels of cytokines and chemokines from CVT fluid and serum samples were measured using the Human Cytokine Array/Chemokine Array 71-403 Plex Panel (Eve Technologies, HD71). Additional details related to CVT fluid and serum collection and processing and the soluble mediator assay are described in the Supplemental Methods.

Statistics

For purposes of data analysis, BV+ was defined by Nugent score of 7-10. BV- was defined as normal flora by Nugent score 0-3. For flow cytometry and soluble mediator analyses, to maximize the size of our BV+ group, we used enrollment visits of participants who were BV+ at enrollment, augmented with the six-month visit from any additional participants who were BV+ at the six-month exit visit. Thus, our BV+ group consisted of one time point per individual,

but in some cases, it was enrollment while for others it was at exit. Among the remaining individuals, enrollment visits from those with normal flora at enrollment served as the BV-reference group. For immunofluorescent imaging, all specimens were from enrollment.

For both flow cytometry and soluble mediator data, results from VT and CX biopsy and CVT fluid samples were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and number of unprotected sex acts in the last thirty days, while results from PBMC samples were *a priori* adjusted for hormonal contraceptive use only. In this exploratory work, we did not adjust p-values for multiple testing. P-values less than or equal to 0.05 were considered significant. Statistical analyses were conducted using R version 4.3.3. Additional details related to statistical analyses performed for laboratory tests are described in the Supplemental Methods.

Study approval

All participants provided written informed consent using documents reviewed and approved by the University of Washington institutional review board, and the Scientific and Ethics Review Unit of the Kenya Medical Research Institute.

Data availability

The data for all analyses described in this manuscript are available in the Supporting Data File. Additional data relevant to the Kinga Study may be made available upon reasonable request to the corresponding authors.

Author contributions

FM, JBG, JLS, SCV, NP, ICT, and LW conducted the experiments. FM, ATT, JLS, AS, CM, and KKT analyzed data. MM provided reagents. JD and LKS contributed analysis methods. BHC, KN, NM, JRL, and JML designed the research study. FM, MCS, JRL, and JML wrote the first draft of the manuscript. All authors edited and approved the manuscript.

Acknowledgments

We thank all study volunteers for their participation in the Kinga Study and their willingness to provide many different samples. Additionally, we thank the members of the Kinga Study team and the Lund and Prlic labs for their helpful discussions on experimental findings and manuscript preparation throughout this process. Additionally, we thank the Fred Hutchinson Cancer Center Cellular Imaging Shared Resource, supported by the Fred Hutchinson Cancer Center Cellular Imaging Core Facility (RRID:SCR_022609) of the Fred Hutch/University of Washington/Seattle Children's Cancer Consortium (P30 CA015704), for assistance with microscopy and image analysis. SV was supported by T32 AI007509, ICT was supported by T32 AI007140, and LW was supported by T32 AI083203. See Supplemental Acknowledgments for Kinga Study consortium details.

Figure Legends

Figure 1. Bacterial vaginosis did not alter the balance of T cell frequency in the cervicovaginal tract tissues.

Flow cytometry was used to quantify the proportions of T cells within different tissue sites as indicated. (A) The frequency of CD3⁺ (CX BV- N=147, BV+ N=37; VT BV- N=144, BV+ N=40; PBMC BV- N=143, BV+ N=42) among total CD45⁺ cells. (B) The frequency of Tconv (CX BV- N=129, BV+ N=29; VT BV- N=122, BV+ N=32; PBMC BV- N=143, BV+ N=42) and (C) CD3⁺CD8⁺ (CX BV- N=141, BV+ N=33; VT BV- N=128, BV+ N=36; PBMC BV- N=143, BV+ N=42) among the total CD45⁺ cells. (D) The frequency of Tconv (CX BV- N=129, BV+ N=29; VT BV- N=122, BV+ N=32; PBMC BV- N=143, BV+ N=42) and (E) CD8⁺ (CX BV- N=141, BV+ N=33; VT BV- N=130, BV+ N=36; PBMC BV- N=143, BV+ N=42) among the total CD3⁺ cells. Adjusted rank regression analysis was performed to compare frequencies in each tissue between BV- and BV+ individuals. PBMC comparisons were *a priori* adjusted for hormonal contraceptive use, and CX and VT comparisons were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and semen exposure to reduce the effects of potential confounding variables on the analysis of BV-driven T cell alterations. Adjusted p-value displayed for all $p_{\text{adj}} \leq 0.10$; all other comparisons labeled as “ns” for non-significant. Each dot represents a measurement from an individual sample. Each horizontal bar indicates the median for its respective group. Additional statistical information for each comparison is provided in Supplemental Table 1.

Figure 2. Conventional CD4⁺ T cells displayed increased markers of activation and tissue residency in the cervix of individuals with bacterial vaginosis.

Flow cytometry was used to examine Tconv cell phenotypes within different tissue sites as indicated. (A) The frequency of CCR5⁺, (B) HLA-DR⁺, and (C) CD69⁺CD103⁺ among total Tconv cells (CX BV- N=128, BV+ N=29; VT BV- N=122, BV+ N=30; PBMC BV- N=143, BV+ N=42). Adjusted rank regression analysis was performed to compare frequencies in each tissue between BV- and BV+ individuals. PBMC comparisons were *a priori* adjusted for hormonal contraceptive use, and CX and VT comparisons were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and semen exposure to reduce the effects of potential confounding variables on the analysis of BV-driven T cell alterations. Adjusted p-value displayed for all $p_{\text{adj}} \leq 0.10$; all other comparisons labeled as “ns” for non-significant. Each dot represents a measurement from an individual sample. Each horizontal bar indicates the median for its respective group. Additional statistical information for each comparison is provided in Supplemental Table 1.

Fig 3. The total density of T cells and HIV target cells in the cervix and vagina was not altered by bacterial vaginosis.

(A) Representative H&E-stained VT tissue section imaged with brightfield microscopy at 20X magnification. (B) Representative immunofluorescent stained tissue section from the same VT biopsy as Figure 3A. DAPI stain is shown in blue, CD3 stain is shown in red, CD4 stain is shown in cyan, and CCR5 stain is shown in green. All fluorescent signals overlaid. Comparison of the density of CD3⁺ T cells, CD3⁺CD4⁺ T cells, or CD3⁺CD4⁺CD5⁺ HIV target cells in the (C) CX (BV- N=44, BV+ N=11) and (D) VT (BV- N=55, BV+ N=12). Wilcoxon rank sum test was performed for each comparison shown. Comparisons with $p > 0.05$ labeled "ns" for not significant. Each dot represents a measurement from an individual sample. Each horizontal bar

indicates the median for its respective group. Additional statistical information for each comparison is provided in Supplemental Table 2.

Figure 4. Conventional CD4⁺ T cells in the cervix showed signs of dysfunction in individuals with bacterial vaginosis.

Flow cytometry was used to examine Tconv cell phenotypes within different tissue sites as indicated. (A) The frequency of CD39⁺, (B) CD101⁺, and (C) T Cell Factor-1⁺ (TCF-1) among total Tconv cells (CX BV- N=128, BV+ N=29; VT BV- N=122, BV+ N=30; PBMC BV- N=143, BV+ N=42). Adjusted rank regression analysis was performed to compare frequencies in each tissue between BV- and BV+ individuals. PBMC comparisons were *a priori* adjusted for hormonal contraceptive use, and CX and VT comparisons were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and semen exposure to reduce the effects of potential confounding variables on the analysis of BV-driven T cell alterations. Adjusted p-value displayed for all $p_{\text{adj}} \leq 0.10$; all other comparisons labeled as “ns” for non-significant. Each dot represents a measurement from an individual sample. Each horizontal bar indicates the median for its respective group. Additional statistical information for each comparison is provided in Supplemental Table 1.

Figure 5. Th17 cells exhibited increased markers of activation and tissue residency in cervical samples from individuals with bacterial vaginosis.

Flow cytometry was used to examine Th17 phenotypes within different tissue sites as indicated. (A) The frequency of Th17 cells, defined as CD161⁺CCR6⁺ Tconv, among total Tconv cells (CX BV- N=128, BV+ N=29; VT BV- N=122, BV+ N=30; PBMC BV- N=143, BV+ N=42) (B)

HLA-DR⁺, (C) CD69⁺CD103⁺, (D) CD101⁺, and (E) CCR5⁺ frequencies among total Th17 cells (CX BV- N=42, BV+ N=9; VT BV- N=32, BV+ N=9; PBMC BV- N=143, BV+ N=42).

Adjusted rank regression analysis was performed to compare frequencies in each tissue between BV- and BV+ individuals. PBMC comparisons were *a priori* adjusted for hormonal contraceptive use, and CX and VT comparisons were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and semen exposure to reduce the effects of potential confounding variables on the analysis of BV-driven T cell alterations. Adjusted p-value displayed for all $p_{\text{adj}} \leq 0.10$; all other comparisons labeled as “ns” for non-significant. Each dot represents a measurement from an individual sample. Each horizontal bar indicates the median for its respective group. Additional statistical information for each comparison is provided in Supplemental Table 1.

Figure 6. Cervical CD8⁺ T cells displayed a dysfunctional phenotype in individuals with bacterial vaginosis.

Flow cytometry was used to examine CD8⁺ T cell phenotypes within different tissue sites as indicated. (A) The frequency of CD39⁺, (B) Granzyme B⁺, and (C) T-bet⁺ among total CD8⁺ T cells (CX BV- N=127, BV+ N=29; VT BV- N=116, BV+ N=31; PBMC BV- N=143, BV+ N=42). Adjusted rank regression analysis was performed to compare frequencies in each tissue between BV- and BV+ individuals. PBMC comparisons were *a priori* adjusted for hormonal contraceptive use, and CX and VT comparisons were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and semen exposure to reduce the effects of potential confounding variables on the analysis of BV-driven T cell alterations. Adjusted p-value displayed for all $p_{\text{adj}} \leq 0.10$; all other comparisons labeled as “ns” for non-significant. Each dot

represents a measurement from an individual sample. Each horizontal bar indicates the median for its respective group. Additional statistical information for each comparison is provided in Supplemental Table 1.

Figure 7. Bacterial vaginosis was associated with reduced chemokine concentrations and increased inflammatory cytokine concentrations in CVT fluid.

Luminex was used to quantify cytokines and chemokines from (A) CVT fluid (BV- N=134, BV+ N=36 for IL-4; BV- N=135, BV+ N=37 for IL-2; BV- N=136, BV+ N=37 for LIF and TARC; BV- N=136, BV+ N=38 for all other cytokine/chemokine comparisons) and (B) serum (BV- N=149, BV+ N=44). Adjusted estimated mean difference (BV positive – BV negative) for CVT fluid cytokines/chemokines and serum cytokines/chemokines that met the criteria for quantification of cytokine/chemokine concentrations (greater than or equal to 80% of samples were detectable). The adjusted 95% confidence interval is shown for all comparisons. Significant results when $p_{adj} \leq 0.05$ comparing BV- versus BV+ are colored in orange and non-significant differences ($p > 0.05$) comparing BV- versus BV+ are purple. Vertical dashed line at $x=0$ for reference. Serum comparisons were *a priori* adjusted for hormonal contraceptive use, and CVT fluid comparisons were *a priori* adjusted for hormonal contraceptive use, HSV-2 serology, HIV exposure, and semen exposure to reduce the effects of potential confounding variables on the analysis of BV-driven T cell alterations. Additional statistical information for each comparison is provided in Supplemental Table 3.

Tables

Table 1. Clinical and Demographic Participant Data

BV Status	Overall N = 204	Negative N = 158	Positive N = 46
Age			
<=24	66 (32.4%)	48 (30.4%)	18 (39.1%)
25-29	48 (23.5%)	36 (22.8%)	12 (26.1%)
30-34	33 (16.2%)	27 (17.1%)	6 (13.0%)
35-39	29 (14.2%)	23 (14.6%)	6 (13.0%)
40-49	27 (13.2%)	23 (14.6%)	4 (8.7%)
>=50	1 (0.5%)	1 (0.6%)	0 (0%)
HIV Exposure			
HIV exposed (HIV serodifferent couple)	36 (17.6%)	23 (14.8%)	13 (28.3%)
HIV unexposed (HIV negative, concordant couple)	168 (82.4%)	135 (85.4%)	33 (71.7%)
HSV-2 Status			
Seropositive	81 (39.9%)	56 (35.7%)	25 (54.3%)
Seronegative	114 (56.2%)	95 (60.5%)	19 (41.3%)
Indeterminate	8 (3.9%)	6 (3.8%)	2 (4.3%)
Not Tested	1	1	0
Hormonal Contraceptive Use			
Not using hormonal contraceptives and had menses in last 35 days	90 (44.8%)	64 (40.8%)	26 (59.1%)
Not using hormonal contraceptives and experiencing secondary amenorrhea	29 (14.4%)	24 (15.3%)	5 (11.4%)
Uses hormonal contraceptives	82 (40.8%)	69 (43.9%)	13 (29.5%)
Unknown	3	1	2
Number of unprotected vaginal intercourse in last 30 days	8 (3,12) ^A	8 (3,12) ^A	8 (1,12) ^A

^AMedian (IQR)

Demographic and clinical data from the 204 participants included in the flow cytometry and cytokine analyses.

References

1. Dabee S, Passmore JS, Heffron R, and Jaspan HB. The Complex Link between the Female Genital Microbiota, Genital Infections, and Inflammation. *Infect Immun*. 2021;89(5).
2. Peebles K, Velloza J, Balkus JE, McClelland RS, and Barnabas RV. High Global Burden and Costs of Bacterial Vaginosis: A Systematic Review and Meta-Analysis. *Sex Transm Dis*. 2019;46(5):304-11.
3. Sobel JD. Vaginitis. *N Engl J Med*. 1997;337(26):1896-903.
4. Allsworth JE, Lewis VA, and Peipert JF. Viral sexually transmitted infections and bacterial vaginosis: 2001-2004 National Health and Nutrition Examination Survey data. *Sex Transm Dis*. 2008;35(9):791-6.
5. Atashili J, Poole C, Ndumbe PM, Adimora AA, and Smith JS. Bacterial vaginosis and HIV acquisition: a meta-analysis of published studies. *AIDS*. 2008;22(12):1493-501.
6. Cohen CR, Duerr A, Pruithithada N, Ruggao S, Hillier S, Garcia P, et al. Bacterial vaginosis and HIV seroprevalence among female commercial sex workers in Chiang Mai, Thailand. *AIDS*. 1995;9(9):1093-7.
7. Gosmann C, Anahtar MN, Handley SA, Farcasanu M, Abu-Ali G, Bowman BA, et al. Lactobacillus-Deficient Cervicovaginal Bacterial Communities Are Associated with Increased HIV Acquisition in Young South African Women. *Immunity*. 2017;46(1):29-37.
8. Jain JP, Bristow CC, Pines HA, Harvey-Vera A, Rangel G, Staines H, et al. Factors in the HIV risk environment associated with bacterial vaginosis among HIV-negative female

- sex workers who inject drugs in the Mexico-United States border region. *BMC Public Health*. 2018;18(1):1032.
9. Massad LS, Daubert EM, Evans CT, Minkoff H, Kassaye S, Dionne-Odom J, et al. Trends in Bacterial Vaginosis Prevalence in a Cohort of U.S. Women with and at Risk for HIV. *J Womens Health (Larchmt)*. 2022;31(5):726-32.
 10. Sewankambo N, Gray RH, Wawer MJ, Paxton L, McNaim D, Wabwire-Mangen F, et al. HIV-1 infection associated with abnormal vaginal flora morphology and bacterial vaginosis. *Lancet*. 1997;350(9077):546-50.
 11. Taha TE, Hoover DR, Dallabetta GA, Kumwenda NI, Mtimavalye LA, Yang LP, et al. Bacterial vaginosis and disturbances of vaginal flora: association with increased acquisition of HIV. *AIDS*. 1998;12(13):1699-706.
 12. Hillier SL, Nugent RP, Eschenbach DA, Krohn MA, Gibbs RS, Martin DH, et al. Association between bacterial vaginosis and preterm delivery of a low-birth-weight infant. The Vaginal Infections and Prematurity Study Group. *N Engl J Med*. 1995;333(26):1737-42.
 13. Hillier SL, Martius J, Krohn M, Kiviat N, Holmes KK, and Eschenbach DA. A case-control study of chorioamnionic infection and histologic chorioamnionitis in prematurity. *N Engl J Med*. 1988;319(15):972-8.
 14. Watts DH, Krohn MA, Hillier SL, and Eschenbach DA. Bacterial vaginosis as a risk factor for post-cesarean endometritis. *Obstet Gynecol*. 1990;75(1):52-8.
 15. Torrone EA, Morrison CS, Chen PL, Kwok C, Francis SC, Hayes RJ, et al. Prevalence of sexually transmitted infections and bacterial vaginosis among women in sub-Saharan

- Africa: An individual participant data meta-analysis of 18 HIV prevention studies. *PLoS Med.* 2018;15(2):e1002511.
16. Spear GT, St John E, and Zariffard MR. Bacterial vaginosis and human immunodeficiency virus infection. *AIDS Res Ther.* 2007;4:25.
 17. St John E, Mares D, and Spear GT. Bacterial vaginosis and host immunity. *Curr HIV/AIDS Rep.* 2007;4(1):22-8.
 18. Cauci S, Culhane JF, Di Santolo M, and McCollum K. Among pregnant women with bacterial vaginosis, the hydrolytic enzymes sialidase and prolidase are positively associated with interleukin-1beta. *Am J Obstet Gynecol.* 2008;198(1):132 e1-7.
 19. Shvartsman E, Perciani CT, Richmond MEI, Russell JNH, Tough RH, Vancuren SJ, et al. Gardnerella subgroup dominant microbiomes are associated with divergent cervicovaginal immune responses in a longitudinal cohort of Kenyan women. *Front Immunol.* 2022;13:974195.
 20. Monin L, Whettlock EM, and Male V. Immune responses in the human female reproductive tract. *Immunology.* 2020;160(2):106-15.
 21. Woodward Davis AS, Vick SC, Pattacini L, Voillet V, Hughes SM, Lentz GM, et al. The human memory T cell compartment changes across tissues of the female reproductive tract. *Mucosal Immunol.* 2021;14(4):862-72.
 22. Pattacini L, Woodward Davis A, Czartoski J, Mair F, Presnell S, Hughes SM, et al. A pro-inflammatory CD8+ T-cell subset patrols the cervicovaginal tract. *Mucosal Immunol.* 2019;12(5):1118-29.

23. Kumar BV, Ma W, Miron M, Granot T, Guyer RS, Carpenter DJ, et al. Human Tissue-Resident Memory T Cells Are Defined by Core Transcriptional and Functional Signatures in Lymphoid and Mucosal Sites. *Cell Rep.* 2017;20(12):2921-34.
24. Cibrian D, and Sanchez-Madrid F. CD69: from activation marker to metabolic gatekeeper. *Eur J Immunol.* 2017;47(6):946-53.
25. Mackay LK, Braun A, Macleod BL, Collins N, Tebartz C, Bedoui S, et al. Cutting edge: CD69 interference with sphingosine-1-phosphate receptor function regulates peripheral T cell retention. *J Immunol.* 2015;194(5):2059-63.
26. Shioh LR, Rosen DB, Brdickova N, Xu Y, An J, Lanier LL, et al. CD69 acts downstream of interferon-alpha/beta to inhibit S1P1 and lymphocyte egress from lymphoid organs. *Nature.* 2006;440(7083):540-4.
27. Pham TH, Okada T, Matloubian M, Lo CG, and Cyster JG. S1P1 receptor signaling overrides retention mediated by G alpha i-coupled receptors to promote T cell egress. *Immunity.* 2008;28(1):122-33.
28. Keller HR, Ligons DL, Li C, Hwang S, Luckey MA, Prakhar P, et al. The molecular basis and cellular effects of distinct CD103 expression on CD4 and CD8 T cells. *Cell Mol Life Sci.* 2021;78(15):5789-805.
29. Sathaliyawala T, Kubota M, Yudanin N, Turner D, Camp P, Thome JJ, et al. Distribution and compartmentalization of human circulating and tissue-resident memory T cell subsets. *Immunity.* 2013;38(1):187-97.
30. Clark RA, Watanabe R, Teague JE, Schlapbach C, Tawa MC, Adams N, et al. Skin effector memory T cells do not recirculate and provide immune protection in alemtuzumab-treated CTCL patients. *Sci Transl Med.* 2012;4(117):117ra7.

31. Elliott Williams M, Hardnett FP, Sheth AN, Wein AN, Li ZT, Radzio-Basu J, et al. The menstrual cycle regulates migratory CD4 T-cell surveillance in the female reproductive tract via CCR5 signaling. *Mucosal Immunol.* 2024;17(1):41-53.
32. Schenkel JM, Fraser KA, Vezys V, and Masopust D. Sensing and alarm function of resident memory CD8(+) T cells. *Nat Immunol.* 2013;14(5):509-13.
33. Beura LK, Fares-Frederickson NJ, Steinert EM, Scott MC, Thompson EA, Fraser KA, et al. CD4(+) resident memory T cells dominate immunosurveillance and orchestrate local recall responses. *J Exp Med.* 2019;216(5):1214-29.
34. Schenkel JM, Fraser KA, Beura LK, Pauken KE, Vezys V, and Masopust D. T cell memory. Resident memory CD8 T cells trigger protective innate and adaptive immune responses. *Science.* 2014;346(6205):98-101.
35. Kumar BV, Connors TJ, and Farber DL. Human T Cell Development, Localization, and Function throughout Life. *Immunity.* 2018;48(2):202-13.
36. Rodriguez-Garcia M, Connors K, and Ghosh M. HIV Pathogenesis in the Human Female Reproductive Tract. *Curr HIV/AIDS Rep.* 2021;18(2):139-56.
37. Mitchell C, and Marrazzo J. Bacterial vaginosis and the cervicovaginal immune response. *Am J Reprod Immunol.* 2014;71(6):555-63.
38. Rebbapragada A, Howe K, Wachihhi C, Pettengell C, Sunderji S, Huibner S, et al. Bacterial vaginosis in HIV-infected women induces reversible alterations in the cervical immune environment. *J Acquir Immune Defic Syndr.* 2008;49(5):520-2.
39. Giraldo PC, de Carvalho JB, do Amaral RL, da Silveira Goncalves AK, Eleuterio J, Jr., and Guimaraes F. Identification of immune cells by flow cytometry in vaginal lavages

- from women with vulvovaginitis and normal microflora. *Am J Reprod Immunol*. 2012;67(3):198-205.
40. Kyongo JK, Jespers V, Goovaerts O, Michiels J, Menten J, Fichorova RN, et al. Searching for lower female genital tract soluble and cellular biomarkers: defining levels and predictors in a cohort of healthy Caucasian women. *PLoS One*. 2012;7(8):e43951.
 41. Anahtar MN, Byrne EH, Doherty KE, Bowman BA, Yamamoto HS, Soumillon M, et al. Cervicovaginal bacteria are a major modulator of host inflammatory responses in the female genital tract. *Immunity*. 2015;42(5):965-76.
 42. Hedges SR, Barrientes F, Desmond RA, and Schwebke JR. Local and systemic cytokine levels in relation to changes in vaginal flora. *J Infect Dis*. 2006;193(4):556-62.
 43. Byrne EH, Farcasanu M, Bloom SM, Xulu N, Xu J, Hykes BL, Jr., et al. Antigen Presenting Cells Link the Female Genital Tract Microbiome to Mucosal Inflammation, With Hormonal Contraception as an Additional Modulator of Inflammatory Signatures. *Front Cell Infect Microbiol*. 2021;11:733619.
 44. Qulu WP, Mzobe G, Mtshali A, Letsoalo MP, Osman F, San JE, et al. Metronidazole Treatment Failure and Persistent BV Lead to Increased Frequencies of Activated T- and Dendritic-Cell Subsets. *Microorganisms*. 2023;11(11).
 45. Hughes SM, Ferre AL, Yandura SE, Shetler C, Baker CAR, Calienes F, et al. Cryopreservation of human mucosal tissues. *PLoS One*. 2018;13(7):e0200653.
 46. Lund JM, Hladik F, and Prlic M. Advances and challenges in studying the tissue-resident T cell compartment in the human female reproductive tract. *Immunol Rev*. 2023;316(1):52-62.

47. Yagi H, Nomura T, Nakamura K, Yamazaki S, Kitawaki T, Hori S, et al. Crucial role of FOXP3 in the development and function of human CD25+CD4+ regulatory T cells. *Int Immunol.* 2004;16(11):1643-56.
48. Szabo SJ, Kim ST, Costa GL, Zhang X, Fathman CG, and Glimcher LH. A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell.* 2000;100(6):655-69.
49. Annunziato F, Cosmi L, Liotta F, Maggi E, and Romagnani S. The phenotype of human Th17 cells and their precursors, the cytokines that mediate their differentiation and the role of Th17 cells in inflammation. *Int Immunol.* 2008;20(11):1361-8.
50. Ssemaganda A, Nguyen HM, Nuhu F, Jahan N, Card CM, Kiazzyk S, et al. Expansion of cytotoxic tissue-resident CD8(+) T cells and CCR6(+)CD161(+) CD4(+) T cells in the nasal mucosa following mRNA COVID-19 vaccination. *Nat Commun.* 2022;13(1):3357.
51. Schey R, Dornhoff H, Baier JL, Purtak M, Opoka R, Koller AK, et al. CD101 inhibits the expansion of colitogenic T cells. *Mucosal Immunol.* 2016;9(5):1205-17.
52. Van Coillie S, Wiernicki B, and Xu J. Molecular and Cellular Functions of CTLA-4. *Adv Exp Med Biol.* 2020;1248:7-32.
53. Fang F, Yu M, Cavanagh MM, Hutter Saunders J, Qi Q, Ye Z, et al. Expression of CD39 on Activated T Cells Impairs their Survival in Older Individuals. *Cell Rep.* 2016;14(5):1218-31.
54. Zhang J, Lyu T, Cao Y, and Feng H. Role of TCF-1 in differentiation, exhaustion, and memory of CD8(+) T cells: A review. *FASEB J.* 2021;35(5):e21549.
55. Jiang Y, Li Y, and Zhu B. T-cell exhaustion in the tumor microenvironment. *Cell Death Dis.* 2015;6(6):e1792.
56. Groom JR, and Luster AD. CXCR3 in T cell function. *Exp Cell Res.* 2011;317(5):620-31.

57. Trapani JA, and Sutton VR. Granzyme B: pro-apoptotic, antiviral and antitumor functions. *Curr Opin Immunol*. 2003;15(5):533-43.
58. Sallusto F, Lenig D, Forster R, Lipp M, and Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature*. 1999;401(6754):708-12.
59. Xu L, Cao Y, Xie Z, Huang Q, Bai Q, Yang X, et al. The transcription factor TCF-1 initiates the differentiation of T(FH) cells during acute viral infection. *Nat Immunol*. 2015;16(9):991-9.
60. Choi YS, Gullicksrud JA, Xing S, Zeng Z, Shan Q, Li F, et al. LEF-1 and TCF-1 orchestrate T(FH) differentiation by regulating differentiation circuits upstream of the transcriptional repressor Bcl6. *Nat Immunol*. 2015;16(9):980-90.
61. Wu T, Shin HM, Moseman EA, Ji Y, Huang B, Harly C, et al. TCF1 Is Required for the T Follicular Helper Cell Response to Viral Infection. *Cell Rep*. 2015;12(12):2099-110.
62. Yang BH, Wang K, Wan S, Liang Y, Yuan X, Dong Y, et al. TCF1 and LEF1 Control Treg Competitive Survival and Tfr Development to Prevent Autoimmune Diseases. *Cell Rep*. 2019;27(12):3629-45 e6.
63. Yu Q, Sharma A, Oh SY, Moon HG, Hossain MZ, Salay TM, et al. T cell factor 1 initiates the T helper type 2 fate by inducing the transcription factor GATA-3 and repressing interferon-gamma. *Nat Immunol*. 2009;10(9):992-9.
64. Zhang J, He Z, Sen S, Wang F, Zhang Q, and Sun Z. TCF-1 Inhibits IL-17 Gene Expression To Restrain Th17 Immunity in a Stage-Specific Manner. *J Immunol*. 2018;200(10):3397-406.






65. Feinen B, Jerse AE, Gaffen SL, and Russell MW. Critical role of Th17 responses in a murine model of *Neisseria gonorrhoeae* genital infection. *Mucosal Immunol.* 2010;3(3):312-21.
66. Masson L, Salkinder AL, Olivier AJ, McKinnon LR, Gamielien H, Mlisana K, et al. Relationship between female genital tract infections, mucosal interleukin-17 production and local T helper type 17 cells. *Immunology.* 2015;146(4):557-67.
67. Acosta-Rodriguez EV, Rivino L, Geginat J, Jarrossay D, Gattorno M, Lanzavecchia A, et al. Surface phenotype and antigenic specificity of human interleukin 17-producing T helper memory cells. *Nat Immunol.* 2007;8(6):639-46.
68. McKinnon LR, and Kaul R. Quality and quantity: mucosal CD4+ T cells and HIV susceptibility. *Curr Opin HIV AIDS.* 2012;7(2):195-202.
69. McMichael AJ, and Rowland-Jones SL. Cellular immune responses to HIV. *Nature.* 2001;410(6831):980-7.
70. Brenna E, and McMichael AJ. The Importance of Cellular Immune Response to HIV: Implications for Antibody Production and Vaccine Design. *DNA Cell Biol.* 2022;41(1):38-42.
71. Hudson WH, Gensheimer J, Hashimoto M, Wieland A, Valanparambil RM, Li P, et al. Proliferating Transitory T Cells with an Effector-like Transcriptional Signature Emerge from PD-1(+) Stem-like CD8(+) T Cells during Chronic Infection. *Immunity.* 2019;51(6):1043-58 e4.
72. Minnie SA, Waltner OG, Zhang P, Takahashi S, Nemychenkov NS, Ensbey KS, et al. TIM-3(+) CD8 T cells with a terminally exhausted phenotype retain functional capacity in hematological malignancies. *Sci Immunol.* 2024;9(94):eadg1094.

73. Philip M, Fairchild L, Sun L, Horste EL, Camara S, Shakiba M, et al. Chromatin states define tumour-specific T cell dysfunction and reprogramming. *Nature*. 2017;545(7655):452-6.
74. Joag VR, McKinnon LR, Liu J, Kidane ST, Yudin MH, Nyanga B, et al. Identification of preferential CD4+ T-cell targets for HIV infection in the cervix. *Mucosal Immunol*. 2016;9(1):1-12.
75. Stieh DJ, Matias E, Xu H, Fought AJ, Blanchard JL, Marx PA, et al. Th17 Cells Are Preferentially Infected Very Early after Vaginal Transmission of SIV in Macaques. *Cell Host Microbe*. 2016;19(4):529-40.
76. Cantero-Perez J, Grau-Exposito J, Serra-Peinado C, Rosero DA, Luque-Ballesteros L, Astorga-Gamaza A, et al. Resident memory T cells are a cellular reservoir for HIV in the cervical mucosa. *Nat Commun*. 2019;10(1):4739.
77. Gulzar N, and Copeland KF. CD8+ T-cells: function and response to HIV infection. *Curr HIV Res*. 2004;2(1):23-37.
78. Collins DR, Gaiha GD, and Walker BD. CD8(+) T cells in HIV control, cure and prevention. *Nat Rev Immunol*. 2020;20(8):471-82.
79. Phares TW, Stohlman SA, Hwang M, Min B, Hinton DR, and Bergmann CC. CD4 T cells promote CD8 T cell immunity at the priming and effector site during viral encephalitis. *J Virol*. 2012;86(5):2416-27.
80. Ryckman KK, Williams SM, and Kalinka J. Correlations of selected vaginal cytokine levels with pregnancy-related traits in women with bacterial vaginosis and mycoplasmas. *J Reprod Immunol*. 2008;78(2):172-80.

81. Masson L, Barnabas S, Deese J, Lennard K, Dabee S, Gamiieldien H, et al. Inflammatory cytokine biomarkers of asymptomatic sexually transmitted infections and vaginal dysbiosis: a multicentre validation study. *Sex Transm Infect.* 2019;95(1):5-12.
82. Masson L, Mlisana K, Little F, Werner L, Mkhize NN, Ronacher K, et al. Defining genital tract cytokine signatures of sexually transmitted infections and bacterial vaginosis in women at high risk of HIV infection: a cross-sectional study. *Sex Transm Infect.* 2014;90(8):580-7.
83. Zielinski CE, Mele F, Aschenbrenner D, Jarrossay D, Ronchi F, Gattorno M, et al. Pathogen-induced human TH17 cells produce IFN-gamma or IL-10 and are regulated by IL-1beta. *Nature.* 2012;484(7395):514-8.
84. Khader SA, Pearl JE, Sakamoto K, Gilmartin L, Bell GK, Jelley-Gibbs DM, et al. IL-23 compensates for the absence of IL-12p70 and is essential for the IL-17 response during tuberculosis but is dispensable for protection and antigen-specific IFN-gamma responses if IL-12p70 is available. *J Immunol.* 2005;175(2):788-95.
85. Tang C, Chen S, Qian H, and Huang W. Interleukin-23: as a drug target for autoimmune inflammatory diseases. *Immunology.* 2012;135(2):112-24.
86. Foucher ED, Blanchard S, Preisser L, Descamps P, Ifrah N, Delneste Y, et al. IL-34- and M-CSF-induced macrophages switch memory T cells into Th17 cells via membrane IL-1alpha. *Eur J Immunol.* 2015;45(4):1092-102.
87. Cao W, Yang Y, Wang Z, Liu A, Fang L, Wu F, et al. Leukemia inhibitory factor inhibits T helper 17 cell differentiation and confers treatment effects of neural progenitor cell therapy in autoimmune disease. *Immunity.* 2011;35(2):273-84.


88. Kleinschek MA, Owyang AM, Joyce-Shaikh B, Langrish CL, Chen Y, Gorman DM, et al. IL-25 regulates Th17 function in autoimmune inflammation. *J Exp Med*. 2007;204(1):161-70.
89. Serebrenik J, Wang T, Hunte R, Srinivasan S, McWalters J, Tharp GK, et al. Differences in Vaginal Microbiota, Host Transcriptome, and Proteins in Women With Bacterial Vaginosis Are Associated With Metronidazole Treatment Response. *J Infect Dis*. 2021;224(12):2094-104.
90. Shannon B, Gajer P, Yi TJ, Ma B, Humphrys MS, Thomas-Pavanel J, et al. Distinct Effects of the Cervicovaginal Microbiota and Herpes Simplex Type 2 Infection on Female Genital Tract Immunology. *J Infect Dis*. 2017;215(9):1366-75.
91. Joag V, Obila O, Gajer P, Scott MC, Dizzell S, Humphrys M, et al. Impact of Standard Bacterial Vaginosis Treatment on the Genital Microbiota, Immune Milieu, and Ex Vivo Human Immunodeficiency Virus Susceptibility. *Clin Infect Dis*. 2019;68(10):1675-83.
92. Mack M, Luckow B, Nelson PJ, Cihak J, Simmons G, Clapham PR, et al. Aminooxypentane-RANTES induces CCR5 internalization but inhibits recycling: a novel inhibitory mechanism of HIV infectivity. *J Exp Med*. 1998;187(8):1215-24.
93. Iqbal SM, Ball TB, Kimani J, Kiama P, Thottingal P, Embree JE, et al. Elevated T cell counts and RANTES expression in the genital mucosa of HIV-1-resistant Kenyan commercial sex workers. *J Infect Dis*. 2005;192(5):728-38.
94. Secchi M, Grampa V, and Vangelista L. Rational CCL5 mutagenesis integration in a lactobacilli platform generates extremely potent HIV-1 blockers. *Sci Rep*. 2018;8(1):1890.

95. Losikoff P, Fichorova R, Snyder B, Rodriguez I, Cu-Uvin S, Harwell J, et al. Genital tract interleukin-8 but not interleukin-1beta or interleukin-6 concentration is associated with bacterial vaginosis and its clearance in HIV-infected and HIV-uninfected women. *Infect Dis Obstet Gynecol.* 2007;2007:92307.
96. Mtshali A, San JE, Osman F, Garrett N, Balle C, Giandhari J, et al. Temporal Changes in Vaginal Microbiota and Genital Tract Cytokines Among South African Women Treated for Bacterial Vaginosis. *Front Immunol.* 2021;12:730986.
97. Cauci S, Guaschino S, De Aloysio D, Driussi S, De Santo D, Penacchioni P, et al. Interrelationships of interleukin-8 with interleukin-1beta and neutrophils in vaginal fluid of healthy and bacterial vaginosis positive women. *Mol Hum Reprod.* 2003;9(1):53-8.
98. Weissenbacher T, Walter C, Mylonas I, Scholz C, Gingelmaier A, and Friese K. Interleukin-6, interleukin-10 and interleukin-12 in vaginal fluid from women with bacterial vaginosis. *Arch Gynecol Obstet.* 2010;281(1):77-80.
99. Teh HE, Pung CK, Arasoo VJT, and Yap PSX. A Landscape View of the Female Genital Tract Microbiome in Healthy Controls and Women With Reproductive Health Conditions Associated With Ectopic Pregnancy. *Br J Biomed Sci.* 2023;80:12098.
100. Spira AI, Marx PA, Patterson BK, Mahoney J, Koup RA, Wolinsky SM, et al. Cellular targets of infection and route of viral dissemination after an intravaginal inoculation of simian immunodeficiency virus into rhesus macaques. *J Exp Med.* 1996;183(1):215-25.
101. Nugent RP, Krohn MA, and Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol.* 1991;29(2):297-301.




• BV Negative
• BV Positive

A




B




• BV Negative
• BV Positive


C




A




B




C




D



A




B




• BV Negative
• BV Positive


C




A



B




C







• BV Negative
• BV Positive

D



E





• BV Negative
• BV Positive

