

Sample	Acrp30	F4/80	Csf1r	Cd68	TNF-a	iNOS
Adipocytes	5.17E-05 ±		1.49E-03 ±	4.77E-03 ±	1.34E-03 ±	1.42E-04 ±
Day 0	3.61E-06	ND	6.92E-04	7.57E-04	1.24E-03	1.91E-05
Adipocytes	4.19E-01 ±		4.40E-04 ±	4.51E-03 ±	8.49E-04 ±	5.13E-04 ±
Day 2	2.61E-02	ND	9.82E-05	1.03E-03	1.19E-03	7.96E-04
Adipocytes	4.38E+00 ±		1.57E-03 ±	4.99E-03 ±	3.50E-04 ±	5.62E-04 ±
Day 4	2.02E+00	ND	8.15E-04	6.44E-04	4.02E-04	3.41E-04
Adipocytes	2.77E+00 ±		8.37E-04 ±	6.72E-03 ±	3.74E-04 ±	1.50E-03 ±
Day 6	5.31E-01	ND	9.39E-05	3.25E-03	1.50E-04	1.90E-03
Adipocytes	9.20E+00 ±		1.40E-03 ±	4.60E-03 ±	2.82E-04 ±	2.20E-03 ±
Day 12	2.42E+00	ND	8.64E-04	1.21E-03	3.99E-04	4.28E-04
Adipocytes	9.02E-01 ±		9.76E-04 ±	1.46E-02 ±	8.48E-07 ±	1.27E-02 ±
Day 24	7.07E-01	ND	6.83E-04	1.41E-02	7.36E-07	1.73E-02
	8.63E-04 ±	1.07E+01±	1.32E+01 ±	1.02E+01 ±	4.32E+00 ±	4.41E-01 ±
Macrophages	1.05E-04	1.55E+00	7.28E+00	1.84E+00	2.42E+00	1.79E-01

**Table .** 3T3-L1 cells were differentiated in-vitro. Gene expression was measured using quantitative PCR at different time points and compared to gene expression in isolated adipose tissue macrophages. Data are presented as means ± SD. Not detectable