

Full unedited blots for

MESH1-mediated Coenzyme A degradation drives ferroptosis sensitivity and muscle pathology

Chao-Chieh Lin^{1,2*}, Joshua Rose^{3*}, Alexander A Mestre^{1,2,3*}, Chien-Kuang Cornelia Ding^{1,2}, Ssu-Yu Chen^{1,2}, Sze Mun Choy⁹, Kah Yong Goh⁹, Wen-Xing Lee⁹, Yanting Chen⁷, Tianai Sun^{1,2}, Jianli Wu^{1,2}, Yueqi Chen⁸, Yunju Oh⁸, Pyeonghwa Jeong⁸, Jiyong Hong^{6,8}, Michael C. Fitzgerald⁸, Guo-Fang Zhang^{4,5}, Hong-Wen Tang^{9,10†}, Pei Zhou^{3†}, Jen-Tsan Chi^{1,2†}


Corresponding author:

jentsan.chi@duke.edu


peizhou@biochem.duke.edu

hongwen.tang@duke-nus.edu.sg


Full unedited blot/gel for Figure 3E




Full unedited blot/gel for Figure 5D




Full unedited blot/gel for Figure S2A



Full unedited blot/gel for Figure S2B







Full unedited blot/gel for Figure S2H




Full unedited blot/gel for Figure S2U

siNC	+			
siMESH1	+	+	+	+
siNADK		+		+
siCOASY			+	+



Full unedited blot/gel for Figure S3J



Full unedited blot/gel for Figure S3N

