

Supplemental references

- S1. Slavotinek, A.M., Stone, E.M., Mykytyn, K., Heckenlively, J.R., Green, J.S., Heon, E., Musarella, M.A., Parfrey, P.S. and Sheffield, V.C. 2000. Mutations in MKKS cause Bardet-Biedl syndrome. *Nature Genetics* 26:15-16.
- S2. Katsanis, N., Beales, P.L., Woods, M.O., Lewis, R.A., Green, J.S., Parfrey, P.S., Ansley, S.J., Davidson, W.S. and Lupski, J.R. 2000. Mutations in MKKS cause obesity, retinal dystrophy and renal malformations associated with Bardet-Biedl syndrome. *Nature Genetics* 26:67-70.
- S3. Mykytyn, K., Braun, T., Carmi, R., Haider, N.B., Searby, C.C., Shastri, M., Beck, G., Wright, A.F., Iannaccone, A., Elbedour, K., Riise, R., Baldi, A., Raas-Rothschild, A., Gorman, S.W., Duhl, D.M., Jacobson, S.G., Casavant, T., Stone, E.M. and Sheffield, V.C. 2001. Identification of the gene that, when mutated, causes the human obesity syndrome BBS4. *Nature Genetics* 28:188-191.
- S4. Nishimura, D.Y., Searby, C.C., Carmi, R., Elbedour, K., Van Maldergem, L., Fulton, A.B., Lam, B.L., Powell, B.R., Swiderski, R.E., Bugge, K.E., Haider, N.B., Kwitek-Black, A.E., Ying, L., Duhl, D.M., Gorman, S.W., Heon, E., Iannaccone, A., Bonneau, D., Biesecker, L.G., Jacobson, S.G., Stone, E.M. and Sheffield, V.C. 2001. Positional cloning of a novel gene on chromosome 16q causing Bardet-Biedl syndrome (BBS2). *Human Molecular Genetics* 10:865-874.
- S5. Mykytyn, K., Nishimura, D.Y., Searby, C.C., Shastri, M., Yen, H.J., Beck, J.S., Braun, T., Streb, L.M., Cornier, A.S., Cox, G.F., Fulton, A.B., Carmi, R., Luleci, G., Chandrasekharappa, S.C., Collins, F.S., Jacobson, S.G., Heckenlively, J.R., Weleber, R.G., Stone, E.M. and Sheffield, V.C. 2002. Identification of the gene (BBS1) most commonly involved in Bardet-Biedl syndrome, a complex human obesity syndrome. *Nature Genetics* 31:434-438.
- S6. Chiang, A.P., Nishimura, D., Searby, C., Elbedour, K., Carmi, R., Ferguson, A.L., Secrist, J., Braun, T., Casavant, T., Stone, E.M. and Sheffield, V.C. 2004. Comparative genomic analysis identifies an ADP-ribosylation factor-like gene as the cause of Bardet-Biedl syndrome (BBS3). *American Journal Human of Genetics* 75:475-484.

- S7. Fan, Y., Esmail, M.A., Ansley, S.J., Blacque, O.E., Boroevich, K., Ross, A.J., Moore, S.J., Badano, J.L., May-Simera, H., Compton, D.S., Green, J.S., Lewis, R.A., van Haelst, M.M., Parfrey, P.S., Baillie, D.L., Beales, P.L., Katsanis, N., Davidson, W.S. and Leroux, M.R. 2004. Mutations in a member of the Ras superfamily of small GTP-binding proteins causes Bardet-Biedl syndrome. *Nature Genetics* 36:989-993.
- S8. Badano, J.L., Ansley, S.J., Leitch, C.C., Lewis, R.A., Lupski, J.R. and Katsanis, N. 2003. Identification of a novel Bardet-Biedl syndrome protein, BBS7, that shares structural features with BBS1 and BBS2. *American Journal Human of Genetics* 72:650-658.
- S9. Ansley, S.J., Badano, J.L., Blacque, O.E., Hill, J., Hoskins, B.E., Leitch, C.C., Kim, J.C., Ross, A.J., Eichers, E.R., Teslovich, T.M., Mah, A.K., Johnsen, R.C., Cavender, J.C., Lewis, R.A., Leroux, M.R., Beales, P.L. and Katsanis, N. 2003. Basal body dysfunction is a likely cause of pleiotropic Bardet-Biedl syndrome. *Nature* 425:628-633.
- S10. Li, J.B., Gerdes, J.M., Haycraft, C.J., Fan, Y., Teslovich, T.M., May-Simera, H., Li, H., Blacque, O.E., Li, L., Leitch, C.C., et al. 2004. Comparative genomics identifies a flagellar and basal body proteome that includes the BBS5 human disease gene. *Cell* 117:541-552.
- S11. Nishimura, D.Y., Swiderski, R.E., Searby, C.C., Berg, E.M., Ferguson, A.L., Hennekam, R., Merin, S., Weleber, R.G., Biesecker, L.G., Stone, E.M. and Sheffield, V.C. 2005. Comparative genomics and gene expression analysis identifies BBS9, a new Bardet-Biedl syndrome gene. *American Journal Human of Genetics* 77:1021-1033.
- S12. Stoetzel, C., Laurier, V., Davis, E.E., Muller, J., Rix, S., Badano, J.L., Leitch, C.C., Salem, N., Chouery, E., Corbani, S., Jalk, N., Vicaire, S., Sarda, P., Hamel, C., Lacombe, D., Holder, M., Odent, S., Holder, S., Brooks, A.S., Elcioglu, N.H., Silva, E.D., Rossillion, B., Sigaudy, S., de Ravel, T.J., Lewis, R.A., Leheup, B., Verloes, A., Amati-Bonneau, P., Megarbane, A., Poch, O., Bonneau, D., Beales, P.L., Mandel, J.L., Katsanis, N. and Dollfus, H. 2006. BBS10 encodes a

- vertebrate-specific chaperonin-like protein and is a major BBS locus. *Nature Genetics* 38:521-524.
- S13. White, D.R., Ganesh, A., Nishimura, D., Rattenberry, E., Ahmed, S., Smith, U.M., Pasha, S., Raeburn, S., Trembath, R.C., Rajab, A., et al. 2007. Autozygosity mapping of Bardet-Biedl syndrome to 12q21.2 and confirmation of FLJ23560 as BBS10. *Eur J Hum Genet* 15:173-178.
- S14. Chiang, A.P., Beck, J.S., Yen, H.J., Tayeh, M.K., Scheetz, T.E., Swiderski, R.E., Nishimura, D.Y., Braun, T.A., Kim, K.Y., Huang, J., Elbedour, K., Carmi, R., Slusarski, D.C., Casavant, T.L., Stone, E.M. and Sheffield, V.C. 2006. Homozygosity mapping with SNP arrays identifies TRIM32, an E3 ubiquitin ligase, as a Bardet-Biedl syndrome gene (BBS11). *Proceedings of the National Academy of Sciences USA* 103:6287-6292.
- S15. Stoetzel, C., Muller, J., Laurier, V., Davis, E.E., Zaghloul, N.A., Vicaire, S., Jacquelin, C., Plewniak, F., Leitch, C.C., Sarda, P., et al. 2007. Identification of a novel BBS gene (BBS12) highlights the major role of a vertebrate-specific branch of chaperonin-related proteins in Bardet-Biedl syndrome. *Am J Hum Genet* 80:1-11.
- S16. Stearns, T., and Botstein, D. 1988. Unlinked noncomplementation: isolation of new conditional-lethal mutations in each of the tubulin genes of *Saccharomyces cerevisiae*. *Genetics* 119:249-260.
- S17. Scholey, J.M. 2008. Intraflagellar transport motors in cilia: moving along the cell's antenna. *J Cell Biol* 180:23-29.
- S18. Zimmermann, K.W. 1898. Beiträge zur Kenntnis einiger Drüsen und Epithelien. *Arch. Mikrosk. Anat.* 52:552-706.
- S19. Satir, P. 1999. The cilium as a biological nanomachine. *FASEB Journal* 13:S235-S237.
- S20. den Hollander, A.I., Koenekoop, R.K., Yzer, S., Lopez, I., Arends, M.L., Voeselek, K.E., Zonneveld, M.N., Strom, T.M., Meitinger, T., Brunner, H.G., Hoyng, C.B., van den Born, L.I., Rohrschneider, K. and Cremers, F.P. 2006. Mutations in the CEP290 (NPHP6) gene are a frequent cause of Leber congenital amaurosis. *American Journal Human of Genetics* 79:556-561.

- S21. Sayer, J.A., Otto, E.A., O'Toole, J.F., Nurnberg, G., Kennedy, M.A., Becker, C., Hennies, H.C., Helou, J., Attanasio, M., Fausett, B.V., Utsch, B., Khanna, H., Liu, Y., Drummond, I., Kawakami, I., Kusakabe, T., Tsuda, M., Ma, L., Lee, H., Larson, R.G., Allen, S.J., Wilkinson, C.J., Nigg, E.A., Shou, C., Lillo, C., Williams, D.S., Hoppe, B., Kemper, M.J., Neuhaus, T., Parisi, M.A., Glass, I.A., Petry, M., Kispert, A., Gloy, J., Ganner, A., Walz, G., Zhu, X., Goldman, D., Nurnberg, P., Swaroop, A., Leroux, M.R. and Hildebrandt, F. 2006. The centrosomal protein nephrocystin-6 is mutated in Joubert syndrome and activates transcription factor ATF4. *Nature Genetics* 38:674-681.
- S22. Valente, E.M., Silhavy, J.L., Brancati, F., Barrano, G., Krishnaswami, S.R., Castori, M., Lancaster, M.A., Boltshauser, E., Boccone, L., Al-Gazali, L., Fazzi, E., Signorini, S., Louie, C.M., Bellacchio, E.; International Joubert Syndrome Related Disorders Study Group, Bertini, E., Dallapiccola, B. and Gleeson, J.G. 2006. Mutations in CEP290, which encodes a centrosomal protein, cause pleiotropic forms of Joubert syndrome. *Nature Genetics* 38:623-625.
- S23. Insinna, C., and Besharse, J.C. 2008. Intraflagellar transport and the sensory outer segment of vertebrate photoreceptors. *Dev Dyn* 237:1982-1992.
- S24. Tetsu, O., and McCormick, F. 1999. Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature* 398:422-426.
- S25. Hill, R.E. 2007. How to make a zone of polarizing activity: insights into limb development via the abnormality preaxial polydactyly. *Dev Growth Differ* 49:439-448.
- S26. Niswander, L., Jeffrey, S., Martin, G.R., and Tickle, C. 1994. A positive feedback loop coordinates growth and patterning in the vertebrate limb. *Nature* 371:609-612.
- S27. Krauss, S., Concordet, J.P., and Ingham, P.W. 1993. A functionally conserved homolog of the Drosophila segment polarity gene hh is expressed in tissues with polarizing activity in zebrafish embryos. *Cell* 75:1431-1444.
- S28. Wang, B., Fallon, J.F., and Beachy, P.A. 2000. Hedgehog-regulated processing of Gli3 produces an anterior/posterior repressor gradient in the developing vertebrate limb. *Cell* 100:423-434.

- S29. Litingtung, Y., Dahn, R.D., Li, Y., Fallon, J.F., and Chiang, C. 2002. Shh and Gli3 are dispensable for limb skeleton formation but regulate digit number and identity. *Nature* 418:979-983.
- S30. Masuya, H., Sagai, T., Wakana, S., Moriwaki, K., and Shiroishi, T. 1995. A duplicated zone of polarizing activity in polydactylous mouse mutants. *Genes Dev* 9:1645-1653.
- S31. Mo, R., Freer, A.M., Zinyk, D.L., Crackower, M.A., Michaud, J., Heng, H.H., Chik, K.W., Shi, X.M., Tsui, L.C., Cheng, S.H., et al. 1997. Specific and redundant functions of Gli2 and Gli3 zinc finger genes in skeletal patterning and development. *Development* 124:113-123.
- S32. Park, H.L., Bai, C., Platt, K.A., Matisse, M.P., Beeghly, A., Hui, C.C., Nakashima, M., and Joyner, A.L. 2000. Mouse Gli1 mutants are viable but have defects in SHH signaling in combination with a Gli2 mutation. *Development* 127:1593-1605.
- S33. Kang, S., Graham, J.M., Jr., Olney, A.H., and Biesecker, L.G. 1997. GLI3 frameshift mutations cause autosomal dominant Pallister-Hall syndrome. *Nat Genet* 15:266-268.
- S34. Torres, V.E., and Harris, P.C. 2006. Mechanisms of Disease: autosomal dominant and recessive polycystic kidney diseases. *Nat Clin Pract Nephrol* 2:40-55; quiz 55.